

### 3. Classification of differential equations (Notes on Diffy Qs, 0.3)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

There are many types of differential equations.

There are many types of differential equations.

Most broadly:

There are many types of differential equations.

Most broadly:

- *Ordinary differential equations* (ODE):  
Only one independent variable.

There are many types of differential equations.

Most broadly:

- *Ordinary differential equations* (ODE):  
Only one independent variable.
- *Partial differential equations* (PDE):  
Several independent variables, using partial derivatives.

**Examples:**

ODE:

## Examples:

ODE:

$$\frac{dy}{dt} = ky$$

(Exponential growth)

## Examples:

ODE:

$$\frac{dy}{dt} = ky$$

(Exponential growth)

$$\frac{dy}{dt} = k(A - y)$$

(Newton's law of cooling)



## Examples:

ODE:

$$\frac{dy}{dt} = ky$$

(Exponential growth)

$$\frac{dy}{dt} = k(A - y)$$

(Newton's law of cooling)

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = f(t)$$

(Mechanical vibrations)

## Examples:

ODE:

$$\frac{dy}{dt} = ky$$

(Exponential growth)

$$\frac{dy}{dt} = k(A - y)$$

(Newton's law of cooling)

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = f(t)$$

(Mechanical vibrations)

PDE:

## Examples:

ODE:

$$\frac{dy}{dt} = ky \quad (\text{Exponential growth})$$

$$\frac{dy}{dt} = k(A - y) \quad (\text{Newton's law of cooling})$$

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = f(t) \quad (\text{Mechanical vibrations})$$

PDE:

$$\frac{\partial y}{\partial t} + c \frac{\partial y}{\partial x} = 0 \quad (\text{Transport equation})$$

## Examples:

ODE:

$$\frac{dy}{dt} = ky \quad (\text{Exponential growth})$$

$$\frac{dy}{dt} = k(A - y) \quad (\text{Newton's law of cooling})$$

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = f(t) \quad (\text{Mechanical vibrations})$$

PDE:

$$\frac{\partial y}{\partial t} + c \frac{\partial y}{\partial x} = 0 \quad (\text{Transport equation})$$

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} \quad (\text{Heat equation})$$

## Examples:

ODE:

$$\frac{dy}{dt} = ky$$

(Exponential growth)

$$\frac{dy}{dt} = k(A - y)$$

(Newton's law of cooling)

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = f(t)$$

(Mechanical vibrations)

PDE:

$$\frac{\partial y}{\partial t} + c \frac{\partial y}{\partial x} = 0$$

(Transport equation)

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$$

(Heat equation)

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}$$

(Wave equation in 2 dimensions)

If there is more than one equation it is a *system of differential equations*.

If there is more than one equation it is a *system of differential equations*.

**Examples:**

If there is more than one equation it is a *system of differential equations*.

**Examples:**

A *system of ordinary differential equations* (system of ODE):

$$y' = x, \quad x' = y.$$



If there is more than one equation it is a *system of differential equations*.

**Examples:**

A *system of ordinary differential equations* (system of ODE):

$$y' = x, \quad x' = y.$$

Maxwell's equations for electromagnetics are a *system of partial differential equations* (system of PDE):

$$\begin{aligned} \nabla \cdot \vec{D} &= \rho, & \nabla \cdot \vec{B} &= 0, \\ \nabla \times \vec{E} &= -\frac{\partial \vec{B}}{\partial t}, & \nabla \times \vec{H} &= \vec{J} + \frac{\partial \vec{D}}{\partial t}. \end{aligned}$$

(Note: divergence  $\nabla \cdot$  and curl  $\nabla \times$  are written in partial derivatives in  $x, y, z$ .)

The highest order derivative that appears is the *order* of the equation (or system).

The highest order derivative that appears is the *order* of the equation (or system).

$$\frac{dy}{dt} = ky$$

is a first order ODE equation.

The highest order derivative that appears is the *order* of the equation (or system).

$$\frac{dy}{dt} = ky$$

is a first order ODE equation.

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} + y = \sin(x)$$

is a second order ODE equation.

The highest order derivative that appears is the *order* of the equation (or system).

$$\frac{dy}{dt} = ky$$

is a first order ODE equation.

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} + y = \sin(x)$$

is a second order ODE equation.

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

is a fourth order PDE equation.

The highest order derivative that appears is the *order* of the equation (or system).

$$\frac{dy}{dt} = ky$$

is a first order ODE equation.

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} + y = \sin(x)$$

is a second order ODE equation.

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

is a fourth order PDE equation.

**Remark:** The most common equations in physics are first and second order.

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly.

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as  $\sin$  or  $\exp$ .)



An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as sin or exp.)

An ODE of order  $n$  can be put into the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = b(x).$$

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as sin or exp.)

An ODE of order  $n$  can be put into the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = b(x).$$

E.g., linear 2<sup>nd</sup> order ODE:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as sin or exp.)

An ODE of order  $n$  can be put into the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = b(x).$$

E.g., linear 2<sup>nd</sup> order ODE:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

Note: The dependence on  $x$  need not be linear, only the dependence on  $y$ .

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as sin or exp.)

An ODE of order  $n$  can be put into the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = b(x).$$

E.g., linear 2<sup>nd</sup> order ODE:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

Note: The dependence on  $x$  need not be linear, only the dependence on  $y$ .

E.g., **nonlinear** 2<sup>nd</sup> order PDE (Burger’s equation):

$$\frac{\partial y}{\partial t} + y \frac{\partial y}{\partial x} = \nu \frac{\partial^2 y}{\partial x^2}.$$

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as sin or exp.)

An ODE of order  $n$  can be put into the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = b(x).$$

E.g., linear 2<sup>nd</sup> order ODE:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

Note: The dependence on  $x$  need not be linear, only the dependence on  $y$ .

E.g., **nonlinear** 2<sup>nd</sup> order PDE (Burger’s equation):

$$\frac{\partial y}{\partial t} + y \frac{\partial y}{\partial x} = \nu \frac{\partial^2 y}{\partial x^2}.$$

E.g., **nonlinear** 1<sup>st</sup> order ODE:

$$\frac{dx}{dt} = x^2$$

An equation is *linear* if the dependent variable(s) and their derivatives appear linearly. (“linearly”: only first powers, not multiplied together, divided, or composed with functions such as sin or exp.)

An ODE of order  $n$  can be put into the form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = b(x).$$

E.g., linear 2<sup>nd</sup> order ODE:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

Note: The dependence on  $x$  need not be linear, only the dependence on  $y$ .

E.g., **nonlinear** 2<sup>nd</sup> order PDE (Burger’s equation):

$$\frac{\partial y}{\partial t} + y \frac{\partial y}{\partial x} = \nu \frac{\partial^2 y}{\partial x^2}.$$

E.g., **nonlinear** 1<sup>st</sup> order ODE:

$$\frac{dx}{dt} = x^2$$

**Remark:** Nonlinear equations are notoriously difficult to handle.

A linear equation is *homogeneous* if all terms depend on the dependent variable.

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.



A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

**Examples:**

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

**Examples:**

Homogeneous:

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

**Examples:**

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

**Examples:**

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

### Examples:

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0$$

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

### Examples:

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0$$

Inhomogeneous:

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

### Examples:

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0$$

Inhomogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

### Examples:

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0$$

Inhomogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

$$\frac{dx}{dt} + x + t = 0$$



A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

### Examples:

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0$$

Inhomogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

$$\frac{dx}{dt} + x + t = 0$$

The inhomogeneity is often some “outside input” into the physical system.

A linear equation is *homogeneous* if all terms depend on the dependent variable. Otherwise, it is *nonhomogeneous* or *inhomogeneous*.

### Examples:

Homogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = 0$$

$$a^4 \frac{\partial^4 y}{\partial x^4} + \frac{\partial^2 y}{\partial t^2} = 0$$

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x) y = 0$$

Inhomogeneous:

$$e^x \frac{d^2 y}{dx^2} + \sin(x) \frac{dy}{dx} + x^2 y = \frac{1}{x}$$

$$\frac{dx}{dt} + x + t = 0$$

The inhomogeneity is often some “outside input” into the physical system.

We solve an inhomogeneous equation using the solution to the corresponding homogeneous equation.

A linear equation has *constant coefficients* if the coefficients are constants (except for perhaps the inhomogeneity).

A linear equation has *constant coefficients* if the coefficients are constants (except for perhaps the inhomogeneity). A general constant coefficient linear ODE:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 \frac{dy}{dx} + a_0 y = b(x),$$

$a_0, a_1, \dots, a_n$  are constants,  $b$  may depend on  $x$ .

A linear equation has *constant coefficients* if the coefficients are constants (except for perhaps the inhomogeneity). A general constant coefficient linear ODE:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 \frac{dy}{dx} + a_0 y = b(x),$$

$a_0, a_1, \dots, a_n$  are constants,  $b$  may depend on  $x$ .

Finally, an equation (or system) is *autonomous* if the equation does not depend on the independent variable at all.

A linear equation has *constant coefficients* if the coefficients are constants (except for perhaps the inhomogeneity). A general constant coefficient linear ODE:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 \frac{dy}{dx} + a_0 y = b(x),$$

$a_0, a_1, \dots, a_n$  are constants,  $b$  may depend on  $x$ .

Finally, an equation (or system) is *autonomous* if the equation does not depend on the independent variable at all.

E.g.,  $\frac{dx}{dt} = x^2$  is autonomous, but  $\frac{dx}{dt} = xt$  is not.

A linear equation has *constant coefficients* if the coefficients are constants (except for perhaps the inhomogeneity). A general constant coefficient linear ODE:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1 \frac{dy}{dx} + a_0 y = b(x),$$

$a_0, a_1, \dots, a_n$  are constants,  $b$  may depend on  $x$ .

Finally, an equation (or system) is *autonomous* if the equation does not depend on the independent variable at all.

E.g.,  $\frac{dx}{dt} = x^2$  is autonomous, but  $\frac{dx}{dt} = xt$  is not.

Autonomous equations often appear when the setup is independent of time.

**Examples:** (see if you can guess the properties before they are revealed)



**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

- $\frac{d^2x}{dt^2} + \frac{dx}{dt} + x = \sin(t)$

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

- $\frac{d^2x}{dt^2} + \frac{dx}{dt} + x = \sin(t)$

Second order, linear, inhomogeneous, constant coefficient ODE equation.

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

- $\frac{d^2x}{dt^2} + \frac{dx}{dt} + x = \sin(t)$

Second order, linear, inhomogeneous, constant coefficient ODE equation.

- $\theta'' + \sin(\theta) = 0$

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

- $\frac{d^2x}{dt^2} + \frac{dx}{dt} + x = \sin(t)$

Second order, linear, inhomogeneous, constant coefficient ODE equation.

- $\theta'' + \sin(\theta) = 0$

Second order, autonomous, nonlinear ODE equation.



**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

- $\frac{d^2x}{dt^2} + \frac{dx}{dt} + x = \sin(t)$

Second order, linear, inhomogeneous, constant coefficient ODE equation.

- $\theta'' + \sin(\theta) = 0$

Second order, autonomous, nonlinear ODE equation.

- $\frac{\partial u}{\partial t} + \frac{\partial v}{\partial x} = x^2 \quad \frac{\partial u}{\partial t} - \frac{\partial v}{\partial x} = xt$

**Examples:** (see if you can guess the properties before they are revealed)

- $\frac{dy}{dt} = ky$

First order, autonomous, linear, homogeneous, constant coefficient ODE equation.

- $\frac{dy}{dt} = k(A - y), \quad (A \neq 0)$

First order, autonomous, linear, inhomogeneous, constant coefficient ODE equation.

- $\frac{d^2x}{dt^2} + \frac{dx}{dt} + x = \sin(t)$

Second order, linear, inhomogeneous, constant coefficient ODE equation.

- $\theta'' + \sin(\theta) = 0$

Second order, autonomous, nonlinear ODE equation.

- $\frac{\partial u}{\partial t} + \frac{\partial v}{\partial x} = x^2 \quad \frac{\partial u}{\partial t} - \frac{\partial v}{\partial x} = xt$

First order, linear, inhomogeneous, constant coefficient PDE system.