

## 4. Integrals as solutions (Notes on Diffy Qs, 1.1)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C,$$

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C, \quad \text{or} \quad y(x) = \int f(x) dx + C.$$

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C, \quad \text{or} \quad y(x) = \int f(x) dx + C.$$

That's the *general solution*.

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C, \quad \text{or} \quad y(x) = \int f(x) dx + C.$$

That's the *general solution*.

**Example:** Find the general solution of  $y' = 3x^2$ .



Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C, \quad \text{or} \quad y(x) = \int f(x) dx + C.$$

That's the *general solution*.

**Example:** Find the general solution of  $y' = 3x^2$ .

Integrate:  $y = x^3 + C$ .

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C, \quad \text{or} \quad y(x) = \int f(x) dx + C.$$

That's the *general solution*.

**Example:** Find the general solution of  $y' = 3x^2$ .

Integrate:  $y = x^3 + C$ .

Check (differentiate):  $y' =$

Consider a general first order ODE:

$$\frac{dy}{dx} = f(x, y), \quad \text{or} \quad y' = f(x, y).$$

There is no single, simple procedure where you turn a crank and out pops a solution.

In the case

$$y' = f(x),$$

antidifferentiate (integrate) both sides:

$$\int y'(x) dx = \int f(x) dx + C, \quad \text{or} \quad y(x) = \int f(x) dx + C.$$

That's the *general solution*.

**Example:** Find the general solution of  $y' = 3x^2$ .

Integrate:  $y = x^3 + C$ .

Check (differentiate):  $y' = 3x^2$ . ✓

Note that  $\int f(x) dx$  means an antiderivative.

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .



Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE?

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Does it satisfy the IC?

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Does it satisfy the IC?  $y(x_0) =$

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Does it satisfy the IC?  $y(x_0) = \int_{x_0}^{x_0} f(t) dt + y_0$

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Does it satisfy the IC?  $y(x_0) = \int_{x_0}^{x_0} f(t) dt + y_0 = y_0$ . ✓

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Does it satisfy the IC?  $y(x_0) = \int_{x_0}^{x_0} f(t) dt + y_0 = y_0$ . ✓

**Example:** Solve  $y' = e^{-x^2}$ ,  $y(0) = 1$ .

Note that  $\int f(x) dx$  means an antiderivative.

An antiderivative can be computed via a definite integral:

$$y(x) = \int_{x_0}^x f(t) dt + C.$$

Even if there is no “closed form” answer, that gives you an exact formula.

Usually also an initial condition (IC)  $y(x_0) = y_0$ .

The solution to  $y' = f(x)$ ,  $y(x_0) = y_0$  is  $y(x) = \int_{x_0}^x f(t) dt + y_0$ .

Is it a solution to the DE? Yes by FTC!

Does it satisfy the IC?  $y(x_0) = \int_{x_0}^{x_0} f(t) dt + y_0 = y_0$ . ✓

**Example:** Solve  $y' = e^{-x^2}$ ,  $y(0) = 1$ .

$$y(x) = \int_0^x e^{-t^2} dt + 1.$$



So far, that's just regular Calc 1.

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

Write it in the *Leibniz notation*:  $\frac{dy}{dx} = f(y)$ .

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

Write it in the *Leibniz notation*:  $\frac{dy}{dx} = f(y)$ .

Inverse function theorem says:  $\frac{dx}{dy} = \frac{1}{f(y)}$ .

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

Write it in the *Leibniz notation*:  $\frac{dy}{dx} = f(y)$ .

Inverse function theorem says:  $\frac{dx}{dy} = \frac{1}{f(y)}$ .

Now integrate:  $x(y) = \int \frac{1}{f(y)} dy + C$ .

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

Write it in the *Leibniz notation*:  $\frac{dy}{dx} = f(y)$ .

Inverse function theorem says:  $\frac{dx}{dy} = \frac{1}{f(y)}$ .

Now integrate:  $x(y) = \int \frac{1}{f(y)} dy + C$ .

What's wrong?

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

Write it in the *Leibniz notation*:  $\frac{dy}{dx} = f(y)$ .

Inverse function theorem says:  $\frac{dx}{dy} = \frac{1}{f(y)}$ .

Now integrate:  $x(y) = \int \frac{1}{f(y)} dy + C$ .

What's wrong? We have  $x$  in terms of  $y$ , not  $y$  in terms  $x$ .

So far, that's just regular Calc 1.

How about  $y' = f(y)$  ?

Write it in the *Leibniz notation*:  $\frac{dy}{dx} = f(y)$ .

Inverse function theorem says:  $\frac{dx}{dy} = \frac{1}{f(y)}$ .

Now integrate:  $x(y) = \int \frac{1}{f(y)} dy + C$ .

What's wrong? We have  $x$  in terms of  $y$ , not  $y$  in terms  $x$ .

So solve for  $y$ !



**Example:** Let's use the method to solve  $y' = ky$  ( $k > 0$ ).

**Example:** Let's use the method to solve  $y' = ky$  ( $k > 0$ ).

Note that  $y = 0$  is a solution.

**Example:** Let's use the method to solve  $y' = ky$  ( $k > 0$ ).

Note that  $y = 0$  is a solution.

Assume  $y \neq 0$  and write  $\frac{dx}{dy} = \frac{1}{ky}$ .

**Example:** Let's use the method to solve  $y' = ky$  ( $k > 0$ ).

Note that  $y = 0$  is a solution.

Assume  $y \neq 0$  and write  $\frac{dx}{dy} = \frac{1}{ky}$ .

Integrate:  $x(y) = x = \frac{1}{k} \ln |y| + D,$

$D$  is an arbitrary constant.

**Example:** Let's use the method to solve  $y' = ky$  ( $k > 0$ ).

Note that  $y = 0$  is a solution.

Assume  $y \neq 0$  and write  $\frac{dx}{dy} = \frac{1}{ky}$ .

Integrate:  $x(y) = x = \frac{1}{k} \ln |y| + D,$

$D$  is an arbitrary constant.

Solve for  $y$ :  $|y| = e^{kx-kD} = e^{-kD} e^{kx}.$

**Example:** Let's use the method to solve  $y' = ky$  ( $k > 0$ ).

Note that  $y = 0$  is a solution.

Assume  $y \neq 0$  and write  $\frac{dx}{dy} = \frac{1}{ky}$ .

Integrate:  $x(y) = x = \frac{1}{k} \ln |y| + D,$

$D$  is an arbitrary constant.

Solve for  $y$ :  $|y| = e^{kx-kD} = e^{-kD} e^{kx}.$

If we let  $C$  be arbitrary, then  $y = Ce^{kx}$  includes all the possibilities, including  $y = 0$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .



**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$       or       $y = 0$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$       or       $y = 0$ .

IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$       or       $y = 0$ .

IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .

This solution has a singularity.

**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .


Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$  or  $y = 0$ .

IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .

This solution has a singularity.

$y$  "blows up" as  $x \rightarrow 1$ .



**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .


The general solution is  $y = \frac{1}{C - x}$  or  $y = 0$ .

IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .

This solution has a singularity.

$y$  "blows up" as  $x \rightarrow 1$ .

$y' = y^2$  seems so nice,  
yet  $y$  is so badly behaved.





**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$  or  $y = 0$ .


IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .

This solution has a singularity.

$y$  "blows up" as  $x \rightarrow 1$ .

$y' = y^2$  seems so nice,  
yet  $y$  is so badly behaved.

$y$  is only defined on  $(-\infty, 1)$



**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$  or  $y = 0$ .

IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .


This solution has a singularity.

$y$  "blows up" as  $x \rightarrow 1$ .

$y' = y^2$  seems so nice,  
yet  $y$  is so badly behaved.

$y$  is only defined on  $(-\infty, 1)$

$(1, \infty)$ , doesn't include our IC.



**Example:** Find the solution of  $y' = y^2$ ,  $y(0) = 1$ .

Note that  $y = 0$  is a solution, so we can now assume  $y \neq 0$ .

Write:  $\frac{dx}{dy} = \frac{1}{y^2}$ .      Integrate:  $x = \frac{-1}{y} + C$ .      Solve for  $y$ :  $y = \frac{1}{C - x}$ .

The general solution is  $y = \frac{1}{C - x}$  or  $y = 0$ .

IC  $y(0) = 1$  leads to  $C = 1$  or  $y = \frac{1}{1-x}$ .

This solution has a singularity.


$y$  "blows up" as  $x \rightarrow 1$ .

$y' = y^2$  seems so nice,  
yet  $y$  is so badly behaved.

$y$  is only defined on  $(-\infty, 1)$

$(1, \infty)$ , doesn't include our IC.

So the right hand side of the graph  
isn't really part of our solution.



Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled.

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .



Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0$ ,  $x = 0$ , so IC is:  $0 = x(0)$

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0$ ,  $x = 0$ , so IC is:  $0 = x(0) = 2e^{0/2} + C$

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0$ ,  $x = 0$ , so IC is:  $0 = x(0) = 2e^{0/2} + C = 2 + C$

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0, x = 0$ , so IC is:  $0 = x(0) = 2e^{0/2} + C = 2 + C \Rightarrow C = -2$ .

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0, x = 0$ , so IC is:  $0 = x(0) = 2e^{0/2} + C = 2 + C \Rightarrow C = -2$ .

Solution:  $x(t) = 2e^{t/2} - 2$ .

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0, x = 0$ , so IC is:  $0 = x(0) = 2e^{0/2} + C = 2 + C \Rightarrow C = -2$ .

Solution:  $x(t) = 2e^{t/2} - 2$ .

At 2 and 10 seconds we are at

$$x(2) = 2e^{2/2} - 2 \approx 3.44 \text{ meters,}$$

Problems solvable by integration often deal with *velocity*, *acceleration*, and *distance*.

**Example:** A car drives at a speed of  $e^{t/2}$  meters per second, where  $t$  is time in seconds. How far did the car get in 2 seconds (starting at  $t = 0$ )? How far in 10 seconds?

Denote by  $x$  the distance traveled. Equation is  $x' = e^{t/2}$ .

General solution is:  $x(t) = 2e^{t/2} + C$ .

At  $t = 0, x = 0$ , so IC is:  $0 = x(0) = 2e^{0/2} + C = 2 + C \Rightarrow C = -2$ .

Solution:  $x(t) = 2e^{t/2} - 2$ .

At 2 and 10 seconds we are at

$$x(2) = 2e^{2/2} - 2 \approx 3.44 \text{ meters}, \quad x(10) = 2e^{10/2} - 2 \approx 294 \text{ meters}.$$

**Example:** Suppose that the car accelerates at  $t^2$  m/s<sup>2</sup>.



**Example:** Suppose that the car accelerates at  $t^2 \text{ m/s}^2$ . At time  $t = 0$  the car is at the 1 meter mark and is traveling at  $10 \text{ m/s}$ .

**Example:** Suppose that the car accelerates at  $t^2 \text{ m/s}^2$ . At time  $t = 0$  the car is at the 1 meter mark and is traveling at  $10 \text{ m/s}$ . Where is the car at time  $t = 10$ ?

**Example:** Suppose that the car accelerates at  $t^2 \text{ m/s}^2$ . At time  $t = 0$  the car is at the 1 meter mark and is traveling at  $10 \text{ m/s}$ . Where is the car at time  $t = 10$ ?

Actually a second order problem.

**Example:** Suppose that the car accelerates at  $t^2 \text{ m/s}^2$ . At time  $t = 0$  the car is at the 1 meter mark and is traveling at  $10 \text{ m/s}$ . Where is the car at time  $t = 10$ ?

Actually a second order problem.

Let  $x$  be the distance. Then  $x'$  is the velocity. And  $x''$  is the acceleration.

**Example:** Suppose that the car accelerates at  $t^2$  m/s<sup>2</sup>. At time  $t = 0$  the car is at the 1 meter mark and is traveling at 10 m/s. Where is the car at time  $t = 10$ ?

Actually a second order problem.

Let  $x$  be the distance. Then  $x'$  is the velocity. And  $x''$  is the acceleration.

The full problem:  $x'' = t^2$ ,  $x(0) = 1$ ,  $x'(0) = 10$ .

**Example:** Suppose that the car accelerates at  $t^2$  m/s<sup>2</sup>. At time  $t = 0$  the car is at the 1 meter mark and is traveling at 10 m/s. Where is the car at time  $t = 10$ ?

Actually a second order problem.

Let  $x$  be the distance. Then  $x'$  is the velocity. And  $x''$  is the acceleration.

The full problem:  $x'' = t^2$ ,  $x(0) = 1$ ,  $x'(0) = 10$ .

Give  $x'$  a name, say  $v = x'$ .

**Example:** Suppose that the car accelerates at  $t^2$  m/s<sup>2</sup>. At time  $t = 0$  the car is at the 1 meter mark and is traveling at 10 m/s. Where is the car at time  $t = 10$ ?

Actually a second order problem.

Let  $x$  be the distance. Then  $x'$  is the velocity. And  $x''$  is the acceleration.

The full problem:  $x'' = t^2$ ,  $x(0) = 1$ ,  $x'(0) = 10$ .

Give  $x'$  a name, say  $v = x'$ .

Then we have the subproblem:  $v' = t^2$ ,  $v(0) = 10$ .

**Example:** Suppose that the car accelerates at  $t^2$  m/s<sup>2</sup>. At time  $t = 0$  the car is at the 1 meter mark and is traveling at 10 m/s. Where is the car at time  $t = 10$ ?

Actually a second order problem.

Let  $x$  be the distance. Then  $x'$  is the velocity. And  $x''$  is the acceleration.

The full problem:  $x'' = t^2$ ,  $x(0) = 1$ ,  $x'(0) = 10$ .

Give  $x'$  a name, say  $v = x'$ .

Then we have the subproblem:  $v' = t^2$ ,  $v(0) = 10$ .

So solve for  $v$ , then as  $x' = v$  solve for  $x$  by integration.



**Example:** Suppose that the car accelerates at  $t^2$  m/s<sup>2</sup>. At time  $t = 0$  the car is at the 1 meter mark and is traveling at 10 m/s. Where is the car at time  $t = 10$ ?

Actually a second order problem.

Let  $x$  be the distance. Then  $x'$  is the velocity. And  $x''$  is the acceleration.

The full problem:  $x'' = t^2$ ,  $x(0) = 1$ ,  $x'(0) = 10$ .

Give  $x'$  a name, say  $v = x'$ .

Then we have the subproblem:  $v' = t^2$ ,  $v(0) = 10$ .

So solve for  $v$ , then as  $x' = v$  solve for  $x$  by integration.

### Exercise

*Solve for  $v$ , and then solve for  $x$ . Find  $x(10)$  to answer the question.*