

9. Autonomous equations (Notes on Diffy Qs, 1.6)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.


$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.

Example: The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

x =temperature of coffee, t =time,

A =ambient temperature, k =constant



$$x' = 0.3(5 - x) \quad (\text{i.e., } k = 0.3, A = 5)$$

$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.


Example: The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

x =temperature of coffee, t =time,

A =ambient temperature, k =constant

The solution $x = A$ is called
an *equilibrium solution*.



$$x' = 0.3(5 - x) \quad (\text{i.e., } k = 0.3, A = 5)$$

$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.

Example: The cooling coffee problem


$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

x =temperature of coffee, t =time,

A =ambient temperature, k =constant

The solution $x = A$ is called
an *equilibrium solution*.

Equilibrium solutions happen when $f(x) = 0$.



$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.

Example: The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$


x =temperature of coffee, t =time,

A =ambient temperature, k =constant

The solution $x = A$ is called
an *equilibrium solution*.

Equilibrium solutions happen when $f(x) = 0$.

Call an x such that $f(x) = 0$ a *critical point*
($f(5) = 0$ in the plot)



$$x' = 0.3(5 - x) \quad (\text{i.e., } k = 0.3, A = 5)$$

$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.

Example: The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

x =temperature of coffee, t =time,


A =ambient temperature, k =constant

The solution $x = A$ is called
an *equilibrium solution*.

Equilibrium solutions happen when $f(x) = 0$.

Call an x such that $f(x) = 0$ a *critical point*
($f(5) = 0$ in the plot)

The critical point $x = A = 5$ is *stable*,
small changes in initial condition still lead to the solution approaching $x = 5$ as $t \rightarrow \infty$.



$\frac{dx}{dt} = f(x)$ (equation independent of time) is called an *autonomous equation*.

Example: The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

x =temperature of coffee, t =time,

A =ambient temperature, k =constant


The solution $x = A$ is called
an *equilibrium solution*.

Equilibrium solutions happen when $f(x) = 0$.

Call an x such that $f(x) = 0$ a *critical point*
($f(5) = 0$ in the plot)

The critical point $x = A = 5$ is *stable*,
small changes in initial condition still lead to the solution approaching $x = 5$ as $t \rightarrow \infty$.

A critical point that is not stable is called *unstable*.



Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.


Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Example: $x' = 0.1x(5 - x)$




Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.




Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.

$x = 0$ is unstable,




Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.
 $x = 0$ is unstable, $x = 5$ is stable.



Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.


Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.
 $x = 0$ is unstable, $x = 5$ is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$



Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)


Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.
 $x = 0$ is unstable, $x = 5$ is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$

Note: Picture not enough to decide for $x(0) < 0$.



Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.

Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)


Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.
 $x = 0$ is unstable, $x = 5$ is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$

Note: Picture not enough to decide for $x(0) < 0$. $x(t)$ may not exist for all time t .



Consider the *logistic equation*: $x' = kx(M - x)$, for some $k > 0$ and $M > 0$.


Common model for population where there is a maximum sustainable population M .
(less catastrophic than $x' = kx$)

Example: $x' = 0.1x(5 - x)$

Two critical points $x = 0$ and $x = 5$.
 $x = 0$ is unstable, $x = 5$ is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$



Note: Picture not enough to decide for $x(0) < 0$. $x(t)$ may not exist for all time t .

So to find long term behavior, $x(t)$ for very large t , we don't need to solve.

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

1) Draw the x axis (vertical).

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):


- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.


E.g.: $x' = f(x) = 0.1x(5 - x)$



To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.


E.g.: $x' = f(x) = 0.1x(5 - x)$



To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$




Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$




Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.

Graph $f(x) = 0.1x(5 - x)$,

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$




Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.
Graph $f(x) = 0.1x(5 - x)$, or plug in a few points,

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$



Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.


Graph $f(x) = 0.1x(5 - x)$, or plug in a few points,

e.g., $f(6) = -0.6 < 0$,

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$



Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.


Graph $f(x) = 0.1x(5 - x)$, or plug in a few points,

e.g., $f(6) = -0.6 < 0$, $f(1) = 0.4 > 0$,

To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$




Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.

Graph $f(x) = 0.1x(5 - x)$, or plug in a few points,
e.g., $f(6) = -0.6 < 0$, $f(1) = 0.4 > 0$, $f(-1) = -0.6 < 0$


To find $\lim_{t \rightarrow \infty} x(t)$, enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the x axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where $f(x) > 0$ and down arrow where $f(x) < 0$.

E.g.: $x' = f(x) = 0.1x(5 - x)$



Enough to note that $x = 0$ and $x = 5$ solve $0.1x(5 - x) = 0$.
Graph $f(x) = 0.1x(5 - x)$, or plug in a few points,
e.g., $f(6) = -0.6 < 0$, $f(1) = 0.4 > 0$, $f(-1) = -0.6 < 0$



Graph of $y = f(x) = 0.1x(5 - x)$

Armed with a phase diagram, easy to sketch solutions.

Classifying critical points is also easy:




Classifying critical points is also easy:



Remark: If one arrow points in and one out, sometimes called *semistable*.

Classifying critical points is also easy:



Remark: If one arrow points in and one out, sometimes called *semistable*.

Unstable points tend to be bad news:

Classifying critical points is also easy:



Remark: If one arrow points in and one out, sometimes called *semistable*.

Unstable points tend to be bad news:

Small changes in initial conditions lead to different outcomes.

Example: Consider the logistic equation with harvesting.

Example: Consider the logistic equation with harvesting.
An alien race really likes to eat humans,

Example: Consider the logistic equation with harvesting.
An alien race really likes to eat humans,
they harvest h million humans per year.

Example: Consider the logistic equation with harvesting.
An alien race really likes to eat humans,
they harvest h million humans per year.
 x =millions of humans on the planet,

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

M =limiting population when no harvesting.

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

M =limiting population when no harvesting.

$k > 0$ is a number depending on how quickly humans multiply.

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

M =limiting population when no harvesting.

$k > 0$ is a number depending on how quickly humans multiply.

Equation becomes: $x' = kx(M - x) - h$.

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

M =limiting population when no harvesting.

$k > 0$ is a number depending on how quickly humans multiply.

Equation becomes: $x' = kx(M - x) - h$.

Find critical points (solve $kx(M - x) - h = -kx^2 + kMx - h = 0$)

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

M =limiting population when no harvesting.

$k > 0$ is a number depending on how quickly humans multiply.

Equation becomes: $x' = kx(M - x) - h$.

Find critical points (solve $kx(M - x) - h = -kx^2 + kMx - h = 0$)

There are two (quadratic formula). Give them names:

$$A = \frac{kM + \sqrt{(kM)^2 - 4hk}}{2k}, \quad B = \frac{kM - \sqrt{(kM)^2 - 4hk}}{2k}.$$

Example: Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest h million humans per year.

x =millions of humans on the planet, t =time in years,

M =limiting population when no harvesting.

$k > 0$ is a number depending on how quickly humans multiply.

Equation becomes: $x' = kx(M - x) - h$.

Find critical points (solve $kx(M - x) - h = -kx^2 + kMx - h = 0$)

There are two (quadratic formula). Give them names:


$$A = \frac{kM + \sqrt{(kM)^2 - 4hk}}{2k}, \quad B = \frac{kM - \sqrt{(kM)^2 - 4hk}}{2k}.$$

3 possibilities: $A > B$, or $A = B$, or A and B both complex (i.e. no real solutions).

For example, let $M = 8$ and $k = 0.1$.

For example, let $M = 8$ and $k = 0.1$.


Harvest $h = 1$ million:



$$x' = 0.1x(8-x) - 1$$

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:




$$x' = 0.1x(8-x) - 1$$

$$A \approx 6.45,$$

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$, stable

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$, stable

$B \approx 1.55$,

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$, stable

$B \approx 1.55$, unstable

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:




$$x' = 0.1x(8 - x) - 1$$

$A \approx 6.45$, stable $B \approx 1.55$, unstable

As long as there are at least 1.55 million humans, the harvesting is sustainable.

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:



$$x' = 0.1x(8-x) - 1$$


$A \approx 6.45$, stable $B \approx 1.55$, unstable

As long as there are at least 1.55 million humans, the harvesting is sustainable.

If there is an earthquake and population drops below B ,

For example, let $M = 8$ and $k = 0.1$.

Harvest $h = 1$ million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$, stable $B \approx 1.55$, unstable

As long as there are at least 1.55 million humans, the harvesting is sustainable.


If there is an earthquake and population drops below B ,
the alien race becomes vegetarian. (not good for us either)

Now harvest $h = 1.6$:



$$x' = 0.1x(8-x) - 1.6.$$


Now harvest $h = 1.6$:



$$x' = 0.1x(8 - x) - 1.6.$$

$A = B = 4$, only one (unstable) critical point.

Now harvest $h = 1.6$:




$$x' = 0.1x(8 - x) - 1.6.$$

$A = B = 4$, only one (unstable) critical point.

If population starts above 4 million, it will tend towards 4 million.

Now harvest $h = 1.6$:



$$x' = 0.1x(8 - x) - 1.6.$$


$A = B = 4$, only one (unstable) critical point.

If population starts above 4 million, it will tend towards 4 million.

If the population drops below 4 million, humans die out.


Now harvest $h = 2$ million per year.

Now harvest $h = 2$ million per year.



$$x' = 0.1x(8-x) - 2.$$


Now harvest $h = 2$ million per year.



$$x' = 0.1x(8-x) - 2.$$

No critical points.

Now harvest $h = 2$ million per year.



$$x' = 0.1x(8-x) - 2.$$

No critical points. Population always goes to 0.