

12. Exact equations (part 2) (Notes on Diffy Qs, 1.8)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Brief review:

Brief review:

$M dx + N dy = 0$ is exact if there is a function F such that $F_x = M$ and $F_y = N$.

Brief review:

$M dx + N dy = 0$ is exact if there is a function F such that $F_x = M$ and $F_y = N$.

If such an F exists, then $M_y = N_x$.

Brief review:

$M dx + N dy = 0$ is exact if there is a function F such that $F_x = M$ and $F_y = N$.

If such an F exists, then $M_y = N_x$.

Conversely, if $M_y = N_x$, then locally such a function F exists.

Brief review:

$M dx + N dy = 0$ is exact if there is a function F such that $F_x = M$ and $F_y = N$.

If such an F exists, then $M_y = N_x$.

Conversely, if $M_y = N_x$, then locally such a function F exists.

To solve for F , integrate M in x , differentiate in y , and set equal to N .

Brief review:

$M dx + N dy = 0$ is exact if there is a function F such that $F_x = M$ and $F_y = N$.

If such an F exists, then $M_y = N_x$.

Conversely, if $M_y = N_x$, then locally such a function F exists.

To solve for F , integrate M in x , differentiate in y , and set equal to N .

(Or reverse the roles of x and y and M and N).

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$.

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y)$$

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) \quad \Rightarrow \quad 2y(x + 1) = 2xy + A'(y)$$

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) \Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y$$

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) \Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2$$

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$\begin{aligned} F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) &\Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2 \\ \Rightarrow F(x, y) &= \frac{1}{3}x^3 + xy^2 + y^2. \end{aligned}$$

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$\begin{aligned} F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) &\Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2 \\ \Rightarrow F(x, y) = \frac{1}{3}x^3 + xy^2 + y^2. &\quad \text{Now solve } F(x, y) = C. \end{aligned}$$

Example: Solve $x^2 + y^2 + 2y(x+1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x+1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$\begin{aligned} F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) &\Rightarrow 2y(x+1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2 \\ \Rightarrow F(x, y) = \frac{1}{3}x^3 + xy^2 + y^2. &\text{ Now solve } F(x, y) = C. \end{aligned}$$

$$y^2 = \frac{C - (1/3)x^3}{x+1}, \quad \text{so} \quad y = \pm \sqrt{\frac{C - (1/3)x^3}{x+1}}.$$

If $x = -1$, the explicit solution invalid.

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$\begin{aligned} F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) &\Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2 \\ \Rightarrow F(x, y) = \frac{1}{3}x^3 + xy^2 + y^2. &\text{ Now solve } F(x, y) = C. \end{aligned}$$

$$y^2 = \frac{C - (1/3)x^3}{x + 1}, \quad \text{so} \quad y = \pm \sqrt{\frac{C - (1/3)x^3}{x + 1}}.$$

If $x = -1$, the explicit solution invalid.

$x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$ has no solution there, note that term before $\frac{dy}{dx}$ is zero.

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$\begin{aligned} F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) &\Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2 \\ \Rightarrow F(x, y) = \frac{1}{3}x^3 + xy^2 + y^2. &\text{ Now solve } F(x, y) = C. \end{aligned}$$

$$y^2 = \frac{C - (1/3)x^3}{x + 1}, \quad \text{so} \quad y = \pm \sqrt{\frac{C - (1/3)x^3}{x + 1}}.$$

If $x = -1$, the explicit solution invalid.

$x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$ has no solution there, note that term before $\frac{dy}{dx}$ is zero.

$(x^2 + y^2) dx + 2y(x + 1) dy = 0$ does have a solution $x = -1$.

Example: Solve $x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$. $M = x^2 + y^2$ and $N = 2y(x + 1)$.

Exercise: Check the equation is exact.

Integrate M in x , differentiate in y and set to N :

$$\begin{aligned} F(x, y) = \frac{1}{3}x^3 + xy^2 + A(y) &\Rightarrow 2y(x + 1) = 2xy + A'(y) \Rightarrow A'(y) = 2y \Rightarrow A(y) = y^2 \\ \Rightarrow F(x, y) = \frac{1}{3}x^3 + xy^2 + y^2. &\text{ Now solve } F(x, y) = C. \end{aligned}$$

$$y^2 = \frac{C - (1/3)x^3}{x + 1}, \quad \text{so} \quad y = \pm \sqrt{\frac{C - (1/3)x^3}{x + 1}}.$$

If $x = -1$, the explicit solution invalid.

$x^2 + y^2 + 2y(x + 1)\frac{dy}{dx} = 0$ has no solution there, note that term before $\frac{dy}{dx}$ is zero.

$(x^2 + y^2)dx + 2y(x + 1)dy = 0$ does have a solution $x = -1$.

One could solve for x in terms of y for any initial condition (messy).

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x) \quad \text{and} \quad \frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x)$$

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x)$ and $\frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x) \Rightarrow$ Exact!

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x)$ and $\frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x) \Rightarrow$ Exact!

How to find the u in general?

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x) \quad \text{and} \quad \frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x) \Rightarrow \text{Exact!}$$

How to find the u in general?

$$u \text{ must satisfy } \frac{\partial}{\partial y} [uM] = u_y M + u M_y = \frac{\partial}{\partial x} [uN] = u_x N + u N_x.$$

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x) \quad \text{and} \quad \frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x) \Rightarrow \text{Exact!}$$

How to find the u in general?

$$u \text{ must satisfy } \frac{\partial}{\partial y} [uM] = u_y M + u M_y = \frac{\partial}{\partial x} [uN] = u_x N + u N_x.$$

$$\Rightarrow (M_y - N_x)u = u_x N - u_y M.$$

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x) \quad \text{and} \quad \frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x) \Rightarrow \text{Exact!}$$

How to find the u in general?

$$u \text{ must satisfy } \frac{\partial}{\partial y} [uM] = u_y M + u M_y = \frac{\partial}{\partial x} [uN] = u_x N + u N_x.$$

$$\Rightarrow (M_y - N_x)u = u_x N - u_y M.$$

OK, that's another differential equation, how does that help?

Integrating factors:

Sometimes $M dx + N dy = 0$ is not exact, but $uM dx + uN dy = 0$ is.

The linear $\frac{dy}{dx} + p(x)y = f(x)$, or $(p(x)y - f(x)) dx + dy = 0$ is an example:

Let $r(x) = e^{\int p(x) dx}$ and multiply $(r(x)p(x)y - r(x)f(x)) dx + r(x) dy = 0$.

$$\frac{\partial}{\partial y} [r(x)p(x)y - r(x)f(x)] = r(x)p(x) \quad \text{and} \quad \frac{\partial}{\partial x} [r(x)] = r'(x) = r(x)p(x) \Rightarrow \text{Exact!}$$

How to find the u in general?

$$u \text{ must satisfy } \frac{\partial}{\partial y} [uM] = u_y M + u M_y = \frac{\partial}{\partial x} [uN] = u_x N + u N_x.$$

$$\Rightarrow (M_y - N_x)u = u_x N - u_y M.$$

OK, that's another differential equation, how does that help?

One strategy: Look for u that is a function of x alone, or y alone.

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M$$

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \quad \Rightarrow \quad (M_y - N_x)u = u' N$$

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \quad \Rightarrow \quad (M_y - N_x)u = u' N \quad \Rightarrow \quad \frac{M_y - N_x}{N} u = u'$$

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \quad \Rightarrow \quad (M_y - N_x)u = u' N \quad \Rightarrow \quad \frac{M_y - N_x}{N} u = u'$$

If $\frac{M_y - N_x}{N}$ is a function of x alone, then $u' - \frac{M_y - N_x}{N} u = 0$ is a linear equation!

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \Rightarrow (M_y - N_x)u = u' N \Rightarrow \frac{M_y - N_x}{N} u = u'$$

If $\frac{M_y - N_x}{N}$ is a function of x alone, then $u' - \frac{M_y - N_x}{N} u = 0$ is a linear equation!

$$\text{Let } P(x) = \frac{M_y - N_x}{N} \Rightarrow u(x) = e^{\int P(x) dx}$$

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \Rightarrow (M_y - N_x)u = u' N \Rightarrow \frac{M_y - N_x}{N} u = u'$$

If $\frac{M_y - N_x}{N}$ is a function of x alone, then $u' - \frac{M_y - N_x}{N} u = 0$ is a linear equation!

$$\text{Let } P(x) = \frac{M_y - N_x}{N} \Rightarrow u(x) = e^{\int P(x) dx}$$

Similarly if $u = u(y)$.

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \Rightarrow (M_y - N_x)u = u' N \Rightarrow \frac{M_y - N_x}{N} u = u'$$

If $\frac{M_y - N_x}{N}$ is a function of x alone, then $u' - \frac{M_y - N_x}{N} u = 0$ is a linear equation!

$$\text{Let } P(x) = \frac{M_y - N_x}{N} \Rightarrow u(x) = e^{\int P(x) dx}$$

$$\text{Similarly if } u = u(y). \Rightarrow \frac{M_y - N_x}{M} u = -u'$$

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \Rightarrow (M_y - N_x)u = u' N \Rightarrow \frac{M_y - N_x}{N} u = u'$$

If $\frac{M_y - N_x}{N}$ is a function of x alone, then $u' - \frac{M_y - N_x}{N} u = 0$ is a linear equation!

$$\text{Let } P(x) = \frac{M_y - N_x}{N} \Rightarrow u(x) = e^{\int P(x) dx}$$

$$\text{Similarly if } u = u(y). \Rightarrow \frac{M_y - N_x}{M} u = -u'$$

If $\frac{M_y - N_x}{M}$ is a function of y alone, then

Let's look for $u(x)$. Then $u_x = u'$ and $u_y = 0$.

$$(M_y - N_x)u = u_x N - u_y M \Rightarrow (M_y - N_x)u = u' N \Rightarrow \frac{M_y - N_x}{N} u = u'$$

If $\frac{M_y - N_x}{N}$ is a function of x alone, then $u' - \frac{M_y - N_x}{N} u = 0$ is a linear equation!

$$\text{Let } P(x) = \frac{M_y - N_x}{N} \Rightarrow u(x) = e^{\int P(x) dx}$$

$$\text{Similarly if } u = u(y). \Rightarrow \frac{M_y - N_x}{M} u = -u'$$

If $\frac{M_y - N_x}{M}$ is a function of y alone, then

$$\text{Let } Q(y) = \frac{M_y - N_x}{M} \Rightarrow u(y) = e^{-\int Q(y) dy}.$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1}$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \quad \Rightarrow \quad \text{Not exact!}$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \quad \Rightarrow \quad \text{Not exact!}$$

Notice $\frac{M_y - N_x}{N}$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \quad \Rightarrow \quad \text{Not exact!}$$

$$\text{Notice } \frac{M_y - N_x}{N} = \frac{2y}{x + 1} \frac{1}{2y}$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \quad \Rightarrow \quad \text{Not exact!}$$

$$\text{Notice } \frac{M_y - N_x}{N} = \frac{2y}{x + 1} \frac{1}{2y} = \frac{1}{x + 1} = P(x) \quad \text{is a function of } x \text{ alone.}$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \quad \Rightarrow \quad \text{Not exact!}$$

$$\text{Notice } \frac{M_y - N_x}{N} = \frac{2y}{x + 1} \frac{1}{2y} = \frac{1}{x + 1} = P(x) \quad \text{is a function of } x \text{ alone.}$$

$$\text{Compute } u(x) = e^{\int P(x) dx}$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \quad \Rightarrow \quad \text{Not exact!}$$

$$\text{Notice } \frac{M_y - N_x}{N} = \frac{2y}{x + 1} \frac{1}{2y} = \frac{1}{x + 1} = P(x) \quad \text{is a function of } x \text{ alone.}$$

$$\text{Compute } u(x) = e^{\int P(x) dx} = e^{\ln(x+1)} = x + 1.$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \Rightarrow \text{Not exact!}$$

$$\text{Notice } \frac{M_y - N_x}{N} = \frac{2y}{x + 1} \frac{1}{2y} = \frac{1}{x + 1} = P(x) \text{ is a function of } x \text{ alone.}$$

$$\text{Compute } u(x) = e^{\int P(x) dx} = e^{\ln(x+1)} = x + 1.$$

$$\text{The equation becomes: } x^2 + y^2 + 2y(x + 1) \frac{dy}{dx} = 0.$$

Example: Solve $\frac{x^2 + y^2}{x + 1} + 2y \frac{dy}{dx} = 0$.

$$M = \frac{x^2 + y^2}{x + 1} \text{ and } N = 2y.$$

$$M_y - N_x = \frac{2y}{x + 1} - 0 = \frac{2y}{x + 1} \neq 0 \Rightarrow \text{Not exact!}$$

$$\text{Notice } \frac{M_y - N_x}{N} = \frac{2y}{x + 1} \frac{1}{2y} = \frac{1}{x + 1} = P(x) \text{ is a function of } x \text{ alone.}$$

$$\text{Compute } u(x) = e^{\int P(x) dx} = e^{\ln(x+1)} = x + 1.$$

$$\text{The equation becomes: } x^2 + y^2 + 2y(x + 1) \frac{dy}{dx} = 0.$$

This is exact and we solved it a moment ago:

$$y = \pm \sqrt{\frac{C - (1/3)x^3}{x + 1}}$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M}$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy}$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y),$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y}$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

$$A'(y) = 1/y$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

$$A'(y) = 1/y \quad \Rightarrow \quad A(y) = \ln |y|$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

$$A'(y) = 1/y \quad \Rightarrow \quad A(y) = \ln |y| \quad \Rightarrow \quad F(x, y) = xy + \ln |y|.$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

$$A'(y) = 1/y \quad \Rightarrow \quad A(y) = \ln |y| \quad \Rightarrow \quad F(x, y) = xy + \ln |y|.$$

$$\text{Implicit solution: } xy + \ln |y| = C.$$

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

$$A'(y) = 1/y \quad \Rightarrow \quad A(y) = \ln |y| \quad \Rightarrow \quad F(x, y) = xy + \ln |y|.$$

$$\text{Implicit solution: } xy + \ln |y| = C.$$

We divided by y , so we should check $y = 0$:

Example: Solve $y^2 + (xy + 1)\frac{dy}{dx} = 0$.

$$M_y - N_x = 2y - y = y \neq 0 \quad \Rightarrow \quad \text{Not exact.}$$

$$\frac{M_y - N_x}{M} = \frac{y}{y^2} = \frac{1}{y} = Q(y) \quad \text{is a function of } y \text{ alone.}$$

$$u(y) = e^{-\int Q(y) dy} = e^{-\ln y} = \frac{1}{y}.$$

$$\text{Equation becomes} \quad y + \frac{xy + 1}{y} \frac{dy}{dx} = 0. \quad (\text{Exercise: exact})$$

$$\text{Solve:} \quad F(x, y) = xy + A(y), \quad \frac{xy + 1}{y} = x + \frac{1}{y} = x + A'(y).$$

$$A'(y) = 1/y \quad \Rightarrow \quad A(y) = \ln |y| \quad \Rightarrow \quad F(x, y) = xy + \ln |y|.$$

$$\text{Implicit solution: } xy + \ln |y| = C.$$

We divided by y , so we should check $y = 0$: $y = 0$ is a solution too.