

13. Second order linear ODEs (Notes on Diffy Qs, 2.1)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Second order linear differential equation:

$$A(x)y'' + B(x)y' + C(x)y = F(x).$$

Second order linear differential equation:

$$A(x)y'' + B(x)y' + C(x)y = F(x).$$

Or

$$y'' + p(x)y' + q(x)y = f(x).$$

Second order linear differential equation:

$$A(x)y'' + B(x)y' + C(x)y = F(x).$$

Or

$$y'' + p(x)y' + q(x)y = f(x).$$

Linear means that the equation contains no powers nor functions of y , y' , and y'' .

Second order linear differential equation:

$$A(x)y'' + B(x)y' + C(x)y = F(x).$$

Or

$$y'' + p(x)y' + q(x)y = f(x).$$

Linear means that the equation contains no powers nor functions of y , y' , and y'' .

If $f(x) = 0$, then the equation is *homogeneous*:

$$y'' + p(x)y' + q(x)y = 0.$$

Second order linear differential equation:

$$A(x)y'' + B(x)y' + C(x)y = F(x).$$

Or

$$y'' + p(x)y' + q(x)y = f(x).$$

Linear means that the equation contains no powers nor functions of y , y' , and y'' .

If $f(x) = 0$, then the equation is *homogeneous*:

$$y'' + p(x)y' + q(x)y = 0.$$

Examples:

$$y'' + k^2y = 0$$

Two solutions are: $y_1 = \cos(kx)$, $y_2 = \sin(kx)$.

$$y'' - k^2y = 0$$

Two solutions are: $y_1 = e^{kx}$, $y_2 = e^{-kx}$.

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$.

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

We call $C_1y_1 + C_2y_2$ a *linear combination* of y_1 and y_2 .

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

We call $C_1y_1 + C_2y_2$ a *linear combination* of y_1 and y_2 .

Proof: Let $y = C_1y_1 + C_2y_2$.

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

We call $C_1y_1 + C_2y_2$ a *linear combination* of y_1 and y_2 .

Proof: Let $y = C_1y_1 + C_2y_2$. Then

$$y'' + py' + qy = (C_1y_1 + C_2y_2)'' + p(C_1y_1 + C_2y_2)' + q(C_1y_1 + C_2y_2)$$

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

We call $C_1y_1 + C_2y_2$ a *linear combination* of y_1 and y_2 .

Proof: Let $y = C_1y_1 + C_2y_2$. Then

$$\begin{aligned}y'' + py' + qy &= (C_1y_1 + C_2y_2)'' + p(C_1y_1 + C_2y_2)' + q(C_1y_1 + C_2y_2) \\&= C_1y_1'' + C_2y_2'' + C_1py_1' + C_2py_2' + C_1qy_1 + C_2qy_2\end{aligned}$$

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

We call $C_1y_1 + C_2y_2$ a *linear combination* of y_1 and y_2 .

Proof: Let $y = C_1y_1 + C_2y_2$. Then

$$\begin{aligned}y'' + py' + qy &= (C_1y_1 + C_2y_2)'' + p(C_1y_1 + C_2y_2)' + q(C_1y_1 + C_2y_2) \\&= C_1y_1'' + C_2y_2'' + C_1py_1' + C_2py_2' + C_1qy_1 + C_2qy_2 \\&= C_1(y_1'' + py_1' + qy_1) + C_2(y_2'' + py_2' + qy_2)\end{aligned}$$

Theorem (Superposition)

Suppose y_1 and y_2 are two solutions of the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then

$$y(x) = C_1y_1(x) + C_2y_2(x)$$

also solves the equation for arbitrary constants C_1 and C_2 .

We call $C_1y_1 + C_2y_2$ a *linear combination* of y_1 and y_2 .

Proof: Let $y = C_1y_1 + C_2y_2$. Then

$$\begin{aligned}y'' + py' + qy &= (C_1y_1 + C_2y_2)'' + p(C_1y_1 + C_2y_2)' + q(C_1y_1 + C_2y_2) \\&= C_1y_1'' + C_2y_2'' + C_1py_1' + C_2py_2' + C_1qy_1 + C_2qy_2 \\&= C_1(y_1'' + py_1' + qy_1) + C_2(y_2'' + py_2' + qy_2) \\&= C_1 \cdot 0 + C_2 \cdot 0 = 0.\end{aligned}$$



An *operator* is a thing that eats functions and spits out functions.

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

The operator L is *linear* if

$$L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 \quad (\text{almost like multiplying by } L)$$

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

The operator L is *linear* if

$$L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 \quad (\text{almost like multiplying by } L)$$

Nicer proof of the theorem:

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

The operator L is *linear* if

$$L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 \quad (\text{almost like multiplying by } L)$$

Nicer proof of the theorem:

Suppose that $Ly_1 = 0$ and $Ly_2 = 0$.

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

The operator L is *linear* if

$$L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 \quad (\text{almost like multiplying by } L)$$

Nicer proof of the theorem:

Suppose that $Ly_1 = 0$ and $Ly_2 = 0$. Then

$$Ly = L(C_1y_1 + C_2y_2)$$

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

The operator L is *linear* if

$$L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 \quad (\text{almost like multiplying by } L)$$

Nicer proof of the theorem:

Suppose that $Ly_1 = 0$ and $Ly_2 = 0$. Then

$$Ly = L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2$$

An *operator* is a thing that eats functions and spits out functions.

Define the operator L by

$$Ly = y'' + py' + qy.$$

The homogeneous differential equation becomes $Ly = 0$.

The operator L is *linear* if

$$L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 \quad (\text{almost like multiplying by } L)$$

Nicer proof of the theorem:

Suppose that $Ly_1 = 0$ and $Ly_2 = 0$. Then

$$Ly = L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2 = C_1 \cdot 0 + C_2 \cdot 0 = 0.$$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx)$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2}$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx}$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

And $\sinh(kx)$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

And $\sinh(kx) = \frac{e^{kx} - e^{-kx}}{2}$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

And $\sinh(kx) = \frac{e^{kx} - e^{-kx}}{2} = (1/2)e^{kx} + (-1/2)e^{-kx}$

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

And $\sinh(kx) = \frac{e^{kx} - e^{-kx}}{2} = (1/2)e^{kx} + (-1/2)e^{-kx} = (1/2)y_1 + (-1/2)y_2$ is a solution

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

And $\sinh(kx) = \frac{e^{kx} - e^{-kx}}{2} = (1/2)e^{kx} + (-1/2)e^{-kx} = (1/2)y_1 + (-1/2)y_2$ is a solution

\sinh and \cosh are sometimes more convenient than the exponential.

Example: $y'' - k^2y = 0$ has solutions $y_1 = e^{kx}$ and $y_2 = e^{-kx}$.

Recall that $\cosh t = \frac{e^t + e^{-t}}{2}$, $\sinh t = \frac{e^t - e^{-t}}{2}$

So $\cosh(kx) = \frac{e^{kx} + e^{-kx}}{2} = (1/2)e^{kx} + (1/2)e^{-kx} = (1/2)y_1 + (1/2)y_2$ is a solution

And $\sinh(kx) = \frac{e^{kx} - e^{-kx}}{2} = (1/2)e^{kx} + (-1/2)e^{-kx} = (1/2)y_1 + (-1/2)y_2$ is a solution

\sinh and \cosh are sometimes more convenient than the exponential.

Exercise: Verify that

$$\cosh 0 = 1,$$

$$\frac{d}{dt} [\cosh t] = \sinh t,$$

$$\cosh^2 t - \sinh^2 t = 1.$$

$$\sinh 0 = 0,$$

$$\frac{d}{dt} [\sinh t] = \cosh t,$$

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants.

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants. Then the equation

$$y'' + p(x)y' + q(x)y = f(x),$$

has exactly one solution $y(x)$ defined on the interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants. Then the equation

$$y'' + p(x)y' + q(x)y = f(x),$$

has exactly one solution $y(x)$ defined on the interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$

Example: $y'' + k^2y = 0$ with $y(0) = b_0$ and $y'(0) = b_1$ has the solution

$$y(x) = b_0 \cos(kx) + \frac{b_1}{k} \sin(kx).$$

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants. Then the equation

$$y'' + p(x)y' + q(x)y = f(x),$$

has exactly one solution $y(x)$ defined on the interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$

Example: $y'' + k^2y = 0$ with $y(0) = b_0$ and $y'(0) = b_1$ has the solution

$$y(x) = b_0 \cos(kx) + \frac{b_1}{k} \sin(kx).$$

Verify IC: $y(0) = b_0 \cos(0) + \frac{b_1}{k} \sin(0) = b_0.$

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants. Then the equation

$$y'' + p(x)y' + q(x)y = f(x),$$

has exactly one solution $y(x)$ defined on the interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$

Example: $y'' + k^2y = 0$ with $y(0) = b_0$ and $y'(0) = b_1$ has the solution

$$y(x) = b_0 \cos(kx) + \frac{b_1}{k} \sin(kx).$$

Verify IC: $y(0) = b_0 \cos(0) + \frac{b_1}{k} \sin(0) = b_0$.

$$y'(x) = -kb_0 \sin(kx) + b_1 \cos(kx)$$

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants. Then the equation

$$y'' + p(x)y' + q(x)y = f(x),$$

has exactly one solution $y(x)$ defined on the interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$

Example: $y'' + k^2y = 0$ with $y(0) = b_0$ and $y'(0) = b_1$ has the solution

$$y(x) = b_0 \cos(kx) + \frac{b_1}{k} \sin(kx).$$

Verify IC: $y(0) = b_0 \cos(0) + \frac{b_1}{k} \sin(0) = b_0$.

$$y'(x) = -kb_0 \sin(kx) + b_1 \cos(kx) \Rightarrow y'(0) = -kb_0 \sin(0) + b_1 \cos(0) = b_1.$$

Theorem (Existence and uniqueness)

Suppose p, q, f are continuous functions on some interval I , a is a number in I , and b_0, b_1 are constants. Then the equation

$$y'' + p(x)y' + q(x)y = f(x),$$

has exactly one solution $y(x)$ defined on the interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1.$$

Example: $y'' + k^2y = 0$ with $y(0) = b_0$ and $y'(0) = b_1$ has the solution

$$y(x) = b_0 \cos(kx) + \frac{b_1}{k} \sin(kx).$$

Verify IC: $y(0) = b_0 \cos(0) + \frac{b_1}{k} \sin(0) = b_0$.

$$y'(x) = -kb_0 \sin(kx) + b_1 \cos(kx) \Rightarrow y'(0) = -kb_0 \sin(0) + b_1 \cos(0) = b_1.$$

Similarly, $y'' - k^2y = 0$ with $y(0) = b_0$ and $y'(0) = b_1$ has the solution

$$y(x) = b_0 \cosh(kx) + \frac{b_1}{k} \sinh(kx).$$

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$.

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then every other solution is of the form

$$y = C_1y_1 + C_2y_2. \quad (\text{i.e., that's the general solution})$$

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then every other solution is of the form

$$y = C_1y_1 + C_2y_2. \quad (\text{i.e., that's the general solution})$$

Example: $y_1 = \sin x$ and $y_2 = \cos x$ solve $y'' + y = 0$.

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then every other solution is of the form

$$y = C_1y_1 + C_2y_2. \quad (\text{i.e., that's the general solution})$$

Example: $y_1 = \sin x$ and $y_2 = \cos x$ solve $y'' + y = 0$.

\sin and \cos are linearly independent:

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then every other solution is of the form

$$y = C_1y_1 + C_2y_2. \quad (\text{i.e., that's the general solution})$$

Example: $y_1 = \sin x$ and $y_2 = \cos x$ solve $y'' + y = 0$.

\sin and \cos are linearly independent: If $\sin x = A \cos x$ for some constant A , then let $x = 0$ to get $A = 0$.

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then every other solution is of the form

$$y = C_1y_1 + C_2y_2. \quad (\text{i.e., that's the general solution})$$

Example: $y_1 = \sin x$ and $y_2 = \cos x$ solve $y'' + y = 0$.

\sin and \cos are linearly independent: If $\sin x = A \cos x$ for some constant A , then let $x = 0$ to get $A = 0$. But then $\sin x = 0$ for all x , that's nonsense.

Two functions y_1 and y_2 are *linearly independent* if one is not a constant multiple of the other.

Theorem

Let p, q be continuous functions. Let y_1 and y_2 be two linearly independent solutions to the homogeneous equation $y'' + p(x)y' + q(x)y = 0$. Then every other solution is of the form

$$y = C_1 y_1 + C_2 y_2. \quad (\text{i.e., that's the general solution})$$

Example: $y_1 = \sin x$ and $y_2 = \cos x$ solve $y'' + y = 0$.

\sin and \cos are linearly independent: If $\sin x = A \cos x$ for some constant A , then let $x = 0$ to get $A = 0$. But then $\sin x = 0$ for all x , that's nonsense.

y_1 and y_2 are linearly independent and

$$y = C_1 \cos x + C_2 \sin x$$

is the general solution to $y'' + y = 0$.

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

To see y_1 and y_2 are linearly independent, suppose $y_1 = Ay_2$.

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

To see y_1 and y_2 are linearly independent, suppose $y_1 = Ay_2$.

Solve for A

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

To see y_1 and y_2 are linearly independent, suppose $y_1 = Ay_2$.

Solve for $A = \frac{y_1}{y_2} = x^3$.

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

To see y_1 and y_2 are linearly independent, suppose $y_1 = Ay_2$.

Solve for $A = \frac{y_1}{y_2} = x^3$. That's not a constant!

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

To see y_1 and y_2 are linearly independent, suppose $y_1 = Ay_2$.

Solve for $A = \frac{y_1}{y_2} = x^3$. That's not a constant! $\Rightarrow y_1$ and y_2 are linearly independent.

Example: $y'' - 2x^{-2}y = 0$ has solutions $y_1 = x^2$ and $y_2 = \frac{1}{x}$.

To see y_1 and y_2 are linearly independent, suppose $y_1 = Ay_2$.

Solve for $A = \frac{y_1}{y_2} = x^3$. That's not a constant! $\Rightarrow y_1$ and y_2 are linearly independent.

$\Rightarrow y = C_1x^2 + C_2\frac{1}{x}$ is the general solution.

If you have one solution we can find the second via the *reduction of order method*.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$0 = y_2'' + p(x)y_2' + q(x)y_2$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$0 = y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2}$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 = y_2'' + p(x)y_2' + q(x)y_2 &= \underbrace{y_1''v + 2y_1'y' + y_1v''}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)}v. \end{aligned}$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 = y_2'' + p(x)y_2' + q(x)y_2 &= \underbrace{y_1''v + 2y_1'y' + y_1v''}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \xrightarrow{0} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)}v. \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 = y_2'' + p(x)y_2' + q(x)y_2 &= \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)}v. \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v} \rightarrow 0 \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 = y_2'' + p(x)y_2' + q(x)y_2 &= \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)}v. \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \xrightarrow{0} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v}. \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

$$\Rightarrow xw' + 3w = 0$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v} \rightarrow 0 \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

$$\Rightarrow xw' + 3w = 0 \quad \Rightarrow w = Cx^{-3}$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v} \rightarrow 0 \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

$$\Rightarrow xw' + 3w = 0 \quad \Rightarrow w = Cx^{-3} \quad \Rightarrow v = \frac{-C}{2x^2}$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y_1v'}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v} \rightarrow 0 \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

$$\Rightarrow xw' + 3w = 0 \quad \Rightarrow w = Cx^{-3} \quad \Rightarrow v = \frac{-C}{2x^2} \quad \Rightarrow y_2 = y_1v = \frac{-C}{2x}$$

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y' + y_1v''}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \xrightarrow{0} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v}. \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

$$\Rightarrow xw' + 3w = 0 \quad \Rightarrow w = Cx^{-3} \quad \Rightarrow v = \frac{-C}{2x^2} \quad \Rightarrow y_2 = y_1v = \frac{-C}{2x}$$

Any C works, e.g., $C = -2$ for $y_2 = \frac{1}{x}$.

If you have one solution we can find the second via the *reduction of order method*.

Suppose y_1 solves $y'' + p(x)y' + q(x)y = 0$.

Try to find a second solution of the form $y_2(x) = y_1(x)v(x)$. Plug in:

$$\begin{aligned} 0 &= y_2'' + p(x)y_2' + q(x)y_2 = \underbrace{y_1''v + 2y_1'y' + y_1v''}_{y_2''} + \underbrace{p(x)(y_1'v + y_1v')}_{y_2'} + \underbrace{q(x)y_1v}_{y_2} \\ &= y_1v'' + (2y_1' + p(x)y_1)v' + \cancel{(y_1'' + p(x)y_1' + q(x)y_1)v} \rightarrow 0 \end{aligned}$$

So $y_1v'' + (2y_1' + p(x)y_1)v' = 0$.

Write $w = v'$ and we have a first order ODE $y_1w' + (2y_1' + p(x)y_1)w = 0$.

Solve for w , and find v by antidifferentiating.

Example: $y_1 = x$ is a solution to $y'' + x^{-1}y' - x^{-2}y = 0$, let's find y_2 .

$$\Rightarrow xw' + 3w = 0 \quad \Rightarrow w = Cx^{-3} \quad \Rightarrow v = \frac{-C}{2x^2} \quad \Rightarrow y_2 = y_1v = \frac{-C}{2x}$$

Any C works, e.g., $C = -2$ for $y_2 = \frac{1}{x}$. \Rightarrow The general solution is $y = C_1x + C_2\frac{1}{x}$.

We can even just write down a formula

$$y_2(x) = y_1(x) \int \frac{e^{-\int p(x) dx}}{(y_1(x))^2} dx$$

A useful warm-up for next time:

Exercise: For $x^2y'' - xy' = 0$, find two solutions, show that they are linearly independent and find the general solution.

A useful warm-up for next time:

Exercise: For $x^2y'' - xy' = 0$, find two solutions, show that they are linearly independent and find the general solution.

Hint: Try $y = x^r$.

A useful warm-up for next time:

Exercise: For $x^2y'' - xy' = 0$, find two solutions, show that they are linearly independent and find the general solution.

Hint: Try $y = x^r$.

Equations of the form $ax^2y'' + bxy' + cy = 0$ are called *Euler's equations* or *Cauchy–Euler equations*.