

## 5. Slope fields (Notes on Diffy Qs, 1.2)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Consider the general first order equation  $y' = f(x, y)$ .

Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,

Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ .

Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

E.g.,  $y' = xy$ .



Consider the general first order equation  $y' = f(x, y)$ .


We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

E.g.,  $y' = xy$ .

At  $(2, 1.5)$ , draw a line of slope  $xy = 2 \times 1.5 = 3$



Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.


At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

E.g.,  $y' = xy$ .

At  $(2, 1.5)$ , draw a line of slope  $xy = 2 \times 1.5 = 3$

Now do it for a grid of points.



Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .


So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

E.g.,  $y' = xy$ .

At  $(2, 1.5)$ , draw a line of slope  $xy = 2 \times 1.5 = 3$

Now do it for a grid of points.

This is called the *slope field*.



Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .


E.g.,  $y' = xy$ .

At  $(2, 1.5)$ , draw a line of slope  $xy = 2 \times 1.5 = 3$

Now do it for a grid of points.

This is called the *slope field*.

Given a specific initial condition,  
 $y(x_0) = y_0$ , we have a solution.



Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

E.g.,  $y' = xy$ .


At  $(2, 1.5)$ , draw a line of slope  $xy = 2 \times 1.5 = 3$

Now do it for a grid of points.

This is called the *slope field*.

Given a specific initial condition,  
 $y(x_0) = y_0$ , we have a solution.

E.g.,  $y(0) = 0.2$ ,  $y(0) = 0$ , and  $y(0) = -0.2$ .



Consider the general first order equation  $y' = f(x, y)$ .

We can't always solve such an equation explicitly,  
but we can always figure out the shape and behavior of the solutions.

At each  $(x, y)$  in the plane, the solution would have a slope  $y' = f(x, y)$ .

So why not draw a little line with the slope  $f(x, y)$  at the point  $(x, y)$ . And do it for all  $(x, y)$ .

E.g.,  $y' = xy$ .

At  $(2, 1.5)$ , draw a line of slope  $xy = 2 \times 1.5 = 3$


Now do it for a grid of points.

This is called the *slope field*.


Given a specific initial condition,  
 $y(x_0) = y_0$ , we have a solution.

E.g.,  $y(0) = 0.2$ ,  $y(0) = 0$ , and  $y(0) = -0.2$ .


We can tell just by the slope field that  $y(0) > 0$   
leads to very different behavior from  $y(0) < 0$ .



Here's another example:  $y' = -y$  with a few solutions given:



Here's another example:  $y' = -y$  with a few solutions given:



Note how in this case we can tell from the slope field that all solutions just tend to  $y = 0$ .



Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

(i) Does a solution *exist*?

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

The answer is usually yes to both.

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

The answer is usually yes to both.

If the answer is no, you probably do not have the right model.

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

The answer is usually yes to both.

If the answer is no, you probably do not have the right model.

**Example:** Answer to (i) can be no:

$$y' = \frac{1}{x}, \quad y(0) = 0.$$

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

The answer is usually yes to both.

If the answer is no, you probably do not have the right model.

**Example:** Answer to (i) can be no:

$$y' = \frac{1}{x}, \quad y(0) = 0.$$

(or the more harmless looking  $xy' = 1$ )

Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

The answer is usually yes to both.

If the answer is no, you probably do not have the right model.

**Example:** Answer to (i) can be no:

$$y' = \frac{1}{x}, \quad y(0) = 0.$$

(or the more harmless looking  $xy' = 1$ )

Integration yields:  $y = \ln |x| + C$ .



Consider

$$y' = f(x, y), \quad y(x_0) = y_0.$$

- (i) Does a solution *exist*?
- (ii) Is the solution *unique* (if it exists)?

The answer is usually yes to both.

If the answer is no, you probably do not have the right model.


**Example:** Answer to (i) can be no:

$$y' = \frac{1}{x}, \quad y(0) = 0.$$

(or the more harmless looking  $xy' = 1$ )

Integration yields:  $y = \ln |x| + C$ .

There is no solution at  $x = 0$ .



**Example:**

$$y' = 2\sqrt{|y|}, \quad y(0) = 0.$$

**Example:**

$$y' = 2\sqrt{|y|}, \quad y(0) = 0.$$


Note that  $y = 0$  is a solution.

**Example:**

$$y' = 2\sqrt{|y|}, \quad y(0) = 0.$$

Note that  $y = 0$  is a solution. But another solution is the function

$$y(x) = \begin{cases} x^2 & \text{if } x \geq 0, \\ -x^2 & \text{if } x < 0. \end{cases}$$



So answer to (ii) can also be false.

All hope is not lost!

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

*If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to*

$$y' = f(x, y), \quad y(x_0) = y_0,$$

*exists (for  $x$  in some interval) and is unique.*

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

*If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to*

$$y' = f(x, y), \quad y(x_0) = y_0,$$

*exists (for  $x$  in some interval) and is unique.*

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A \quad ?$



All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

*If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to*

$$y' = f(x, y), \quad y(x_0) = y_0,$$

*exists (for  $x$  in some interval) and is unique.*

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A$  ?

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A$  ?

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

If  $A \neq 0$ , then solution is  $y = \frac{1}{C-x}$ .

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A$  ?

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

If  $A \neq 0$ , then solution is  $y = \frac{1}{C-x}$ .

$$A = y(0) = \frac{1}{C-0} = 1/C$$

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A \quad ?$

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

If  $A \neq 0$ , then solution is  $y = \frac{1}{C-x}$ .

$$A = y(0) = \frac{1}{C-0} = 1/C \quad \Rightarrow \quad C = 1/A$$

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A \quad ?$

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

If  $A \neq 0$ , then solution is  $y = \frac{1}{C-x}$ .

$$A = y(0) = \frac{1}{C-0} = 1/C \quad \Rightarrow \quad C = 1/A \quad \Rightarrow \quad \text{The solution is } y = \frac{1}{1/A - x}.$$

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A \quad ?$

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

If  $A \neq 0$ , then solution is  $y = \frac{1}{C-x}$ .

$$A = y(0) = \frac{1}{C-0} = 1/C \quad \Rightarrow \quad C = 1/A \quad \Rightarrow \quad \text{The solution is } y = \frac{1}{1/A - x}.$$

If  $A = 1$ , the solution blows up when  $x = 1$ . The solution exists for  $x < 1$ .

All hope is not lost!

### Theorem (Picard's theorem on existence and uniqueness)

If  $f(x, y)$  is continuous and  $\frac{\partial f}{\partial y}$  exists and is continuous near  $(x_0, y_0)$ , then a solution to

$$y' = f(x, y), \quad y(x_0) = y_0,$$

exists (for  $x$  in some interval) and is unique.

E.g.,  $y' = 1/x, y(0) = 0$  and  $y' = 2\sqrt{|y|}, y(0) = 0$  do not satisfy the hypotheses.

**Example:** Remember  $y' = y^2, \quad y(0) = A$  ?

If  $A = 0$ , then  $y = 0$  is the solution. It exists for all  $x$ .

If  $A \neq 0$ , then solution is  $y = \frac{1}{C-x}$ .

$$A = y(0) = \frac{1}{C-0} = 1/C \quad \Rightarrow \quad C = 1/A \quad \Rightarrow \quad \text{The solution is } y = \frac{1}{1/A - x}.$$

If  $A = 1$ , the solution blows up when  $x = 1$ . The solution exists for  $x < 1$ .

If  $A = 100$ , the solution blows up when  $x = \frac{1}{100} = 0.01$ . The solution exists for  $x < 0.01$ .