

# BA: 4.1

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ .

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ . If the limit

$$L := \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad \text{exists,}$$

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ . If the limit

$$L := \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad \text{exists,}$$

then  $f$  is *differentiable* at  $c$ ,  $L$  is the *derivative* of  $f$  at  $c$ , and write  $f'(c) := L$ .

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ . If the limit

$$L := \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad \text{exists,}$$

then  $f$  is *differentiable* at  $c$ ,  $L$  is the *derivative* of  $f$  at  $c$ , and write  $f'(c) := L$ .

If  $f$  is differentiable at all  $c \in I$ , say  $f$  is *differentiable*, and obtain a function  $f': I \rightarrow \mathbb{R}$ .

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ . If the limit

$$L := \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad \text{exists,}$$

then  $f$  is *differentiable* at  $c$ ,  $L$  is the *derivative* of  $f$  at  $c$ , and write  $f'(c) := L$ .

If  $f$  is differentiable at all  $c \in I$ , say  $f$  is *differentiable*, and obtain a function  $f': I \rightarrow \mathbb{R}$ .

Other notation:  $\frac{df}{dx}$  or  $\frac{d}{dx}(f(x))$ .

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ . If the limit

$$L := \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad \text{exists,}$$

then  $f$  is *differentiable* at  $c$ ,  $L$  is the *derivative* of  $f$  at  $c$ , and write  $f'(c) := L$ .

If  $f$  is differentiable at all  $c \in I$ , say  $f$  is *differentiable*, and obtain a function  $f': I \rightarrow \mathbb{R}$ .

Other notation:  $\frac{df}{dx}$  or  $\frac{d}{dx}(f(x))$ .

The expression  $\frac{f(x) - f(c)}{x - c}$  is called the *difference quotient*.

## Definition

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  be a function, and let  $c \in I$ . If the limit

$$L := \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \quad \text{exists,}$$

then  $f$  is *differentiable* at  $c$ ,  $L$  is the *derivative* of  $f$  at  $c$ , and write  $f'(c) := L$ .

If  $f$  is differentiable at all  $c \in I$ , say  $f$  is *differentiable*, and obtain a function  $f': I \rightarrow \mathbb{R}$ .



Other notation:  $\frac{df}{dx}$  or  $\frac{d}{dx}(f(x))$ .

The expression  $\frac{f(x) - f(c)}{x - c}$  is called the *difference quotient*.


**Remark:** We allow  $I$  to be a closed interval, and we allow  $c$  to be an endpoint of  $I$ .



The graphical interpretation of the derivative:




The graphical interpretation of the derivative:



$\frac{f(x) - f(c)}{x - c}$  is the slope of the *secant line* through  $(c, f(c))$  and  $(x, f(x))$ .

The graphical interpretation of the derivative:



$\frac{f(x) - f(c)}{x - c}$  is the slope of the *secant line* through  $(c, f(c))$  and  $(x, f(x))$ .

$f'(c)$  is the limit of these slopes as  $x \rightarrow c$ .

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ .

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c}$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c}$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c}$$



**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c)$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ .

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ .

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ . For  $x \neq c$ ,

$$\frac{f(x) - f(c)}{x - c}$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ . For  $x \neq c$ ,

$$\frac{f(x) - f(c)}{x - c} = \frac{a(x - c)}{x - c}$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ . For  $x \neq c$ ,

$$\frac{f(x) - f(c)}{x - c} = \frac{a(x - c)}{x - c} = a.$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ . For  $x \neq c$ ,

$$\frac{f(x) - f(c)}{x - c} = \frac{a(x - c)}{x - c} = a.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$



**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ . For  $x \neq c$ ,

$$\frac{f(x) - f(c)}{x - c} = \frac{a(x - c)}{x - c} = a.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = \lim_{x \rightarrow c} a = a.$$

**Example:** Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be  $f(x) := x^2$  and  $c \in \mathbb{R}$ . If  $x \neq c$ , then

$$\frac{x^2 - c^2}{x - c} = \frac{(x + c)(x - c)}{x - c} = (x + c).$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{x^2 - c^2}{x - c} = \lim_{x \rightarrow c} (x + c) = 2c.$$

**Example:** Let  $f(x) := ax + b$  for  $a, b \in \mathbb{R}$ . Let  $c \in \mathbb{R}$ . For  $x \neq c$ ,

$$\frac{f(x) - f(c)}{x - c} = \frac{a(x - c)}{x - c} = a.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = \lim_{x \rightarrow c} a = a.$$

**Remark:** Every differentiable  $f$  “infinitesimally” behaves like the affine function  $ax + b$ .

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ .

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c}$$

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})}$$

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c}$$



**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c} = \lim_{x \rightarrow c} \frac{1}{\sqrt{x} + \sqrt{c}}$$

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c} = \lim_{x \rightarrow c} \frac{1}{\sqrt{x} + \sqrt{c}} = \frac{1}{2\sqrt{c}}.$$

□

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c} = \lim_{x \rightarrow c} \frac{1}{\sqrt{x} + \sqrt{c}} = \frac{1}{2\sqrt{c}}.$$

□

**Example:**  $f(x) := |x|$  is not differentiable at 0.

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c} = \lim_{x \rightarrow c} \frac{1}{\sqrt{x} + \sqrt{c}} = \frac{1}{2\sqrt{c}}.$$

□

**Example:**  $f(x) := |x|$  is not differentiable at 0.

**Proof:** When  $x > 0$ ,

$$\frac{|x| - |0|}{x - 0} = \frac{x - 0}{x - 0} = 1.$$

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c} = \lim_{x \rightarrow c} \frac{1}{\sqrt{x} + \sqrt{c}} = \frac{1}{2\sqrt{c}}.$$

□

**Example:**  $f(x) := |x|$  is not differentiable at 0.

**Proof:** When  $x > 0$ ,

$$\frac{|x| - |0|}{x - 0} = \frac{x - 0}{x - 0} = 1.$$

When  $x < 0$ ,

$$\frac{|x| - |0|}{x - 0} = \frac{-x - 0}{x - 0} = -1.$$

**Example:**  $f(x) := \sqrt{x}$  is differentiable for  $x > 0$ . **Proof:** Fix  $c > 0$  and take  $x \neq c, x > 0$ .

$$\frac{\sqrt{x} - \sqrt{c}}{x - c} = \frac{\sqrt{x} - \sqrt{c}}{(\sqrt{x} - \sqrt{c})(\sqrt{x} + \sqrt{c})} = \frac{1}{\sqrt{x} + \sqrt{c}}.$$

Therefore,

$$f'(c) = \lim_{x \rightarrow c} \frac{\sqrt{x} - \sqrt{c}}{x - c} = \lim_{x \rightarrow c} \frac{1}{\sqrt{x} + \sqrt{c}} = \frac{1}{2\sqrt{c}}.$$

□

**Example:**  $f(x) := |x|$  is not differentiable at 0.

**Proof:** When  $x > 0$ ,

$$\frac{|x| - |0|}{x - 0} = \frac{x - 0}{x - 0} = 1.$$

When  $x < 0$ ,

$$\frac{|x| - |0|}{x - 0} = \frac{-x - 0}{x - 0} = -1.$$

So the limit  $x \rightarrow 0$  does not exist.

□

A continuous function exists which is not differentiable at any point (Weierstrass).

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

### Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*



A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c)$$

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

$\Rightarrow$  limit of  $f(x) - f(c)$  as  $x \rightarrow c$  exists

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

$\Rightarrow$  limit of  $f(x) - f(c)$  as  $x \rightarrow c$  exists and

$$\lim_{x \rightarrow c} (f(x) - f(c))$$

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

$\Rightarrow$  limit of  $f(x) - f(c)$  as  $x \rightarrow c$  exists and

$$\lim_{x \rightarrow c} (f(x) - f(c)) = \left( \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right) \left( \lim_{x \rightarrow c} (x - c) \right)$$

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

$\Rightarrow$  limit of  $f(x) - f(c)$  as  $x \rightarrow c$  exists and

$$\lim_{x \rightarrow c} (f(x) - f(c)) = \left( \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right) \left( \lim_{x \rightarrow c} (x - c) \right) = f'(c) \cdot 0 = 0.$$

A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

$\Rightarrow$  limit of  $f(x) - f(c)$  as  $x \rightarrow c$  exists and

$$\lim_{x \rightarrow c} (f(x) - f(c)) = \left( \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right) \left( \lim_{x \rightarrow c} (x - c) \right) = f'(c) \cdot 0 = 0.$$

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c)$$



A continuous function exists which is not differentiable at any point (Weierstrass).  
However,

## Proposition

*Let  $f: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , then it is continuous at  $c$ .*

**Proof:**

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) \quad \text{and} \quad \lim_{x \rightarrow c} (x - c) = 0.$$

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

$\Rightarrow$  limit of  $f(x) - f(c)$  as  $x \rightarrow c$  exists and

$$\lim_{x \rightarrow c} (f(x) - f(c)) = \left( \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \right) \left( \lim_{x \rightarrow c} (x - c) \right) = f'(c) \cdot 0 = 0.$$

$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c) \Rightarrow f$  is continuous at  $c$ .



## Proposition (Linearity)

*Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .*

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I$ ,  $x \neq c$ ,

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c}$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c}$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$



## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c}$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

$$\frac{h(x) - h(c)}{x - c}$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

$$\frac{h(x) - h(c)}{x - c} = \frac{(f(x) + g(x)) - (f(c) + g(c))}{x - c}$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

$$\frac{h(x) - h(c)}{x - c} = \frac{(f(x) + g(x)) - (f(c) + g(c))}{x - c} = \frac{f(x) - f(c)}{x - c} + \frac{g(x) - g(c)}{x - c}.$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

$$\frac{h(x) - h(c)}{x - c} = \frac{(f(x) + g(x)) - (f(c) + g(c))}{x - c} = \frac{f(x) - f(c)}{x - c} + \frac{g(x) - g(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of RHS exists



## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

$$\frac{h(x) - h(c)}{x - c} = \frac{(f(x) + g(x)) - (f(c) + g(c))}{x - c} = \frac{f(x) - f(c)}{x - c} + \frac{g(x) - g(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of RHS exists and

$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c}$$

## Proposition (Linearity)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c \in I$ , and  $\alpha \in \mathbb{R}$ .

- (i) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := \alpha f(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = \alpha f'(c)$ .
- (ii) Define  $h: I \rightarrow \mathbb{R}$  by  $h(x) := f(x) + g(x)$ . Then  $h$  is differentiable at  $c$  and  $h'(c) = f'(c) + g'(c)$ .

**Proof:** (i) For  $x \in I, x \neq c$ , 
$$\frac{h(x) - h(c)}{x - c} = \frac{\alpha f(x) - \alpha f(c)}{x - c} = \alpha \frac{f(x) - f(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of the RHS exists and 
$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \alpha \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

(ii) For  $x \in I, x \neq c$ ,

$$\frac{h(x) - h(c)}{x - c} = \frac{(f(x) + g(x)) - (f(c) + g(c))}{x - c} = \frac{f(x) - f(c)}{x - c} + \frac{g(x) - g(c)}{x - c}.$$

The limit as  $x \rightarrow c$  of RHS exists and

$$\lim_{x \rightarrow c} \frac{h(x) - h(c)}{x - c} = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} + \lim_{x \rightarrow c} \frac{g(x) - g(c)}{x - c}.$$

□

## Proposition (Product rule / Leibniz rule)

*Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be functions differentiable at  $c$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := f(x)g(x)$ ,*

## Proposition (Product rule / Leibniz rule)

*Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be functions differentiable at  $c$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := f(x)g(x)$ , then  $h$  is differentiable at  $c$  and*

$$h'(c) = f(c)g'(c) + f'(c)g(c).$$

## Proposition (Product rule / Leibniz rule)

*Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be functions differentiable at  $c$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := f(x)g(x)$ , then  $h$  is differentiable at  $c$  and*

$$h'(c) = f(c)g'(c) + f'(c)g(c).$$

**Proof:** Exercise.

## Proposition (Product rule / Leibniz rule)

*Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be functions differentiable at  $c$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := f(x)g(x)$ , then  $h$  is differentiable at  $c$  and*

$$h'(c) = f(c)g'(c) + f'(c)g(c).$$

**Proof:** Exercise. Hint:  $f(x)g(x) - f(c)g(c) = f(x)(g(x) - g(c)) + (f(x) - f(c))g(c)$

## Proposition (Product rule / Leibniz rule)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be functions differentiable at  $c$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := f(x)g(x)$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f(c)g'(c) + f'(c)g(c).$$


**Proof:** Exercise. Hint:  $f(x)g(x) - f(c)g(c) = f(x)(g(x) - g(c)) + (f(x) - f(c))g(c)$   
or  $\Delta(f \cdot g) = f \cdot \Delta g + \Delta f \cdot g$ .

## Proposition (Product rule / Leibniz rule)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be functions differentiable at  $c$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := f(x)g(x)$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f(c)g'(c) + f'(c)g(c).$$

**Proof:** Exercise. Hint:  $f(x)g(x) - f(c)g(c) = f(x)(g(x) - g(c)) + (f(x) - f(c))g(c)$   
or  $\Delta(f \cdot g) = f \cdot \Delta g + \Delta f \cdot g.$





### Proposition (Quotient rule)

*Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c$  and  $g(x) \neq 0$  for all  $x \in I$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := \frac{f(x)}{g(x)}$ ,*

### Proposition (Quotient rule)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c$  and  $g(x) \neq 0$  for all  $x \in I$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := \frac{f(x)}{g(x)}$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = \frac{f'(c)g(c) - f(c)g'(c)}{(g(c))^2}.$$

### Proposition (Quotient rule)

Let  $I$  be an interval, let  $f: I \rightarrow \mathbb{R}$  and  $g: I \rightarrow \mathbb{R}$  be differentiable at  $c$  and  $g(x) \neq 0$  for all  $x \in I$ . If  $h: I \rightarrow \mathbb{R}$  is defined by  $h(x) := \frac{f(x)}{g(x)}$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = \frac{f'(c)g(c) - f(c)g'(c)}{(g(c))^2}.$$

**Proof:** Exercise.

## Proposition (Chain rule)

*Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ .*

## Proposition (Chain rule)

*Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ ,*

## Proposition (Chain rule)

*Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and*

$$h'(c) = f'(g(c))g'(c).$$

## Proposition (Chain rule)

*Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and*

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases}$$



## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

$$f(y) - f(d) = u(y)(y - d)$$

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

$$f(y) - f(d) = u(y)(y - d) \quad \text{and} \quad g(x) - g(c) = v(x)(x - c) \quad \forall x, y.$$

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

$$f(y) - f(d) = u(y)(y - d) \quad \text{and} \quad g(x) - g(c) = v(x)(x - c) \quad \forall x, y.$$

$$\implies h(x) - h(c)$$

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

$$f(y) - f(d) = u(y)(y - d) \quad \text{and} \quad g(x) - g(c) = v(x)(x - c) \quad \forall x, y.$$

$$\implies h(x) - h(c) = f(g(x)) - f(g(c))$$

## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

$$f(y) - f(d) = u(y)(y - d) \quad \text{and} \quad g(x) - g(c) = v(x)(x - c) \quad \forall x, y.$$

$$\implies h(x) - h(c) = f(g(x)) - f(g(c)) = u(g(x))(g(x) - g(c))$$



## Proposition (Chain rule)

Let  $I_1, I_2$  be intervals, let  $g: I_1 \rightarrow I_2$  be differentiable at  $c \in I_1$ , and  $f: I_2 \rightarrow \mathbb{R}$  be differentiable at  $g(c)$ . If  $h: I_1 \rightarrow \mathbb{R}$  is defined by  $h(x) := (f \circ g)(x) = f(g(x))$ , then  $h$  is differentiable at  $c$  and

$$h'(c) = f'(g(c))g'(c).$$

**Proof:** Let  $d := g(c)$ .

Define  $u: I_2 \rightarrow \mathbb{R}$  and  $v: I_1 \rightarrow \mathbb{R}$  by

$$u(y) := \begin{cases} \frac{f(y)-f(d)}{y-d} & \text{if } y \neq d, \\ f'(d) & \text{if } y = d, \end{cases} \quad v(x) := \begin{cases} \frac{g(x)-g(c)}{x-c} & \text{if } x \neq c, \\ g'(c) & \text{if } x = c. \end{cases}$$

$f$  differentiable at  $d = g(c) \implies u$  is continuous at  $d$ .

$g$  differentiable at  $c \implies v$  is continuous at  $c$ .

$$f(y) - f(d) = u(y)(y - d) \quad \text{and} \quad g(x) - g(c) = v(x)(x - c) \quad \forall x, y.$$

$$\implies h(x) - h(c) = f(g(x)) - f(g(c)) = u(g(x))(g(x) - g(c)) = u(g(x))(v(x)(x - c)).$$

If  $x \neq c$ , then 
$$\frac{h(x) - h(c)}{x - c}$$

If  $x \neq c$ , then 
$$\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$$

If  $x \neq c$ , then  $\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$

$u$  is continuous at  $d \implies \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c)).$

If  $x \neq c$ , then 
$$\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$$

$u$  is continuous at  $d \implies \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c)).$

$v$  is continuous at  $c \implies \lim_{x \rightarrow c} v(x) = g'(c).$

If  $x \neq c$ , then 
$$\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$$

$u$  is continuous at  $d \implies \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c)).$

$v$  is continuous at  $c \implies \lim_{x \rightarrow c} v(x) = g'(c).$

$g$  is continuous at  $c \implies \lim_{x \rightarrow c} g(x) = g(c).$

If  $x \neq c$ , then 
$$\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$$

$u$  is continuous at  $d \Rightarrow \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c)).$

$v$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} v(x) = g'(c).$

$g$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} g(x) = g(c).$

$\Rightarrow \lim_{x \rightarrow c} u(g(x))v(x) \text{ exists}$

If  $x \neq c$ , then  $\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$

$u$  is continuous at  $d \Rightarrow \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c)).$

$v$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} v(x) = g'(c).$

$g$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} g(x) = g(c).$

$\Rightarrow \lim_{x \rightarrow c} u(g(x))v(x)$  exists and equals  $f'(g(c))g'(c).$



If  $x \neq c$ , then 
$$\frac{h(x) - h(c)}{x - c} = u(g(x))v(x).$$

$u$  is continuous at  $d \Rightarrow \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c)).$

$v$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} v(x) = g'(c).$

$g$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} g(x) = g(c).$

$\Rightarrow \lim_{x \rightarrow c} u(g(x))v(x)$  exists and equals  $f'(g(c))g'(c).$

$\Rightarrow h$  is differentiable at  $c$

If  $x \neq c$ , then  $\frac{h(x) - h(c)}{x - c} = u(g(x))v(x)$ .

$u$  is continuous at  $d \Rightarrow \lim_{y \rightarrow d} u(y) = f'(d) = f'(g(c))$ .

$v$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} v(x) = g'(c)$ .

$g$  is continuous at  $c \Rightarrow \lim_{x \rightarrow c} g(x) = g(c)$ .

$\Rightarrow \lim_{x \rightarrow c} u(g(x))v(x)$  exists and equals  $f'(g(c))g'(c)$ .

$\Rightarrow h$  is differentiable at  $c$  and  $h'(c) = f'(g(c))g'(c)$ .



**Exercise:** Prove the following simple version of L'Hôpital's rule:

**Exercise:** Prove the following simple version of L'Hôpital's rule:

Suppose  $f: (a, b) \rightarrow \mathbb{R}$  and  $g: (a, b) \rightarrow \mathbb{R}$  are differentiable functions whose derivatives  $f'$  and  $g'$  are continuous functions.

**Exercise:** Prove the following simple version of L'Hôpital's rule:

Suppose  $f: (a, b) \rightarrow \mathbb{R}$  and  $g: (a, b) \rightarrow \mathbb{R}$  are differentiable functions whose derivatives  $f'$  and  $g'$  are continuous functions. Suppose that at  $c \in (a, b)$ ,  $f(c) = 0$ ,  $g(c) = 0$ ,  $g'(x) \neq 0$  for all  $x \in (a, b)$ , and  $g(x) \neq 0$  whenever  $x \neq c$ .

**Exercise:** Prove the following simple version of L'Hôpital's rule:

Suppose  $f: (a, b) \rightarrow \mathbb{R}$  and  $g: (a, b) \rightarrow \mathbb{R}$  are differentiable functions whose derivatives  $f'$  and  $g'$  are continuous functions. Suppose that at  $c \in (a, b)$ ,  $f(c) = 0$ ,  $g(c) = 0$ ,  $g'(x) \neq 0$  for all  $x \in (a, b)$ , and  $g(x) \neq 0$  whenever  $x \neq c$ .

Note that the limit of  $f'(x)/g'(x)$  as  $x$  goes to  $c$  exists.

**Exercise:** Prove the following simple version of L'Hôpital's rule:

Suppose  $f: (a, b) \rightarrow \mathbb{R}$  and  $g: (a, b) \rightarrow \mathbb{R}$  are differentiable functions whose derivatives  $f'$  and  $g'$  are continuous functions. Suppose that at  $c \in (a, b)$ ,  $f(c) = 0$ ,  $g(c) = 0$ ,  $g'(x) \neq 0$  for all  $x \in (a, b)$ , and  $g(x) \neq 0$  whenever  $x \neq c$ .

Note that the limit of  $f'(x)/g'(x)$  as  $x$  goes to  $c$  exists.

Show that

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}.$$