

BA: 2.5

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Given a sequence $\{x_n\}_{n=1}^{\infty}$, a *series* is the formal object

$$\sum_{n=1}^{\infty} x_n$$

Definition

Given a sequence $\{x_n\}_{n=1}^{\infty}$, a *series* is the formal object

$$\sum_{n=1}^{\infty} x_n$$

A series *converges* if the sequence $\{s_k\}_{k=1}^{\infty}$ defined by

$$s_k := \sum_{n=1}^k x_n = x_1 + x_2 + \cdots + x_k \quad \text{converges.}$$

Definition

Given a sequence $\{x_n\}_{n=1}^{\infty}$, a *series* is the formal object

$$\sum_{n=1}^{\infty} x_n$$

A series *converges* if the sequence $\{s_k\}_{k=1}^{\infty}$ defined by

$$s_k := \sum_{n=1}^k x_n = x_1 + x_2 + \cdots + x_k \quad \text{converges.}$$

The numbers s_k are called *partial sums*.

Definition

Given a sequence $\{x_n\}_{n=1}^{\infty}$, a *series* is the formal object

$$\sum_{n=1}^{\infty} x_n$$

A series *converges* if the sequence $\{s_k\}_{k=1}^{\infty}$ defined by

$$s_k := \sum_{n=1}^k x_n = x_1 + x_2 + \cdots + x_k \quad \text{converges.}$$

The numbers s_k are called *partial sums*. If the series converges, we write

$$\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} s_k \quad (\text{cheating a little by treating the series as a number}).$$

Definition

Given a sequence $\{x_n\}_{n=1}^{\infty}$, a *series* is the formal object

$$\sum_{n=1}^{\infty} x_n$$

A series *converges* if the sequence $\{s_k\}_{k=1}^{\infty}$ defined by

$$s_k := \sum_{n=1}^k x_n = x_1 + x_2 + \cdots + x_k \quad \text{converges.}$$

The numbers s_k are called *partial sums*. If the series converges, we write

$$\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} s_k \quad (\text{cheating a little by treating the series as a number}).$$

If $\{s_k\}_{k=1}^{\infty}$ diverges, we say the series is *divergent*.

If a series is convergent, then

$$\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n.$$

If a series is convergent, then $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n$.

If the series does not converge, the right hand side is nonsense.

If a series is convergent, then $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n$.

If the series does not converge, the right hand side is nonsense.

So (unlike for sequences) $\sum_{n=1}^{\infty} x_n$ means two different things:

If a series is convergent, then $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n$.

If the series does not converge, the right hand side is nonsense.

So (unlike for sequences) $\sum_{n=1}^{\infty} x_n$ means two different things:

1) notation for the series,

If a series is convergent, then $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n$.

If the series does not converge, the right hand side is nonsense.

So (unlike for sequences) $\sum_{n=1}^{\infty} x_n$ means two different things:

- 1) notation for the series,
- 2) the actual limit.

If a series is convergent, then $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n$.

If the series does not converge, the right hand side is nonsense.

So (unlike for sequences) $\sum_{n=1}^{\infty} x_n$ means two different things:

- 1) notation for the series,
- 2) the actual limit.

Remark: We could start at a different index: $\sum_{n=0}^{\infty} r^n = \sum_{n=1}^{\infty} r^{n-1}$.

If a series is convergent, then $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} \sum_{n=1}^k x_n$.

If the series does not converge, the right hand side is nonsense.

So (unlike for sequences) $\sum_{n=1}^{\infty} x_n$ means two different things:

- 1) notation for the series,
- 2) the actual limit.

Remark: We could start at a different index: $\sum_{n=0}^{\infty} r^n = \sum_{n=1}^{\infty} r^{n-1}$.

Remark: It is common (but we will not) to write $\sum_{n=1}^{\infty} x_n$ informally as

$$x_1 + x_2 + x_3 + \cdots$$


Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.

Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.

Proof: By induction (exercise), $\left(\sum_{n=1}^k \frac{1}{2^n} \right) + \frac{1}{2^k} = 1$.


Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.

Proof: By induction (exercise), $\left(\sum_{n=1}^k \frac{1}{2^n} \right) + \frac{1}{2^k} = 1$.



Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.


Proof: By induction (exercise), $\left(\sum_{n=1}^k \frac{1}{2^n} \right) + \frac{1}{2^k} = 1$.



Let s_k be the partial sum, then $|1 - s_k| = \left| 1 - \sum_{n=1}^k \frac{1}{2^n} \right|$

Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.


Proof: By induction (exercise), $\left(\sum_{n=1}^k \frac{1}{2^n} \right) + \frac{1}{2^k} = 1$.



Let s_k be the partial sum, then $|1 - s_k| = \left| 1 - \sum_{n=1}^k \frac{1}{2^n} \right| = \left| \frac{1}{2^k} \right| = \frac{1}{2^k}$.

Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.

Proof: By induction (exercise), $\left(\sum_{n=1}^k \frac{1}{2^n} \right) + \frac{1}{2^k} = 1$.




Let s_k be the partial sum, then $|1 - s_k| = \left| 1 - \sum_{n=1}^k \frac{1}{2^n} \right| = \left| \frac{1}{2^k} \right| = \frac{1}{2^k}$.

$\left\{ \frac{1}{2^k} \right\}_{k=1}^{\infty} = \left\{ |1 - s_k| \right\}_{k=1}^{\infty}$ converges to 0

Example: $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges and the limit is 1. That is, $\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{k \rightarrow \infty} \sum_{n=1}^k \frac{1}{2^n} = 1$.

Proof: By induction (exercise), $\left(\sum_{n=1}^k \frac{1}{2^n} \right) + \frac{1}{2^k} = 1$.



Let s_k be the partial sum, then $|1 - s_k| = \left| 1 - \sum_{n=1}^k \frac{1}{2^n} \right| = \left| \frac{1}{2^k} \right| = \frac{1}{2^k}$.

$\left\{ \frac{1}{2^k} \right\}_{k=1}^{\infty} = \{|1 - s_k|\}_{k=1}^{\infty}$ converges to 0 $\Rightarrow \{s_k\}_{k=1}^{\infty}$ converges to 1.

□

Proposition

Suppose $-1 < r < 1$. Then the geometric series $\sum_{n=0}^{\infty} r^n$ converges, and $\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$.

Proposition

Suppose $-1 < r < 1$. Then the geometric series $\sum_{n=0}^{\infty} r^n$ converges, and $\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$.

Proof: Exercise.

Proposition

Suppose $-1 < r < 1$. Then the geometric series $\sum_{n=0}^{\infty} r^n$ converges, and $\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$.

Proof: Exercise.

One proves

$$\sum_{n=0}^{k-1} r^n = \frac{1-r^k}{1-r},$$

Proposition

Suppose $-1 < r < 1$. Then the geometric series $\sum_{n=0}^{\infty} r^n$ converges, and $\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$.

Proof: Exercise.

One proves

$$\sum_{n=0}^{k-1} r^n = \frac{1-r^k}{1-r},$$

and then one takes the limit $k \rightarrow \infty$.

Proposition (tail of a series)

Let $\sum_{n=1}^{\infty} x_n$ be a series. Let $M \in \mathbb{N}$. Then

$$\sum_{n=1}^{\infty} x_n \text{ converges if and only if } \sum_{n=M}^{\infty} x_n \text{ converges.}$$

Proposition (tail of a series)

Let $\sum_{n=1}^{\infty} x_n$ be a series. Let $M \in \mathbb{N}$. Then

$$\sum_{n=1}^{\infty} x_n \text{ converges if and only if } \sum_{n=M}^{\infty} x_n \text{ converges.}$$

Proof: Consider partial sums of the two series (for $k \geq M$)

$$\sum_{n=1}^k x_n = \left(\sum_{n=1}^{M-1} x_n \right) + \sum_{n=M}^k x_n.$$

Proposition (tail of a series)

Let $\sum_{n=1}^{\infty} x_n$ be a series. Let $M \in \mathbb{N}$. Then

$$\sum_{n=1}^{\infty} x_n \text{ converges if and only if } \sum_{n=M}^{\infty} x_n \text{ converges.}$$

Proof: Consider partial sums of the two series (for $k \geq M$)

$$\sum_{n=1}^k x_n = \left(\sum_{n=1}^{M-1} x_n \right) + \sum_{n=M}^k x_n.$$

Note that $\sum_{n=1}^{M-1} x_n$ is a fixed number.

Proposition (tail of a series)

Let $\sum_{n=1}^{\infty} x_n$ be a series. Let $M \in \mathbb{N}$. Then

$$\sum_{n=1}^{\infty} x_n \text{ converges if and only if } \sum_{n=M}^{\infty} x_n \text{ converges.}$$

Proof: Consider partial sums of the two series (for $k \geq M$)

$$\sum_{n=1}^k x_n = \left(\sum_{n=1}^{M-1} x_n \right) + \sum_{n=M}^k x_n.$$

Note that $\sum_{n=1}^{M-1} x_n$ is a fixed number.

Adding a constant does not change the convergence of a sequence. □

Definition

$\sum_{n=1}^{\infty} x_n$ is *Cauchy* (*Cauchy series*) if the sequence of partial sums $\{s_n\}_{n=1}^{\infty}$ is Cauchy.

Definition

$\sum_{n=1}^{\infty} x_n$ is *Cauchy* (*Cauchy series*) if the sequence of partial sums $\{s_n\}_{n=1}^{\infty}$ is Cauchy.

A sequence converges \Leftrightarrow it is Cauchy.

Definition

$\sum_{n=1}^{\infty} x_n$ is *Cauchy* (*Cauchy series*) if the sequence of partial sums $\{s_n\}_{n=1}^{\infty}$ is Cauchy.

A sequence converges \Leftrightarrow it is Cauchy. So: A series converges \Leftrightarrow it is Cauchy.

Definition

$\sum_{n=1}^{\infty} x_n$ is *Cauchy* (*Cauchy series*) if the sequence of partial sums $\{s_n\}_{n=1}^{\infty}$ is Cauchy.

A sequence converges \Leftrightarrow it is Cauchy. So: A series converges \Leftrightarrow it is Cauchy.

$$\sum_{n=1}^{\infty} x_n \text{ is Cauchy} \Leftrightarrow \forall \epsilon > 0, \exists M \in \mathbb{N}, \text{ s. t. } \forall n, k \geq M, \left| \left(\sum_{i=1}^k x_i \right) - \left(\sum_{i=1}^n x_i \right) \right| < \epsilon.$$

Definition

$\sum_{n=1}^{\infty} x_n$ is *Cauchy* (*Cauchy series*) if the sequence of partial sums $\{s_n\}_{n=1}^{\infty}$ is Cauchy.

A sequence converges \Leftrightarrow it is Cauchy. So: A series converges \Leftrightarrow it is Cauchy.

$$\sum_{n=1}^{\infty} x_n \text{ is Cauchy} \Leftrightarrow \forall \epsilon > 0, \exists M \in \mathbb{N}, \text{ s. t. } \forall n, k \geq M, \left| \left(\sum_{i=1}^k x_i \right) - \left(\sum_{i=1}^n x_i \right) \right| < \epsilon.$$

WLOG assume $n < k$. Then

$$\left| \left(\sum_{i=1}^k x_i \right) - \left(\sum_{i=1}^n x_i \right) \right| = \left| \sum_{i=n+1}^k x_i \right| < \epsilon.$$

Definition

$\sum_{n=1}^{\infty} x_n$ is *Cauchy* (*Cauchy series*) if the sequence of partial sums $\{s_n\}_{n=1}^{\infty}$ is Cauchy.

A sequence converges \Leftrightarrow it is Cauchy. So: A series converges \Leftrightarrow it is Cauchy.

$$\sum_{n=1}^{\infty} x_n \text{ is Cauchy} \Leftrightarrow \forall \epsilon > 0, \exists M \in \mathbb{N}, \text{ s. t. } \forall n, k \geq M, \left| \left(\sum_{i=1}^k x_i \right) - \left(\sum_{i=1}^n x_i \right) \right| < \epsilon.$$

WLOG assume $n < k$. Then $\left| \left(\sum_{i=1}^k x_i \right) - \left(\sum_{i=1}^n x_i \right) \right| = \left| \sum_{i=n+1}^k x_i \right| < \epsilon. \Rightarrow$

Proposition

$$\sum_{n=1}^{\infty} x_n \text{ is Cauchy} \Leftrightarrow \forall \epsilon > 0, \exists M \in \mathbb{N} \text{ such that } \forall n \geq M \text{ and } \forall k > n, \left| \sum_{i=n+1}^k x_i \right| < \epsilon.$$

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given.

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M, \quad \epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right|$

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M, \quad \epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right| = |x_{n+1}|.$

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M, \quad \epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right| = |x_{n+1}|.$

\Rightarrow for every $n \geq M + 1$, we have $|x_n| < \epsilon$.



Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M$, $\epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right| = |x_{n+1}|$.

\Rightarrow for every $n \geq M + 1$, we have $|x_n| < \epsilon$. □

Example: If $r \geq 1$ or $r \leq -1$, then the geometric series $\sum_{n=0}^{\infty} r^n$ diverges.

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M, \quad \epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right| = |x_{n+1}|$.

\Rightarrow for every $n \geq M + 1$, we have $|x_n| < \epsilon$. □

Example: If $r \geq 1$ or $r \leq -1$, then the geometric series $\sum_{n=0}^{\infty} r^n$ diverges.

Proof: $|r^n| = |r|^n \geq 1^n = 1$

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M, \quad \epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right| = |x_{n+1}|$.

\Rightarrow for every $n \geq M + 1$, we have $|x_n| < \epsilon$. □

Example: If $r \geq 1$ or $r \leq -1$, then the geometric series $\sum_{n=0}^{\infty} r^n$ diverges.

Proof: $|r^n| = |r|^n \geq 1^n = 1 \Rightarrow$ terms do not go to 0

Proposition

Let $\sum_{n=1}^{\infty} x_n$ be a convergent series. Then $\{x_n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} x_n = 0$.

Proof: $\sum_{n=1}^{\infty} x_n$ is convergent \Rightarrow it is Cauchy.

Let $\epsilon > 0$ be given. $\exists M$ such that $\forall n \geq M, \quad \epsilon > \left| \sum_{i=n+1}^{n+1} x_i \right| = |x_{n+1}|$.

\Rightarrow for every $n \geq M + 1$, we have $|x_n| < \epsilon$. □

Example: If $r \geq 1$ or $r \leq -1$, then the geometric series $\sum_{n=0}^{\infty} r^n$ diverges.

Proof: $|r^n| = |r|^n \geq 1^n = 1 \Rightarrow$ terms do not go to 0 $\Rightarrow \sum_{n=1}^{\infty} r^n$ diverges. □

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1,$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right),$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right),$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right).$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k} = \frac{1}{2}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k} = \frac{1}{2} \Rightarrow s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right)$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k} = \frac{1}{2} \Rightarrow s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right) \geq 1 + \sum_{i=1}^k \frac{1}{2} = 1 + \frac{k}{2}.$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k} = \frac{1}{2} \Rightarrow s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right) \geq 1 + \sum_{i=1}^k \frac{1}{2} = 1 + \frac{k}{2}.$$

$\{\frac{k}{2}\}_{k=1}^{\infty}$ is unbounded

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k} = \frac{1}{2} \Rightarrow s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right) \geq 1 + \sum_{i=1}^k \frac{1}{2} = 1 + \frac{k}{2}.$$

$$\left\{ \frac{k}{2} \right\}_{k=1}^{\infty} \text{ is unbounded} \Rightarrow \{s_{2^k}\}_{k=1}^{\infty} \text{ is unbounded}$$

Example: The *harmonic series* $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (despite the fact that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$).

Proof: Write the partial sums s_n for $n = 2^k$ as:

$$s_1 = 1, \quad s_2 = (1) + \left(\frac{1}{2}\right), \quad s_4 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right), \quad s_8 = (1) + \left(\frac{1}{2}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right),$$

$$s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right). \quad \text{Note } \frac{1}{3} + \frac{1}{4} \geq \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ and } \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} \geq \frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{1}{2}.$$

$$\sum_{m=2^{k-1}+1}^{2^k} \frac{1}{m} \geq \sum_{m=2^{k-1}+1}^{2^k} \frac{1}{2^k} = (2^{k-1}) \frac{1}{2^k} = \frac{1}{2} \Rightarrow s_{2^k} = 1 + \sum_{i=1}^k \left(\sum_{m=2^{i-1}+1}^{2^i} \frac{1}{m} \right) \geq 1 + \sum_{i=1}^k \frac{1}{2} = 1 + \frac{k}{2}.$$

$$\left\{ \frac{k}{2} \right\}_{k=1}^{\infty} \text{ is unbounded} \Rightarrow \{s_{2^k}\}_{k=1}^{\infty} \text{ is unbounded} \Rightarrow \{s_n\}_{n=1}^{\infty} \text{ is unbounded.}$$

So $\{s_n\}_{n=1}^{\infty}$ diverges, that is, $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges. □

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series.

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series. Then

(i) $\sum_{n=1}^{\infty} \alpha x_n$ converges and $\sum_{n=1}^{\infty} \alpha x_n = \alpha \sum_{n=1}^{\infty} x_n$.

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series. Then

(i) $\sum_{n=1}^{\infty} \alpha x_n$ converges and $\sum_{n=1}^{\infty} \alpha x_n = \alpha \sum_{n=1}^{\infty} x_n$.

(ii) $\sum_{n=1}^{\infty} (x_n + y_n)$ converges and $\sum_{n=1}^{\infty} (x_n + y_n) = \left(\sum_{n=1}^{\infty} x_n \right) + \left(\sum_{n=1}^{\infty} y_n \right)$.

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series. Then

(i) $\sum_{n=1}^{\infty} \alpha x_n$ converges and $\sum_{n=1}^{\infty} \alpha x_n = \alpha \sum_{n=1}^{\infty} x_n$.

(ii) $\sum_{n=1}^{\infty} (x_n + y_n)$ converges and $\sum_{n=1}^{\infty} (x_n + y_n) = \left(\sum_{n=1}^{\infty} x_n \right) + \left(\sum_{n=1}^{\infty} y_n \right)$.

Proof: (i) The k th partial sum is $\sum_{n=1}^k \alpha x_n = \alpha \left(\sum_{n=1}^k x_n \right)$.

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series. Then

(i) $\sum_{n=1}^{\infty} \alpha x_n$ converges and $\sum_{n=1}^{\infty} \alpha x_n = \alpha \sum_{n=1}^{\infty} x_n$.

(ii) $\sum_{n=1}^{\infty} (x_n + y_n)$ converges and $\sum_{n=1}^{\infty} (x_n + y_n) = \left(\sum_{n=1}^{\infty} x_n \right) + \left(\sum_{n=1}^{\infty} y_n \right)$.

Proof: (i) The k th partial sum is $\sum_{n=1}^k \alpha x_n = \alpha \left(\sum_{n=1}^k x_n \right)$.

Constant multiple of a convergent sequence is convergent.

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series. Then

(i) $\sum_{n=1}^{\infty} \alpha x_n$ converges and $\sum_{n=1}^{\infty} \alpha x_n = \alpha \sum_{n=1}^{\infty} x_n$.

(ii) $\sum_{n=1}^{\infty} (x_n + y_n)$ converges and $\sum_{n=1}^{\infty} (x_n + y_n) = \left(\sum_{n=1}^{\infty} x_n \right) + \left(\sum_{n=1}^{\infty} y_n \right)$.

Proof: (i) The k th partial sum is $\sum_{n=1}^k \alpha x_n = \alpha \left(\sum_{n=1}^k x_n \right)$.

Constant multiple of a convergent sequence is convergent.

(ii) The k th partial sum is $\sum_{n=1}^k (x_n + y_n) = \left(\sum_{n=1}^k x_n \right) + \left(\sum_{n=1}^k y_n \right)$.

Proposition (Linearity of series)

Let $\alpha \in \mathbb{R}$ and $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be convergent series. Then

(i) $\sum_{n=1}^{\infty} \alpha x_n$ converges and $\sum_{n=1}^{\infty} \alpha x_n = \alpha \sum_{n=1}^{\infty} x_n$.

(ii) $\sum_{n=1}^{\infty} (x_n + y_n)$ converges and $\sum_{n=1}^{\infty} (x_n + y_n) = \left(\sum_{n=1}^{\infty} x_n \right) + \left(\sum_{n=1}^{\infty} y_n \right)$.

Proof: (i) The k th partial sum is $\sum_{n=1}^k \alpha x_n = \alpha \left(\sum_{n=1}^k x_n \right)$.

Constant multiple of a convergent sequence is convergent.

(ii) The k th partial sum is $\sum_{n=1}^k (x_n + y_n) = \left(\sum_{n=1}^k x_n \right) + \left(\sum_{n=1}^k y_n \right)$.

Sum of convergent sequences converges to the sum of the limits. □

Example: By the first item, if $|r| < 1$ and $i \in \mathbb{N}$, then

$$r^i \sum_{n=0}^{\infty} r^n = \sum_{n=0}^{\infty} r^{n+i} = \sum_{n=i}^{\infty} r^n.$$

Example: By the first item, if $|r| < 1$ and $i \in \mathbb{N}$, then

$$r^i \sum_{n=0}^{\infty} r^n = \sum_{n=0}^{\infty} r^{n+i} = \sum_{n=i}^{\infty} r^n.$$

Hence,

$$\sum_{n=i}^{\infty} r^n = \frac{r^i}{1-r}.$$

Example: By the first item, if $|r| < 1$ and $i \in \mathbb{N}$, then

$$r^i \sum_{n=0}^{\infty} r^n = \sum_{n=0}^{\infty} r^{n+i} = \sum_{n=i}^{\infty} r^n.$$

Hence,

$$\sum_{n=i}^{\infty} r^n = \frac{r^i}{1-r}.$$

Remark: Multiplying series is not as simple as adding (see 2.6).

Example: By the first item, if $|r| < 1$ and $i \in \mathbb{N}$, then

$$r^i \sum_{n=0}^{\infty} r^n = \sum_{n=0}^{\infty} r^{n+i} = \sum_{n=i}^{\infty} r^n.$$

Hence,

$$\sum_{n=i}^{\infty} r^n = \frac{r^i}{1-r}.$$

Remark: Multiplying series is not as simple as adding (see 2.6).

Obviously, we don't multiply term by term: $(a+b)(c+d) \neq ac+bd$.

Monotone sequences are easier than arbitrary sequences

Monotone sequences are easier than arbitrary sequences

$\Rightarrow \sum_{n=1}^{\infty} x_n$ where $x_n \geq 0$ for all n are easier, as partial sums are monotone increasing.

Monotone sequences are easier than arbitrary sequences

$\Rightarrow \sum_{n=1}^{\infty} x_n$ where $x_n \geq 0$ for all n are easier, as partial sums are monotone increasing.

Proposition

If $x_n \geq 0$ for all n , then $\sum_{n=1}^{\infty} x_n$ converges if and only if the sequence of partial sums is bounded above.

Monotone sequences are easier than arbitrary sequences

$\Rightarrow \sum_{n=1}^{\infty} x_n$ where $x_n \geq 0$ for all n are easier, as partial sums are monotone increasing.

Proposition

If $x_n \geq 0$ for all n , then $\sum_{n=1}^{\infty} x_n$ converges if and only if the sequence of partial sums is bounded above.

The limit of a monotone increasing sequence is the supremum:

Monotone sequences are easier than arbitrary sequences

$\Rightarrow \sum_{n=1}^{\infty} x_n$ where $x_n \geq 0$ for all n are easier, as partial sums are monotone increasing.

Proposition

If $x_n \geq 0$ for all n , then $\sum_{n=1}^{\infty} x_n$ converges if and only if the sequence of partial sums is bounded above.

The limit of a monotone increasing sequence is the supremum:

If $x_n \geq 0$ for all n , then

$$\sum_{n=1}^k x_n \leq \sum_{n=1}^{\infty} x_n.$$

Monotone sequences are easier than arbitrary sequences

$\Rightarrow \sum_{n=1}^{\infty} x_n$ where $x_n \geq 0$ for all n are easier, as partial sums are monotone increasing.

Proposition

If $x_n \geq 0$ for all n , then $\sum_{n=1}^{\infty} x_n$ converges if and only if the sequence of partial sums is bounded above.

The limit of a monotone increasing sequence is the supremum:

If $x_n \geq 0$ for all n , then

$$\sum_{n=1}^k x_n \leq \sum_{n=1}^{\infty} x_n.$$

(still true if we allow infinite limits).

Definition

$\sum_{n=1}^{\infty} x_n$ *converges absolutely* if $\sum_{n=1}^{\infty} |x_n|$ converges.

Definition

$\sum_{n=1}^{\infty} x_n$ *converges absolutely* if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Definition

$\sum_{n=1}^{\infty} x_n$ *converges absolutely* if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Proposition

If $\sum_{n=1}^{\infty} x_n$ *converges absolutely*, then it *converges*.

Definition

$\sum_{n=1}^{\infty} x_n$ converges absolutely if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Proposition

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then it converges.

Proof: If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then $\sum_{n=1}^{\infty} |x_n|$ is convergent hence Cauchy.

Definition

$\sum_{n=1}^{\infty} x_n$ converges absolutely if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Proposition

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then it converges.

Proof: If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then $\sum_{n=1}^{\infty} |x_n|$ is convergent hence Cauchy.

$\forall \epsilon > 0, \exists M$ such that $\forall k \geq M$ and all $n > k$,
$$\sum_{i=k+1}^n |x_i| = \left| \sum_{i=k+1}^n |x_i| \right| < \epsilon.$$

Definition

$\sum_{n=1}^{\infty} x_n$ converges absolutely if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Proposition

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then it converges.

Proof: If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then $\sum_{n=1}^{\infty} |x_n|$ is convergent hence Cauchy.

$\forall \epsilon > 0, \exists M$ such that $\forall k \geq M$ and all $n > k$, $\sum_{i=k+1}^n |x_i| = \left| \sum_{i=k+1}^n |x_i| \right| < \epsilon$.

By triangle inequality, $\left| \sum_{i=k+1}^n x_i \right| \leq \sum_{i=k+1}^n |x_i|$

Definition

$\sum_{n=1}^{\infty} x_n$ converges absolutely if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Proposition

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then it converges.

Proof: If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then $\sum_{n=1}^{\infty} |x_n|$ is convergent hence Cauchy.

$\forall \epsilon > 0, \exists M$ such that $\forall k \geq M$ and all $n > k$, $\sum_{i=k+1}^n |x_i| = \left| \sum_{i=k+1}^n |x_i| \right| < \epsilon$.

By triangle inequality, $\left| \sum_{i=k+1}^n x_i \right| \leq \sum_{i=k+1}^n |x_i| < \epsilon$

Definition

$\sum_{n=1}^{\infty} x_n$ converges absolutely if $\sum_{n=1}^{\infty} |x_n|$ converges.

If a series converges, but not absolutely, it *converges conditionally*.

Proposition

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then it converges.

Proof: If $\sum_{n=1}^{\infty} x_n$ converges absolutely, then $\sum_{n=1}^{\infty} |x_n|$ is convergent hence Cauchy.

$\forall \epsilon > 0, \exists M$ such that $\forall k \geq M$ and all $n > k$, $\sum_{i=k+1}^n |x_i| = \left| \sum_{i=k+1}^n |x_i| \right| < \epsilon$.

By triangle inequality, $\left| \sum_{i=k+1}^n x_i \right| \leq \sum_{i=k+1}^n |x_i| < \epsilon \Rightarrow \sum_{n=1}^{\infty} x_n$ is Cauchy \Rightarrow converges. \square

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise),

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise), but $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise), but $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

So $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges conditionally.

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise), but $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

So $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges conditionally.

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, the limits of $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} |x_n|$ are in general different.

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise), but $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

So $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges conditionally.

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, the limits of $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} |x_n|$ are in general different.

However, (exercise) $\left| \sum_{i=1}^{\infty} x_i \right| \leq \sum_{i=1}^{\infty} |x_i|$.

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise), but $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

So $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges conditionally.

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, the limits of $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} |x_n|$ are in general different.

However, (exercise) $\left| \sum_{i=1}^{\infty} x_i \right| \leq \sum_{i=1}^{\infty} |x_i|$.

Absolutely convergent series behave nicely (e.g., can be rearranged, or multiplied, etc.).

Example: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges (exercise), but $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

So $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges conditionally.

If $\sum_{n=1}^{\infty} x_n$ converges absolutely, the limits of $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} |x_n|$ are in general different.

However, (exercise) $\left| \sum_{i=1}^{\infty} x_i \right| \leq \sum_{i=1}^{\infty} |x_i|$.

Absolutely convergent series behave nicely (e.g., can be rearranged, or multiplied, etc.).

Conditionally convergent series don't.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

- (i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n$

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n \geq \sum_{n=1}^k x_n$.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n \geq \sum_{n=1}^k x_n$.

Partial sums of $\sum_{n=1}^{\infty} x_n$ are bounded $\Rightarrow \sum_{n=1}^{\infty} x_n$ converges.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n \geq \sum_{n=1}^k x_n$.

Partial sums of $\sum_{n=1}^{\infty} x_n$ are bounded $\Rightarrow \sum_{n=1}^{\infty} x_n$ converges.

(ii) If $\sum_{n=1}^{\infty} x_n$ diverges, partial sums unbounded

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n \geq \sum_{n=1}^k x_n$.

Partial sums of $\sum_{n=1}^{\infty} x_n$ are bounded $\Rightarrow \sum_{n=1}^{\infty} x_n$ converges.

(ii) If $\sum_{n=1}^{\infty} x_n$ diverges, partial sums unbounded $\Rightarrow \forall B \exists k$ such that $B \leq \sum_{n=1}^k x_n$

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n \geq \sum_{n=1}^k x_n$.

Partial sums of $\sum_{n=1}^{\infty} x_n$ are bounded $\Rightarrow \sum_{n=1}^{\infty} x_n$ converges.

(ii) If $\sum_{n=1}^{\infty} x_n$ diverges, partial sums unbounded $\Rightarrow \forall B \exists k$ such that $B \leq \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

Proposition (Comparison test)

Let $\sum_{n=1}^{\infty} x_n$ and $\sum_{n=1}^{\infty} y_n$ be series such that $0 \leq x_n \leq y_n$ for all $n \in \mathbb{N}$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, then so does $\sum_{n=1}^{\infty} x_n$. (ii) If $\sum_{n=1}^{\infty} x_n$ diverges, then so does $\sum_{n=1}^{\infty} y_n$.

Proof: Partial sums monotone (terms ≥ 0) and $x_n \leq y_n$ for all $n \Rightarrow \forall k, \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

(i) If $\sum_{n=1}^{\infty} y_n$ converges, partial sums bounded $\Rightarrow \exists B$ such that $\forall k, B \geq \sum_{n=1}^k y_n \geq \sum_{n=1}^k x_n$.

Partial sums of $\sum_{n=1}^{\infty} x_n$ are bounded $\Rightarrow \sum_{n=1}^{\infty} x_n$ converges.

(ii) If $\sum_{n=1}^{\infty} x_n$ diverges, partial sums unbounded $\Rightarrow \forall B \exists k$ such that $B \leq \sum_{n=1}^k x_n \leq \sum_{n=1}^k y_n$.

\Rightarrow partial sums of $\sum_{n=1}^{\infty} y_n$ unbounded $\Rightarrow \sum_{n=1}^{\infty} y_n$ diverges. □

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1,$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right),$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right).$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right). \quad \text{Note } \frac{1}{2^p} + \frac{1}{3^p} < \frac{1}{2^p} + \frac{1}{2^p}$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right). \quad \text{Note } \frac{1}{2^p} + \frac{1}{3^p} < \frac{1}{2^p} + \frac{1}{2^p} \text{ and } \frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p} < \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p}.$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right). \quad \text{Note } \frac{1}{2^p} + \frac{1}{3^p} < \frac{1}{2^p} + \frac{1}{2^p} \text{ and } \frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p} < \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p}.$$

For all $k \geq 2$,

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right)$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right). \quad \text{Note } \frac{1}{2^p} + \frac{1}{3^p} < \frac{1}{2^p} + \frac{1}{2^p} \text{ and } \frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p} < \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p}.$$

For all $k \geq 2$,

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right) < 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{(2^i)^p} \right)$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right). \quad \text{Note } \frac{1}{2^p} + \frac{1}{3^p} < \frac{1}{2^p} + \frac{1}{2^p} \text{ and } \frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p} < \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p}.$$

For all $k \geq 2$,

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right) < 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{(2^i)^p} \right) = 1 + \sum_{i=1}^{k-1} \left(\frac{2^i}{(2^i)^p} \right)$$

Proposition (p -series or the p -test)

For $p \in \mathbb{R}$, $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

Proof: For $p \leq 1$, then $\frac{1}{n^p} \geq \frac{1}{n}$. $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$ diverges (comparison).

Suppose $p > 1$. Let s_n denote the n th partial sum (monotone seq. again).

$$s_1 = 1, \quad s_3 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right), \quad s_7 = (1) + \left(\frac{1}{2^p} + \frac{1}{3^p}\right) + \left(\frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p}\right),$$

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right). \quad \text{Note } \frac{1}{2^p} + \frac{1}{3^p} < \frac{1}{2^p} + \frac{1}{2^p} \text{ and } \frac{1}{4^p} + \frac{1}{5^p} + \frac{1}{6^p} + \frac{1}{7^p} < \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p} + \frac{1}{4^p}.$$

For all $k \geq 2$,

$$s_{2^k-1} = 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{m^p} \right) < 1 + \sum_{i=1}^{k-1} \left(\sum_{m=2^i}^{2^{i+1}-1} \frac{1}{(2^i)^p} \right) = 1 + \sum_{i=1}^{k-1} \left(\frac{2^i}{(2^i)^p} \right) = 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}} \right)^i.$$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}} \right)^i$ converges.

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}} \right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}} \right)^i$$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}}\right)^i \leq 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}}\right)^i \leq 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

For every n there is a $k \geq 2$ such that $n \leq 2^k - 1$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}}\right)^i \leq 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

For every n there is a $k \geq 2$ such that $n \leq 2^k - 1$

$$\Rightarrow s_n \leq s_{2^k-1}$$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}}\right)^i \leq 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

For every n there is a $k \geq 2$ such that $n \leq 2^k - 1$

$$\Rightarrow s_n \leq s_{2^k-1} < 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}}\right)^i \leq 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

For every n there is a $k \geq 2$ such that $n \leq 2^k - 1$

$$\Rightarrow s_n \leq s_{2^k-1} < 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

The sequence of partial sums is bounded

$$p > 1 \quad \Rightarrow \quad \frac{1}{2^{p-1}} < 1.$$

The geometric series $\sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i$ converges.

$$\Rightarrow s_{2^k-1} < 1 + \sum_{i=1}^{k-1} \left(\frac{1}{2^{p-1}}\right)^i \leq 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

For every n there is a $k \geq 2$ such that $n \leq 2^k - 1$

$$\Rightarrow s_n \leq s_{2^k-1} < 1 + \sum_{i=1}^{\infty} \left(\frac{1}{2^{p-1}}\right)^i.$$

The sequence of partial sums is bounded \Rightarrow the series converges.



Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test.

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test.

By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges. \square

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test. By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges. \square

Often easier to show a series converges than it is to find the limit.

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test. By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges. \square

Often easier to show a series converges than it is to find the limit.

E.g., easy to see $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges,

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test. By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges. \square

Often easier to show a series converges than it is to find the limit.

E.g., easy to see $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, much harder to show the limit is $\pi^2/6$.

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test. By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges. \square

Often easier to show a series converges than it is to find the limit.

E.g., easy to see $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, much harder to show the limit is $\pi^2/6$.

Remark: $\zeta(p) = \sum_{n=1}^{\infty} \frac{1}{n^p}$ is the “Riemann zeta function”.

Example: $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Proof: $\frac{1}{n^2 + 1} < \frac{1}{n^2}$ for all $n \in \mathbb{N}$.

$\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges by the p -series test. By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges. \square

Often easier to show a series converges than it is to find the limit.

E.g., easy to see $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, much harder to show the limit is $\pi^2/6$.

Remark: $\zeta(p) = \sum_{n=1}^{\infty} \frac{1}{n^p}$ is the “Riemann zeta function”.

Understanding its roots is one of the most famous unsolved problems in mathematics, with many applications.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$$\sum_{n=0}^{\infty} r^n \text{ is } \frac{r^{n+1}}{r^n} = r, \text{ and } \sum_{n=0}^{\infty} r^n \text{ converges whenever } r < 1.$$

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $L < 1$ (clearly $L \geq 0$).

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $L < 1$ (clearly $L \geq 0$). Pick r such that $L < r < 1$.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $L < 1$ (clearly $L \geq 0$). Pick r such that $L < r < 1$.

As $r - L > 0$, $\exists M \in \mathbb{N}$ s.t. $\forall n \geq M$, $\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L \Rightarrow \frac{|x_{n+1}|}{|x_n|} < r$.

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $L < 1$ (clearly $L \geq 0$). Pick r such that $L < r < 1$.

As $r - L > 0$, $\exists M \in \mathbb{N}$ s.t. $\forall n \geq M$, $\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L \Rightarrow \frac{|x_{n+1}|}{|x_n|} < r$.

For $n > M$ (that is for $n \geq M + 1$), write

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|}$$

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $L < 1$ (clearly $L \geq 0$). Pick r such that $L < r < 1$.

As $r - L > 0$, $\exists M \in \mathbb{N}$ s.t. $\forall n \geq M$, $\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L \Rightarrow \frac{|x_{n+1}|}{|x_n|} < r$.

For $n > M$ (that is for $n \geq M + 1$), write

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r = |x_M| r^{n-M}$$

Suppose $r > 0$. Ratio of two subsequent terms in the geometric series

$\sum_{n=0}^{\infty} r^n$ is $\frac{r^{n+1}}{r^n} = r$, and $\sum_{n=0}^{\infty} r^n$ converges whenever $r < 1$. We can generalize this.

Proposition (Ratio test)

Let $\sum_{n=1}^{\infty} x_n$ be a series, $x_n \neq 0$ for all n , and $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$ exists.

(i) If $L < 1$, then $\sum_{n=1}^{\infty} x_n$ converges absolutely. (ii) If $L > 1$, then $\sum_{n=1}^{\infty} x_n$ diverges.

Proof: If $L > 1$, by ratio test for seqs., $\{x_n\}_{n=1}^{\infty}$ diverges \Rightarrow the series cannot converge.

Suppose $L < 1$ (clearly $L \geq 0$). Pick r such that $L < r < 1$.

As $r - L > 0$, $\exists M \in \mathbb{N}$ s.t. $\forall n \geq M$, $\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L \Rightarrow \frac{|x_{n+1}|}{|x_n|} < r$.

For $n > M$ (that is for $n \geq M + 1$), write

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

For $k > M$, $\sum_{n=1}^k |x_n|$

For $k > M$,
$$\sum_{n=1}^k |x_n| = \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right)$$

$$\begin{aligned} \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\ &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) \end{aligned}$$

$$\begin{aligned}
 \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
 &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
 \end{aligned}$$

$$\begin{aligned}
 \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
 &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
 \end{aligned}$$

As $0 < r < 1$, $\sum_{n=0}^{\infty} r^n$ converges

$$\begin{aligned}
 \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
 &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
 \end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

$$\begin{aligned}
 \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
 &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
 \end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\begin{aligned}
 \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
 &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
 \end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\sum_{n=1}^k |x_n| < \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right)$$

$$\begin{aligned}
\text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
&< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
\end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\sum_{n=1}^k |x_n| < \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right) \leq \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^{\infty} r^n \right).$$

$$\begin{aligned}
 \text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\
 &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).
 \end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\sum_{n=1}^k |x_n| < \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right) \leq \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^{\infty} r^n \right).$$

The RHS does not depend on k

$$\begin{aligned}\text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\ &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).\end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\sum_{n=1}^k |x_n| < \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right) \leq \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^{\infty} r^n \right).$$

The RHS does not depend on $k \quad \Rightarrow \quad$ partial sums of $\sum_{n=1}^{\infty} |x_n|$ are bounded

$$\begin{aligned}\text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\ &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).\end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\sum_{n=1}^k |x_n| < \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right) \leq \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^{\infty} r^n \right).$$

The RHS does not depend on $k \quad \Rightarrow \quad$ partial sums of $\sum_{n=1}^{\infty} |x_n|$ are bounded

$$\Rightarrow \sum_{n=1}^{\infty} |x_n| \text{ converges}$$

$$\begin{aligned}\text{For } k > M, \quad \sum_{n=1}^k |x_n| &= \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k |x_n| \right) \\ &< \left(\sum_{n=1}^M |x_n| \right) + \left(\sum_{n=M+1}^k (|x_M| r^{-M}) r^n \right) = \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right).\end{aligned}$$

$$\text{As } 0 < r < 1, \quad \sum_{n=0}^{\infty} r^n \text{ converges} \quad \Rightarrow \quad \sum_{n=M+1}^{\infty} r^n \text{ converges.}$$

Take limit as $k \rightarrow \infty$.

$$\sum_{n=1}^k |x_n| < \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^k r^n \right) \leq \left(\sum_{n=1}^M |x_n| \right) + (|x_M| r^{-M}) \left(\sum_{n=M+1}^{\infty} r^n \right).$$

The RHS does not depend on $k \quad \Rightarrow \quad$ partial sums of $\sum_{n=1}^{\infty} |x_n|$ are bounded

$$\Rightarrow \sum_{n=1}^{\infty} |x_n| \text{ converges} \quad \Rightarrow \quad \sum_{n=1}^{\infty} x_n \text{ converges absolutely.}$$

□

Example:

$$\sum_{n=1}^{\infty} \frac{2^n}{n!} \quad \text{converges absolutely.}$$

Example:

$$\sum_{n=1}^{\infty} \frac{2^n}{n!} \quad \text{converges absolutely.}$$

Proof:

$$\lim_{n \rightarrow \infty} \frac{2^{(n+1)}/(n+1)!}{2^n/n!} = \lim_{n \rightarrow \infty} \frac{2}{n+1} = 0.$$

Example:

$$\sum_{n=1}^{\infty} \frac{2^n}{n!} \quad \text{converges absolutely.}$$

Proof:

$$\lim_{n \rightarrow \infty} \frac{2^{(n+1)}/(n+1)!}{2^n/n!} = \lim_{n \rightarrow \infty} \frac{2}{n+1} = 0.$$

Therefore, the series converges absolutely by the ratio test.



If $\sum_{n=1}^{\infty} x_n$ converges, then $x_n \rightarrow 0$.

If $\sum_{n=1}^{\infty} x_n$ converges, then $x_n \rightarrow 0$.

But $x_n \rightarrow 0$ is not enough to get the converse. It has to go to zero fast enough (let's stick with series with positive terms, it's complicated if negative terms are involved).

If $\sum_{n=1}^{\infty} x_n$ converges, then $x_n \rightarrow 0$.

But $x_n \rightarrow 0$ is not enough to get the converse. It has to go to zero fast enough (let's stick with series with positive terms, it's complicated if negative terms are involved).

One can prove:

Exercise: Suppose $\{x_n\}_{n=1}^{\infty}$ is a decreasing sequence and $\sum_{n=1}^{\infty} x_n$ converges.

Prove $\lim_{n \rightarrow \infty} nx_n = 0$.

If $\sum_{n=1}^{\infty} x_n$ converges, then $x_n \rightarrow 0$.

But $x_n \rightarrow 0$ is not enough to get the converse. It has to go to zero fast enough (let's stick with series with positive terms, it's complicated if negative terms are involved).

One can prove:

Exercise: Suppose $\{x_n\}_{n=1}^{\infty}$ is a decreasing sequence and $\sum_{n=1}^{\infty} x_n$ converges.

Prove $\lim_{n \rightarrow \infty} nx_n = 0$.

Converse still doesn't hold: $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$ diverges even though $\lim_{n \rightarrow \infty} n \left(\frac{1}{n \ln n} \right) = 0$.