

# BA: 2.2

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .  
Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ ,

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .



## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$  and  $b_n < x + \epsilon$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$  and  $b_n < x + \epsilon$ .

$\Rightarrow x - \epsilon < a_n \leq x_n \leq b_n < x + \epsilon$ .

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$  and  $b_n < x + \epsilon$ .

$\Rightarrow x - \epsilon < a_n \leq x_n \leq b_n < x + \epsilon$ .

$\Rightarrow -\epsilon < x_n - x < \epsilon$

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$  and  $b_n < x + \epsilon$ .

$\Rightarrow x - \epsilon < a_n \leq x_n \leq b_n < x + \epsilon$ .

$\Rightarrow -\epsilon < x_n - x < \epsilon \Rightarrow |x_n - x| < \epsilon$

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.

Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$  and  $b_n < x + \epsilon$ .

$\Rightarrow x - \epsilon < a_n \leq x_n \leq b_n < x + \epsilon$ .

$\Rightarrow -\epsilon < x_n - x < \epsilon \Rightarrow |x_n - x| < \epsilon \Rightarrow \{x_n\}_{n=1}^{\infty}$  converges to  $x$ . □

## Lemma (Squeeze lemma)

Let  $\{a_n\}_{n=1}^{\infty}$ ,  $\{b_n\}_{n=1}^{\infty}$ , and  $\{x_n\}_{n=1}^{\infty}$  be sequences such that  $a_n \leq x_n \leq b_n \quad \forall n \in \mathbb{N}$ .

Suppose  $\{a_n\}_{n=1}^{\infty}$  and  $\{b_n\}_{n=1}^{\infty}$  converge and  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$ . Let  $\epsilon > 0$  be given.


Find  $M_1$  s.t.  $\forall n \geq M_1, |a_n - x| < \epsilon$ , and  $M_2$  s.t.  $\forall n \geq M_2, |b_n - x| < \epsilon$ .

Set  $M := \max\{M_1, M_2\}$ .

$n \geq M \Rightarrow x - a_n < \epsilon$  or  $x - \epsilon < a_n$  and  $b_n < x + \epsilon$ .

$\Rightarrow x - \epsilon < a_n \leq x_n \leq b_n < x + \epsilon$ .

$\Rightarrow -\epsilon < x_n - x < \epsilon \Rightarrow |x_n - x| < \epsilon \Rightarrow \{x_n\}_{n=1}^{\infty}$  converges to  $x$ . □





**Example:** Consider  $\left\{\frac{1}{n\sqrt{n}}\right\}_{n=1}^{\infty}$ .

**Example:** Consider  $\left\{\frac{1}{n\sqrt{n}}\right\}_{n=1}^{\infty}$ .

As  $\sqrt{n} \geq 1$  for all  $n \in \mathbb{N}$ ,

$$0 \leq \frac{1}{n\sqrt{n}} \leq \frac{1}{n} \quad \text{for all } n \in \mathbb{N}.$$

**Example:** Consider  $\left\{\frac{1}{n\sqrt{n}}\right\}_{n=1}^{\infty}$ .

As  $\sqrt{n} \geq 1$  for all  $n \in \mathbb{N}$ ,

$$0 \leq \frac{1}{n\sqrt{n}} \leq \frac{1}{n} \quad \text{for all } n \in \mathbb{N}.$$

We already know  $\lim_{n \rightarrow \infty} 1/n = 0$ .

**Example:** Consider  $\left\{\frac{1}{n\sqrt{n}}\right\}_{n=1}^{\infty}$ .

As  $\sqrt{n} \geq 1$  for all  $n \in \mathbb{N}$ ,

$$0 \leq \frac{1}{n\sqrt{n}} \leq \frac{1}{n} \quad \text{for all } n \in \mathbb{N}.$$

We already know  $\lim_{n \rightarrow \infty} 1/n = 0$ .

The constant sequence  $\{0\}_{n=1}^{\infty}$  and  $\{1/n\}_{n=1}^{\infty}$  in the squeeze lemma  $\Rightarrow$

$$\lim_{n \rightarrow \infty} \frac{1}{n\sqrt{n}} = 0.$$

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .

Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.



## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$x_n \leq y_n \forall n$

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$$x_n \leq y_n \quad \forall n \quad \Rightarrow \quad 0 \leq y_n - x_n \quad \forall n$$

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$$x_n \leq y_n \quad \forall n \quad \Rightarrow \quad 0 \leq y_n - x_n \quad \forall n \quad \Rightarrow \quad 0 < y - x + \epsilon$$



## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$x_n \leq y_n \forall n \Rightarrow 0 \leq y_n - x_n \forall n \Rightarrow 0 < y - x + \epsilon \Rightarrow x - y < \epsilon$ .

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$x_n \leq y_n \forall n \Rightarrow 0 \leq y_n - x_n \forall n \Rightarrow 0 < y - x + \epsilon \Rightarrow x - y < \epsilon$ .

$\epsilon > 0$  is arbitrary

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$x_n \leq y_n \forall n \Rightarrow 0 \leq y_n - x_n \forall n \Rightarrow 0 < y - x + \epsilon \Rightarrow x - y < \epsilon$ .

$\epsilon > 0$  is arbitrary  $\Rightarrow x - y \leq 0$

## Lemma

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences and  $x_n \leq y_n$  for all  $n \in \mathbb{N}$ .  
Then  $\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$ .

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$  and  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $|x_n - x| < \epsilon/2$  for all  $n \geq M_1$ ,  
and  $M_2$  such that  $|y_n - y| < \epsilon/2$  for all  $n \geq M_2$ .

So for some  $n \geq \max\{M_1, M_2\}$ ,  $x - x_n < \epsilon/2$  and  $y_n - y < \epsilon/2$ .

Add these to get  $y_n - x_n + x - y < \epsilon$  or  $y_n - x_n < y - x + \epsilon$ .

$x_n \leq y_n \forall n \Rightarrow 0 \leq y_n - x_n \forall n \Rightarrow 0 < y - x + \epsilon \Rightarrow x - y < \epsilon$ .

$\epsilon > 0$  is arbitrary  $\Rightarrow x - y \leq 0 \Rightarrow x \leq y$ .



## Corollary

(i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .

## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

Proof is an exercise.

## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

Proof is an exercise.

The results are not true with strict inequalities.



## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

Proof is an exercise.

The results are not true with strict inequalities.

**Example:** Let  $x_n := -1/n$  and  $y_n := 1/n$ .

## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

Proof is an exercise.

The results are not true with strict inequalities.

**Example:** Let  $x_n := -1/n$  and  $y_n := 1/n$ .

Then  $x_n < y_n$ ,  $x_n < 0$ , and  $y_n > 0$  for all  $n$ .

## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

Proof is an exercise.

The results are not true with strict inequalities.

**Example:** Let  $x_n := -1/n$  and  $y_n := 1/n$ .

Then  $x_n < y_n$ ,  $x_n < 0$ , and  $y_n > 0$  for all  $n$ .

However,  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n = 0$ .

## Corollary

- (i) If  $\{x_n\}_{n=1}^{\infty}$  is convergent such that  $x_n \geq 0 \forall n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} x_n \geq 0$ .
- (ii) Let  $a, b \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  be convergent such that  $a \leq x_n \leq b \forall n \in \mathbb{N}$ .  
Then  $a \leq \lim_{n \rightarrow \infty} x_n \leq b$ .

Proof is an exercise.

The results are not true with strict inequalities.

**Example:** Let  $x_n := -1/n$  and  $y_n := 1/n$ .

Then  $x_n < y_n$ ,  $x_n < 0$ , and  $y_n > 0$  for all  $n$ .

However,  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n = 0$ .

Strict inequalities may become non-strict inequalities when limits are applied.

## Proposition

*Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.*

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.

(i)  $\{x_n + y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.

(i)  $\{x_n + y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$ .

(ii)  $\{x_n - y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n - y_n) = \lim_{n \rightarrow \infty} x_n - \lim_{n \rightarrow \infty} y_n$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.

- (i)  $\{x_n + y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$ .
- (ii)  $\{x_n - y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n - y_n) = \lim_{n \rightarrow \infty} x_n - \lim_{n \rightarrow \infty} y_n$ .
- (iii)  $\{x_n y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n y_n) = \left( \lim_{n \rightarrow \infty} x_n \right) \left( \lim_{n \rightarrow \infty} y_n \right)$ .



## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.

- (i)  $\{x_n + y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$ .
- (ii)  $\{x_n - y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n - y_n) = \lim_{n \rightarrow \infty} x_n - \lim_{n \rightarrow \infty} y_n$ .
- (iii)  $\{x_n y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n y_n) = \left( \lim_{n \rightarrow \infty} x_n \right) \left( \lim_{n \rightarrow \infty} y_n \right)$ .
- (iv) If  $\lim_{n \rightarrow \infty} y_n \neq 0$  and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\left\{ \frac{x_n}{y_n} \right\}_{n=1}^{\infty}$  converges and
$$\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \frac{\lim_{n \rightarrow \infty} x_n}{\lim_{n \rightarrow \infty} y_n}.$$

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.

- (i)  $\{x_n + y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$ .
- (ii)  $\{x_n - y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n - y_n) = \lim_{n \rightarrow \infty} x_n - \lim_{n \rightarrow \infty} y_n$ .
- (iii)  $\{x_n y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n y_n) = \left( \lim_{n \rightarrow \infty} x_n \right) \left( \lim_{n \rightarrow \infty} y_n \right)$ .
- (iv) If  $\lim_{n \rightarrow \infty} y_n \neq 0$  and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\left\{ \frac{x_n}{y_n} \right\}_{n=1}^{\infty}$  converges and
$$\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \frac{\lim_{n \rightarrow \infty} x_n}{\lim_{n \rightarrow \infty} y_n}.$$

Note that you can also use constant sequences.

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  be convergent sequences.

- (i)  $\{x_n + y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$ .
- (ii)  $\{x_n - y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n - y_n) = \lim_{n \rightarrow \infty} x_n - \lim_{n \rightarrow \infty} y_n$ .
- (iii)  $\{x_n y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} (x_n y_n) = \left( \lim_{n \rightarrow \infty} x_n \right) \left( \lim_{n \rightarrow \infty} y_n \right)$ .
- (iv) If  $\lim_{n \rightarrow \infty} y_n \neq 0$  and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\left\{ \frac{x_n}{y_n} \right\}_{n=1}^{\infty}$  converges and
$$\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \frac{\lim_{n \rightarrow \infty} x_n}{\lim_{n \rightarrow \infty} y_n}.$$

Note that you can also use constant sequences.

E.g., if  $c \in \mathbb{R}$  and  $\{x_n\}_{n=1}^{\infty}$  converges, then

$$\lim_{n \rightarrow \infty} c x_n = c \left( \lim_{n \rightarrow \infty} x_n \right), \quad \lim_{n \rightarrow \infty} (c + x_n) = c + \lim_{n \rightarrow \infty} x_n, \quad \text{etc.}$$

**Proof:** (i) (addition).

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .



**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n + y_n) - (x + y)|$$

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n + y_n) - (x + y)| = |x_n - x + y_n - y|$$

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n + y_n) - (x + y)| = |x_n - x + y_n - y| \leq |x_n - x| + |y_n - y|$$

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n + y_n) - (x + y)| = |x_n - x + y_n - y| \leq |x_n - x| + |y_n - y| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n + y_n) - (x + y)| = |x_n - x + y_n - y| \leq |x_n - x| + |y_n - y| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

(i) is proved.

**Proof:** (i) (addition).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent,  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \epsilon/2$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \epsilon/2$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n + y_n) - (x + y)| = |x_n - x + y_n - y| \leq |x_n - x| + |y_n - y| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

(i) is proved.

Proof of (ii) (subtraction) is almost identical (exercise).



Next (iii) (multiplication).

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given.

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,



Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n y_n) - (xy)|$$

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$|(x_n y_n) - (xy)| = |(x_n - x + x)(y_n - y + y) - xy|$$

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$\begin{aligned} |(x_n y_n) - (xy)| &= |(x_n - x + x)(y_n - y + y) - xy| \\ &= |(x_n - x)y + x(y_n - y) + (x_n - x)(y_n - y)| \end{aligned}$$

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$\begin{aligned} |(x_n y_n) - (xy)| &= |(x_n - x + x)(y_n - y + y) - xy| \\ &= |(x_n - x)y + x(y_n - y) + (x_n - x)(y_n - y)| \\ &\leq |(x_n - x)y| + |x(y_n - y)| + |(x_n - x)(y_n - y)| \end{aligned}$$

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$\begin{aligned} |(x_n y_n) - (xy)| &= |(x_n - x + x)(y_n - y + y) - xy| \\ &= |(x_n - x)y + x(y_n - y) + (x_n - x)(y_n - y)| \\ &\leq |(x_n - x)y| + |x(y_n - y)| + |(x_n - x)(y_n - y)| \\ &= |x_n - x| |y| + |x| |y_n - y| + |x_n - x| |y_n - y| \end{aligned}$$

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n, y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$\begin{aligned} |(x_n y_n) - (xy)| &= |(x_n - x + x)(y_n - y + y) - xy| \\ &= |(x_n - x)y + x(y_n - y) + (x_n - x)(y_n - y)| \\ &\leq |(x_n - x)y| + |x(y_n - y)| + |(x_n - x)(y_n - y)| \\ &= |x_n - x| |y| + |x| |y_n - y| + |x_n - x| |y_n - y| \\ &< \frac{\epsilon}{3K} K + K \frac{\epsilon}{3K} + \frac{\epsilon}{3K} \frac{\epsilon}{3K} \quad (\text{now notice that } \frac{\epsilon}{3K} \leq 1 \text{ and } K \geq 1) \end{aligned}$$

Next (iii) (multiplication).

Suppose  $\{x_n\}_{n=1}^{\infty}$  and  $\{y_n\}_{n=1}^{\infty}$  are convergent and  $x := \lim_{n \rightarrow \infty} x_n$ ,  $y := \lim_{n \rightarrow \infty} y_n$ .

Let  $\epsilon > 0$  be given. Let  $K := \max\{|x|, |y|, \epsilon/3, 1\}$ .

Find an  $M_1$  such that  $\forall n \geq M_1, |x_n - x| < \frac{\epsilon}{3K}$ .

Find an  $M_2$  such that  $\forall n \geq M_2, |y_n - y| < \frac{\epsilon}{3K}$ .

Take  $M := \max\{M_1, M_2\}$ .

For all  $n \geq M$ ,

$$\begin{aligned} |(x_n y_n) - (xy)| &= |(x_n - x + x)(y_n - y + y) - xy| \\ &= |(x_n - x)y + x(y_n - y) + (x_n - x)(y_n - y)| \\ &\leq |(x_n - x)y| + |x(y_n - y)| + |(x_n - x)(y_n - y)| \\ &= |x_n - x| |y| + |x| |y_n - y| + |x_n - x| |y_n - y| \\ &< \frac{\epsilon}{3K} K + K \frac{\epsilon}{3K} + \frac{\epsilon}{3K} \frac{\epsilon}{3K} \quad (\text{now notice that } \frac{\epsilon}{3K} \leq 1 \text{ and } K \geq 1) \\ &\leq \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon \end{aligned}$$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

and  $\lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}$ .



(iv) follows from (iii) and the claim:

**Claim:** *If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}$ .*

**Proof of claim:** Let  $\epsilon > 0$  be given.

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}$ .

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}$ .

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

$$\left| \frac{1}{y_n} - \frac{1}{y} \right|$$



(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| = \left| \frac{y - y_n}{yy_n} \right|$$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| = \left| \frac{y - y_n}{y y_n} \right| = \frac{|y - y_n|}{|y| |y_n|}$$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| = \left| \frac{y - y_n}{y y_n} \right| = \frac{|y - y_n|}{|y| |y_n|} \leq \frac{|y - y_n|}{|y|} \frac{2}{|y|}$$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| = \left| \frac{y - y_n}{y y_n} \right| = \frac{|y - y_n|}{|y| |y_n|} \leq \frac{|y - y_n|}{|y|} \frac{2}{|y|} < \frac{|y|^2 \frac{\epsilon}{2}}{|y|} \frac{2}{|y|}$$

(iv) follows from (iii) and the claim:

**Claim:** If  $\{y_n\}_{n=1}^{\infty}$  is convergent,  $\lim_{n \rightarrow \infty} y_n \neq 0$ , and  $y_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\{1/y_n\}_{n=1}^{\infty}$  converges

$$\text{and } \lim_{n \rightarrow \infty} \frac{1}{y_n} = \frac{1}{\lim_{n \rightarrow \infty} y_n}.$$

**Proof of claim:** Let  $\epsilon > 0$  be given.

Let  $y := \lim_{n \rightarrow \infty} y_n$ .

$$|y| \neq 0 \quad \Rightarrow \quad \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\} > 0.$$

Find an  $M$  such that for all  $n \geq M$ ,  $|y_n - y| < \min \left\{ |y|^2 \frac{\epsilon}{2}, \frac{|y|}{2} \right\}$ .

For  $n \geq M$ ,  $|y - y_n| < |y|/2$ , so  $|y| = |y - y_n + y_n| \leq |y - y_n| + |y_n| < \frac{|y|}{2} + |y_n|$ .

Subtract  $|y|/2$  from both sides to get  $|y|/2 < |y_n|$ , or  $\frac{1}{|y_n|} < \frac{2}{|y|}$

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| = \left| \frac{y - y_n}{y y_n} \right| = \frac{|y - y_n|}{|y| |y_n|} \leq \frac{|y - y_n|}{|y|} \frac{2}{|y|} < \frac{|y|^2 \frac{\epsilon}{2}}{|y|} \frac{2}{|y|} = \epsilon.$$

□

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.



**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon$ .

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon$ .

Now suppose  $x > 0$  (and hence  $\sqrt{x} > 0$ ).

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon.$$

Now suppose  $x > 0$  (and hence  $\sqrt{x} > 0$ ).

$$|\sqrt{x_n} - \sqrt{x}|$$



**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon.$$

Now suppose  $x > 0$  (and hence  $\sqrt{x} > 0$ ).

$$|\sqrt{x_n} - \sqrt{x}| = \left| \frac{x_n - x}{\sqrt{x_n} + \sqrt{x}} \right|$$

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left(\lim_{n \rightarrow \infty} x_n\right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon.$$

Now suppose  $x > 0$  (and hence  $\sqrt{x} > 0$ ).

$$|\sqrt{x_n} - \sqrt{x}| = \left| \frac{x_n - x}{\sqrt{x_n} + \sqrt{x}} \right| = \frac{1}{\sqrt{x_n} + \sqrt{x}} |x_n - x|$$

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon$ .

Now suppose  $x > 0$  (and hence  $\sqrt{x} > 0$ ).

$$|\sqrt{x_n} - \sqrt{x}| = \left| \frac{x_n - x}{\sqrt{x_n} + \sqrt{x}} \right| = \frac{1}{\sqrt{x_n} + \sqrt{x}} |x_n - x| \leq \frac{1}{\sqrt{x}} |x_n - x|.$$

**Exercise:** (induction)  $\lim_{n \rightarrow \infty} x_n^k = \left( \lim_{n \rightarrow \infty} x_n \right)^k$  for all  $k \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be convergent and  $x_n \geq 0$  for all  $n \in \mathbb{N}$ . Then  $\lim_{n \rightarrow \infty} \sqrt{x_n} = \sqrt{\lim_{n \rightarrow \infty} x_n}$ .

Notice  $\lim_{n \rightarrow \infty} x_n \geq 0$  so the square root makes sense.

**Proof:** Let  $x := \lim_{n \rightarrow \infty} x_n$ ,  $x \geq 0$ .

First suppose  $x = 0$ .

Let  $\epsilon > 0$  be given.

Find  $M$  such that  $\forall n \geq M$ ,  $x_n = |x_n| < \epsilon^2$ , or  $\sqrt{x_n} < \epsilon$ .

$$\Rightarrow |\sqrt{x_n} - \sqrt{x}| = \sqrt{x_n} < \epsilon.$$

Now suppose  $x > 0$  (and hence  $\sqrt{x} > 0$ ).

$$|\sqrt{x_n} - \sqrt{x}| = \left| \frac{x_n - x}{\sqrt{x_n} + \sqrt{x}} \right| = \frac{1}{\sqrt{x_n} + \sqrt{x}} |x_n - x| \leq \frac{1}{\sqrt{x}} |x_n - x|.$$

The rest is an exercise. □

## Proposition

*If  $\{x_n\}_{n=1}^{\infty}$  is a convergent sequence, then  $\{|x_n|\}_{n=1}^{\infty}$  is convergent and*

$$\lim_{n \rightarrow \infty} |x_n| = \left| \lim_{n \rightarrow \infty} x_n \right|.$$

## Proposition

If  $\{x_n\}_{n=1}^{\infty}$  is a convergent sequence, then  $\{|x_n|\}_{n=1}^{\infty}$  is convergent and

$$\lim_{n \rightarrow \infty} |x_n| = \left| \lim_{n \rightarrow \infty} x_n \right|.$$

**Proof sketch:** The reverse triangle inequality:  $\left| |x_n| - |x| \right| \leq |x_n - x|.$



**Example:**

Note that  $\lim_{n \rightarrow \infty} 1/n = 0$ .

**Example:**

Note that  $\lim_{n \rightarrow \infty} 1/n = 0$ .

Then

$$\lim_{n \rightarrow \infty} \left| \sqrt{1 + 1/n} - 100/n^2 \right| = \left| \sqrt{1 + \left( \lim_{n \rightarrow \infty} 1/n \right)} - 100 \left( \lim_{n \rightarrow \infty} 1/n \right) \left( \lim_{n \rightarrow \infty} 1/n \right) \right| = 1.$$



**Example:**

Note that  $\lim_{n \rightarrow \infty} 1/n = 0$ .

Then

$$\lim_{n \rightarrow \infty} \left| \sqrt{1 + 1/n} - 100/n^2 \right| = \left| \sqrt{1 + \left( \lim_{n \rightarrow \infty} 1/n \right)} - 100 \left( \lim_{n \rightarrow \infty} 1/n \right) \left( \lim_{n \rightarrow \infty} 1/n \right) \right| = 1.$$

Read this from right to left. The propositions  $\Rightarrow$  LHS exists.

**Example:**

Note that  $\lim_{n \rightarrow \infty} 1/n = 0$ .

Then

$$\lim_{n \rightarrow \infty} \left| \sqrt{1 + 1/n} - 100/n^2 \right| = \left| \sqrt{1 + \left( \lim_{n \rightarrow \infty} 1/n \right)} - 100 \left( \lim_{n \rightarrow \infty} 1/n \right) \left( \lim_{n \rightarrow \infty} 1/n \right) \right| = 1.$$

Read this from right to left. The propositions  $\Rightarrow$  LHS exists.

**Example:** Be careful:

$$\lim_{n \rightarrow \infty} \left( \frac{n^2}{n+1} - n \right) = -1,$$

**Example:**

Note that  $\lim_{n \rightarrow \infty} 1/n = 0$ .

Then

$$\lim_{n \rightarrow \infty} \left| \sqrt{1 + 1/n} - 100/n^2 \right| = \left| \sqrt{1 + \left( \lim_{n \rightarrow \infty} 1/n \right)} - 100 \left( \lim_{n \rightarrow \infty} 1/n \right) \left( \lim_{n \rightarrow \infty} 1/n \right) \right| = 1.$$

Read this from right to left. The propositions  $\Rightarrow$  LHS exists.

**Example:** Be careful:

$$\lim_{n \rightarrow \infty} \left( \frac{n^2}{n+1} - n \right) = -1, \quad \text{but} \quad \left( \lim_{n \rightarrow \infty} \frac{n^2}{n+1} \right) - \left( \lim_{n \rightarrow \infty} n \right) \text{ is nonsense.}$$

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

1) the sequence is well-defined,

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.



**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}$$

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n}$$

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n} = \frac{x_n^2 + 2}{2x_n}.$$

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n} = \frac{x_n^2 + 2}{2x_n}.$$

$$x_n^2 + 2 > 0 \text{ and } x_n > 0,$$

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n} = \frac{x_n^2 + 2}{2x_n}.$$

$x_n^2 + 2 > 0$  and  $x_n > 0$ , and so  $x_{n+1} = \frac{x_n^2 + 2}{2x_n} > 0$ .

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n} = \frac{x_n^2 + 2}{2x_n}.$$

$x_n^2 + 2 > 0$  and  $x_n > 0$ , and so  $x_{n+1} = \frac{x_n^2 + 2}{2x_n} > 0$ .

By induction  $\{x_n\}_{n=1}^{\infty}$  exists and  $x_n > 0$  for all  $n$ .



**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n} = \frac{x_n^2 + 2}{2x_n}.$$

$x_n^2 + 2 > 0$  and  $x_n > 0$ , and so  $x_{n+1} = \frac{x_n^2 + 2}{2x_n} > 0$ .

By induction  $\{x_n\}_{n=1}^{\infty}$  exists and  $x_n > 0$  for all  $n$ .

We claim  $\{x_n\}_{n=1}^{\infty}$  is monotone decreasing.

**Example:** Define  $\{x_n\}_{n=1}^{\infty}$  by  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ .

We must prove:

- 1) the sequence is well-defined,
- 2) the sequence converges,
- 3) only then try to find the limit.

First,  $x_1 = 2 > 0$  (so  $x_2$  exists:  $x_2 = 2 - \frac{2^2-2}{2 \cdot 2} = 1.5 > 0$ )

Suppose for some  $n$ ,  $x_n$  exists and  $x_n > 0$ .

$$x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n} = \frac{2x_n^2 - x_n^2 + 2}{2x_n} = \frac{x_n^2 + 2}{2x_n}.$$

$x_n^2 + 2 > 0$  and  $x_n > 0$ , and so  $x_{n+1} = \frac{x_n^2 + 2}{2x_n} > 0$ .

By induction  $\{x_n\}_{n=1}^{\infty}$  exists and  $x_n > 0$  for all  $n$ .

We claim  $\{x_n\}_{n=1}^{\infty}$  is monotone decreasing.

If we show that  $x_n^2 - 2 \geq 0$  for all  $n$ , then  $x_{n+1} \leq x_n$  for all  $n$ .

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2}$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2}$$



(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2}$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^{\infty}$  is monotone decreasing and bounded below, it must converge.

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^{\infty}$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^\infty$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

$\{x_{n+1}\}_{n=1}^\infty$  is the 1-tail of  $\{x_n\}_{n=1}^\infty$ , so it converges to the same limit, say  $x$ .

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^\infty$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

$\{x_{n+1}\}_{n=1}^\infty$  is the 1-tail of  $\{x_n\}_{n=1}^\infty$ , so it converges to the same limit, say  $x$ . So

$$2x^2 = x^2 + 2$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^\infty$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

$\{x_{n+1}\}_{n=1}^\infty$  is the 1-tail of  $\{x_n\}_{n=1}^\infty$ , so it converges to the same limit, say  $x$ . So

$$2x^2 = x^2 + 2 \quad \Rightarrow \quad x^2 = 2.$$

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^\infty$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

$\{x_{n+1}\}_{n=1}^\infty$  is the 1-tail of  $\{x_n\}_{n=1}^\infty$ , so it converges to the same limit, say  $x$ . So

$$2x^2 = x^2 + 2 \quad \Rightarrow \quad x^2 = 2.$$

As  $x_n > 0$  for all  $n$ ,  $x \geq 0$ .



(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^\infty$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

$\{x_{n+1}\}_{n=1}^\infty$  is the 1-tail of  $\{x_n\}_{n=1}^\infty$ , so it converges to the same limit, say  $x$ . So

$$2x^2 = x^2 + 2 \quad \Rightarrow \quad x^2 = 2.$$

As  $x_n > 0$  for all  $n$ ,  $x \geq 0$ .  $\Rightarrow x = \sqrt{2}$ .

(recall:  $x_1 := 2$  and  $x_{n+1} := x_n - \frac{x_n^2 - 2}{2x_n}$ )

$$x_1^2 - 2 = 4 - 2 = 2 > 0$$

$$x_{n+1}^2 - 2 = \left( \frac{x_n^2 + 2}{2x_n} \right)^2 - 2 = \frac{x_n^4 + 4x_n^2 + 4 - 8x_n^2}{4x_n^2} = \frac{x_n^4 - 4x_n^2 + 4}{4x_n^2} = \frac{(x_n^2 - 2)^2}{4x_n^2} \geq 0.$$

So  $\{x_n\}_{n=1}^\infty$  is monotone decreasing and bounded below, it must converge.

Write

$$2x_n x_{n+1} = x_n^2 + 2.$$

$\{x_{n+1}\}_{n=1}^\infty$  is the 1-tail of  $\{x_n\}_{n=1}^\infty$ , so it converges to the same limit, say  $x$ . So

$$2x^2 = x^2 + 2 \quad \Rightarrow \quad x^2 = 2.$$

As  $x_n > 0$  for all  $n$ ,  $x \geq 0$ .  $\Rightarrow x = \sqrt{2}$ .

You may have noticed the above is *Newton's method* for finding  $\sqrt{2}$ , a common and practical way to find roots of equations.

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ .

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x$$

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x \quad \Rightarrow \quad x = 0.$$

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x \quad \Rightarrow \quad x = 0.$$

But the sequence does **not** converge (unbounded).

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x \quad \Rightarrow \quad x = 0.$$

But the sequence does **not** converge (unbounded).  $\lim_{n \rightarrow \infty} x_n$  does not exist.



**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x \quad \Rightarrow \quad x = 0.$$

But the sequence does **not** converge (unbounded).  $\lim_{n \rightarrow \infty} x_n$  does not exist.

**Moral:** Before you can compute what a limit is, you need to know the sequence converges.

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x \quad \Rightarrow \quad x = 0.$$

But the sequence does **not** converge (unbounded).  $\lim_{n \rightarrow \infty} x_n$  does not exist.

**Moral:** Before you can compute what a limit is, you need to know the sequence converges.

**More general moral:** Before you prove that a limit actually exists do not do any calculations with

$$\lim_{n \rightarrow \infty} x_n.$$

**Example:** Suppose  $x_1 := 1$  and  $x_{n+1} := x_n^2 + x_n$ .

Suppose we blindly write  $x := \lim_{n \rightarrow \infty} x_n$ . Then  $x_{n+1} = x_n^2 + x_n$  gives

$$x = x^2 + x \quad \Rightarrow \quad x = 0.$$

But the sequence does **not** converge (unbounded).  $\lim_{n \rightarrow \infty} x_n$  does not exist.

**Moral:** Before you can compute what a limit is, you need to know the sequence converges.

**More general moral:** Before you prove that a limit actually exists do not do any calculations with

$$\lim_{n \rightarrow \infty} x_n.$$

Just don't write " $\lim_{n \rightarrow \infty}$ " anywhere before you prove the limit exists.

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence. Suppose  $\exists x \in \mathbb{R}$  and a convergent  $\{a_n\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $|x_n - x| \leq a_n$  for all  $n \in \mathbb{N}$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence. Suppose  $\exists x \in \mathbb{R}$  and a convergent  $\{a_n\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $|x_n - x| \leq a_n$  for all  $n \in \mathbb{N}$ .

Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = x$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence. Suppose  $\exists x \in \mathbb{R}$  and a convergent  $\{a_n\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $|x_n - x| \leq a_n$  for all  $n \in \mathbb{N}$ .  
Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = x$ .

**Proof:** Let  $\epsilon > 0$  be given.

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence. Suppose  $\exists x \in \mathbb{R}$  and a convergent  $\{a_n\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $|x_n - x| \leq a_n$  for all  $n \in \mathbb{N}$ .  
Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = x$ .

**Proof:** Let  $\epsilon > 0$  be given.

Note  $a_n \geq 0$  for all  $n$ .

## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence. Suppose  $\exists x \in \mathbb{R}$  and a convergent  $\{a_n\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $|x_n - x| \leq a_n$  for all  $n \in \mathbb{N}$ .  
Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = x$ .

**Proof:** Let  $\epsilon > 0$  be given.

Note  $a_n \geq 0$  for all  $n$ .

Find  $M \in \mathbb{N}$  such that  $a_n = |a_n - 0| < \epsilon$  for all  $n \geq M$ .



## Proposition

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence. Suppose  $\exists x \in \mathbb{R}$  and a convergent  $\{a_n\}_{n=1}^{\infty}$  such that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $|x_n - x| \leq a_n$  for all  $n \in \mathbb{N}$ .  
Then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = x$ .

**Proof:** Let  $\epsilon > 0$  be given.

Note  $a_n \geq 0$  for all  $n$ .

Find  $M \in \mathbb{N}$  such that  $a_n = |a_n - 0| < \epsilon$  for all  $n \geq M$ .

So for all  $n \geq M$ ,  $|x_n - x| \leq a_n < \epsilon$ .



## Proposition

*Let  $c > 0$ .*

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges.



## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

Now consider  $c > 1$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

Now consider  $c > 1$ . Let  $B > 0$  be arbitrary.

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

Now consider  $c > 1$ . Let  $B > 0$  be arbitrary.

As  $1/c < 1$ , then  $\{(1/c)^n\}_{n=1}^{\infty}$  converges to 0.

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

Now consider  $c > 1$ . Let  $B > 0$  be arbitrary.

As  $1/c < 1$ , then  $\{(1/c)^n\}_{n=1}^{\infty}$  converges to 0.

$\Rightarrow$  for some large enough  $n$ ,  $\frac{1}{c^n} = \left(\frac{1}{c}\right)^n < \frac{1}{B}$ .



## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

Now consider  $c > 1$ . Let  $B > 0$  be arbitrary.

As  $1/c < 1$ , then  $\{(1/c)^n\}_{n=1}^{\infty}$  converges to 0.

$\Rightarrow$  for some large enough  $n$ ,  $\frac{1}{c^n} = \left(\frac{1}{c}\right)^n < \frac{1}{B}$ .

$\Rightarrow c^n > B$ , and  $B$  is not an upper bound for  $\{c^n\}_{n=1}^{\infty}$ .

## Proposition

Let  $c > 0$ .

- (i) If  $c < 1$ , then  $\lim_{n \rightarrow \infty} c^n = 0$ .
- (ii) If  $c > 1$ , then  $\{c^n\}_{n=1}^{\infty}$  is unbounded.

**Proof:** First consider  $c < 1$ .

As  $c > 0$ , then  $c^n > 0$  for all  $n \in \mathbb{N}$  by induction.

As  $c < 1$ , then  $c^{n+1} < c^n$  for all  $n$ .

$\Rightarrow \{c^n\}_{n=1}^{\infty}$  is decreasing bounded below  $\Rightarrow$  it converges. Let  $x := \lim_{n \rightarrow \infty} c^n$ .

The 1-tail  $\{c^{n+1}\}_{n=1}^{\infty}$  converges to  $x$ .

Take the limit of both sides of  $c^{n+1} = c \cdot c^n \Rightarrow x = cx$ , or  $(1 - c)x = 0$ .

$\Rightarrow x = 0$  as  $c \neq 1$ .

Now consider  $c > 1$ . Let  $B > 0$  be arbitrary.

As  $1/c < 1$ , then  $\{(1/c)^n\}_{n=1}^{\infty}$  converges to 0.

$\Rightarrow$  for some large enough  $n$ ,  $\frac{1}{c^n} = \left(\frac{1}{c}\right)^n < \frac{1}{B}$ .

$\Rightarrow c^n > B$ , and  $B$  is not an upper bound for  $\{c^n\}_{n=1}^{\infty}$ .

As  $B$  was arbitrary,  $\{c^n\}_{n=1}^{\infty}$  is unbounded.



## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

(i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .



## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .

$\{n\}_{n=1}^{\infty}$  is unbounded, and  $L = 1$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .

$\{n\}_{n=1}^{\infty}$  is unbounded, and  $L = 1$ .

**Proof:** Suppose  $L < 1$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .

$\{n\}_{n=1}^{\infty}$  is unbounded, and  $L = 1$ .

**Proof:** Suppose  $L < 1$ .

$$\frac{|x_{n+1}|}{|x_n|} \geq 0 \text{ for all } n \quad \Rightarrow \quad L \geq 0.$$

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .

$\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .

$\{n\}_{n=1}^{\infty}$  is unbounded, and  $L = 1$ .

**Proof:** Suppose  $L < 1$ .

$\frac{|x_{n+1}|}{|x_n|} \geq 0$  for all  $n \Rightarrow L \geq 0$ . Pick  $r$  such that  $L < r < 1$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .       $\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .  
 $\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .       $\{n\}_{n=1}^{\infty}$  is unbounded, and  $L = 1$ .

**Proof:** Suppose  $L < 1$ .

$\frac{|x_{n+1}|}{|x_n|} \geq 0$  for all  $n \Rightarrow L \geq 0$ .      Pick  $r$  such that  $L < r < 1$ .

$\frac{|x_{n+1}|}{|x_n|}$  will eventually be less than  $r$  and so we compare the sequence to  $\{r^n\}_{n=1}^{\infty}$ .

## Lemma (Ratio test for sequences)

Let  $\{x_n\}_{n=1}^{\infty}$  be such that  $x_n \neq 0$  for all  $n$  and  $L := \lim_{n \rightarrow \infty} \frac{|x_{n+1}|}{|x_n|}$  exists.

- (i) If  $L < 1$ , then  $\{x_n\}_{n=1}^{\infty}$  converges and  $\lim_{n \rightarrow \infty} x_n = 0$ .
- (ii) If  $L > 1$ , then  $\{x_n\}_{n=1}^{\infty}$  is unbounded (hence diverges).

If  $L = 1$ , no conclusion. E.g.,

$\{1/n\}_{n=1}^{\infty}$  converges to zero, but  $L = 1$ .

$\{1\}_{n=1}^{\infty}$  converges to 1, and  $L = 1$ .


$\{(-1)^n\}_{n=1}^{\infty}$  does not converge, and  $L = 1$ .

$\{n\}_{n=1}^{\infty}$  is unbounded, and  $L = 1$ .

**Proof:** Suppose  $L < 1$ .

$\frac{|x_{n+1}|}{|x_n|} \geq 0$  for all  $n \Rightarrow L \geq 0$ . Pick  $r$  such that  $L < r < 1$ .

$\frac{|x_{n+1}|}{|x_n|}$  will eventually be less than  $r$  and so we compare the sequence to  $\{r^n\}_{n=1}^{\infty}$ .



As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$



As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L$$

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

For  $n > M$  ( $n \geq M + 1$ )

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|}$$

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

For  $n > M$  ( $n \geq M + 1$ )

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r$$

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

For  $n > M$  ( $n \geq M + 1$ )

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

For  $n > M$  ( $n \geq M + 1$ )

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^{\infty}$  converges to zero

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

For  $n > M$  ( $n \geq M + 1$ )

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^{\infty}$  converges to zero  $\Rightarrow \{|x_M| r^{-M} r^n\}_{n=1}^{\infty}$  converges to zero.

As  $r - L > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$ ,

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < r - L.$$

$\Rightarrow$  for  $n \geq M$ ,

$$\frac{|x_{n+1}|}{|x_n|} - L < r - L \quad \text{or} \quad \frac{|x_{n+1}|}{|x_n|} < r.$$

For  $n > M$  ( $n \geq M + 1$ )

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} < |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^{\infty}$  converges to zero  $\Rightarrow \{|x_M| r^{-M} r^n\}_{n=1}^{\infty}$  converges to zero.

The  $M$ -tail of  $\{x_n\}_{n=1}^{\infty}$  converges to zero  $\Rightarrow \{x_n\}_{n=1}^{\infty}$  converges to zero.



Now suppose  $L > 1$ .

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|}$$

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} > |x_M| r r \cdots r$$

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} > |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} > |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^{\infty}$  is unbounded ( $r > 1$ )



Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} > |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^\infty$  is unbounded ( $r > 1$ )  $\Rightarrow \{x_n\}_{n=1}^\infty$  is not bounded

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} > |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^\infty$  is unbounded ( $r > 1$ )  $\Rightarrow \{x_n\}_{n=1}^\infty$  is not bounded

(if  $|x_n| \leq B \forall n \Rightarrow r^n < \frac{B}{|x_M|} r^M$  for all  $n > M$ ,  $\Rightarrow \Leftarrow$ ).

Now suppose  $L > 1$ . Pick  $r$  such that  $1 < r < L$ .

As  $L - r > 0$ ,  $\exists M \in \mathbb{N}$  such that for all  $n \geq M$

$$\left| \frac{|x_{n+1}|}{|x_n|} - L \right| < L - r.$$

$\Rightarrow$

$$\frac{|x_{n+1}|}{|x_n|} > r.$$

For  $n > M$ ,

$$|x_n| = |x_M| \frac{|x_{M+1}|}{|x_M|} \frac{|x_{M+2}|}{|x_{M+1}|} \cdots \frac{|x_n|}{|x_{n-1}|} > |x_M| r r \cdots r = |x_M| r^{n-M} = (|x_M| r^{-M}) r^n.$$

$\{r^n\}_{n=1}^\infty$  is unbounded ( $r > 1$ )  $\Rightarrow \{x_n\}_{n=1}^\infty$  is not bounded

(if  $|x_n| \leq B \forall n \Rightarrow r^n < \frac{B}{|x_M|} r^M$  for all  $n > M$ ,  $\Rightarrow \Leftarrow$ ).

$\Rightarrow \{x_n\}_{n=1}^\infty$  diverges.



**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0.$

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!}$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0.$

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!}$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ .

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □



**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given.

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon}$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$



**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$

$$\Rightarrow \exists M, \text{ s.t. for } n \geq M, \frac{n}{(1+\epsilon)^n} < 1,$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$

$$\Rightarrow \exists M, \text{ s.t. for } n \geq M, \frac{n}{(1+\epsilon)^n} < 1, \quad \text{or} \quad n < (1+\epsilon)^n,$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$

$$\Rightarrow \exists M, \text{ s.t. for } n \geq M, \frac{n}{(1+\epsilon)^n} < 1, \quad \text{or} \quad n < (1+\epsilon)^n, \quad \text{or} \quad n^{1/n} < 1+\epsilon.$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$

$$\Rightarrow \exists M, \text{ s.t. for } n \geq M, \frac{n}{(1+\epsilon)^n} < 1, \quad \text{or} \quad n < (1+\epsilon)^n, \quad \text{or} \quad n^{1/n} < 1+\epsilon.$$

$$n^{1/n} \geq 1 \text{ (as } n \geq 1)$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$

$$\Rightarrow \exists M, \text{ s.t. for } n \geq M, \frac{n}{(1+\epsilon)^n} < 1, \quad \text{or} \quad n < (1+\epsilon)^n, \quad \text{or} \quad n^{1/n} < 1+\epsilon.$$

$$n^{1/n} \geq 1 \text{ (as } n \geq 1) \quad \Rightarrow \quad 0 \leq n^{1/n} - 1 < \epsilon.$$

**Example:**  $\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0$ .

**Proof:**  $\frac{2^{n+1}/(n+1)!}{2^n/n!} = \frac{2^{n+1}}{2^n} \frac{n!}{(n+1)!} = \frac{2}{n+1} \rightarrow 0$ . Conclusion follows. □

**Example:**  $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .

**Proof:** Let  $\epsilon > 0$  be given. Consider  $\left\{ \frac{n}{(1+\epsilon)^n} \right\}_{n=1}^{\infty}$ .  $\frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{n+1}{n} \frac{1}{1+\epsilon}$ .

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right) = 1 \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \frac{(n+1)/(1+\epsilon)^{n+1}}{n/(1+\epsilon)^n} = \frac{1}{1+\epsilon} < 1.$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{(1+\epsilon)^n} = 0$$

$$\Rightarrow \exists M, \text{ s.t. for } n \geq M, \frac{n}{(1+\epsilon)^n} < 1, \quad \text{or} \quad n < (1+\epsilon)^n, \quad \text{or} \quad n^{1/n} < 1+\epsilon.$$

$$n^{1/n} \geq 1 \text{ (as } n \geq 1) \quad \Rightarrow \quad 0 \leq n^{1/n} - 1 < \epsilon.$$

$$\Rightarrow \lim_{n \rightarrow \infty} n^{1/n} = 1. \quad \text{□}$$