

# BA: 3.1

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

Note: A cluster point of  $S$  need not lie in  $S$ .

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

Note: A cluster point of  $S$  need not lie in  $S$ .

### Examples:

- (i)  $\{1/n : n \in \mathbb{N}\}$  has a unique cluster point: 0.

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

Note: A cluster point of  $S$  need not lie in  $S$ .

### Examples:

- (i)  $\{1/n : n \in \mathbb{N}\}$  has a unique cluster point: 0.
- (ii) The set of cluster points of  $(0, 1)$  is  $[0, 1]$ .

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

Note: A cluster point of  $S$  need not lie in  $S$ .

### Examples:

- (i)  $\{1/n : n \in \mathbb{N}\}$  has a unique cluster point: 0.
- (ii) The set of cluster points of  $(0, 1)$  is  $[0, 1]$ .
- (iii) The set of cluster points of  $\mathbb{Q}$  is  $\mathbb{R}$ .

## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

Note: A cluster point of  $S$  need not lie in  $S$ .

### Examples:

- (i)  $\{1/n : n \in \mathbb{N}\}$  has a unique cluster point: 0.
- (ii) The set of cluster points of  $(0, 1)$  is  $[0, 1]$ .
- (iii) The set of cluster points of  $\mathbb{Q}$  is  $\mathbb{R}$ .
- (iv) The set of cluster points of  $[0, 1) \cup \{2\}$  is  $[0, 1]$ .



## Definition

Let  $S \subset \mathbb{R}$  be a set. A number  $x \in \mathbb{R}$  is called a *cluster point* of  $S$  if for every  $\epsilon > 0$ , the set  $(x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$  is not empty.

I.e.,  $x \in \mathbb{R}$  is a cluster point of  $S$  if  $\forall \epsilon > 0 \exists y \in S$  s.t.  $y \neq x$  and  $|x - y| < \epsilon$ .

Note: A cluster point of  $S$  need not lie in  $S$ .

### Examples:

- (i)  $\{1/n : n \in \mathbb{N}\}$  has a unique cluster point: 0.
- (ii) The set of cluster points of  $(0, 1)$  is  $[0, 1]$ .
- (iii) The set of cluster points of  $\mathbb{Q}$  is  $\mathbb{R}$ .
- (iv) The set of cluster points of  $[0, 1) \cup \{2\}$  is  $[0, 1]$ .
- (v)  $\mathbb{N}$  has no cluster points.

## Proposition

*Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .*

## Proposition

*Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .*

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

## Proposition

*Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .*

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

For every  $n \in \mathbb{N}$ , pick  $x_n$  to be an arbitrary point of  $(x - 1/n, x + 1/n) \cap S \setminus \{x\}$ .

( $x_n$  exists as  $x$  is a cluster point of  $S$ ).

## Proposition

*Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .*

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

For every  $n \in \mathbb{N}$ , pick  $x_n$  to be an arbitrary point of  $(x - 1/n, x + 1/n) \cap S \setminus \{x\}$ .

( $x_n$  exists as  $x$  is a cluster point of  $S$ ).

$x_n$  is within  $1/n$  of  $x$ :  $|x - x_n| < 1/n$ .

## Proposition

*Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .*

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

For every  $n \in \mathbb{N}$ , pick  $x_n$  to be an arbitrary point of  $(x - 1/n, x + 1/n) \cap S \setminus \{x\}$ .

( $x_n$  exists as  $x$  is a cluster point of  $S$ ).

$x_n$  is within  $1/n$  of  $x$ :  $|x - x_n| < 1/n$ .

$\{1/n\}_{n=1}^{\infty}$  converges to zero,  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

## Proposition

*Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .*

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

For every  $n \in \mathbb{N}$ , pick  $x_n$  to be an arbitrary point of  $(x - 1/n, x + 1/n) \cap S \setminus \{x\}$ .

( $x_n$  exists as  $x$  is a cluster point of  $S$ ).

$x_n$  is within  $1/n$  of  $x$ :  $|x - x_n| < 1/n$ .

$\{1/n\}_{n=1}^{\infty}$  converges to zero,  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

On the other hand,

suppose  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{x\}$  converging to  $x$ .

## Proposition

Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

For every  $n \in \mathbb{N}$ , pick  $x_n$  to be an arbitrary point of  $(x - 1/n, x + 1/n) \cap S \setminus \{x\}$ .

( $x_n$  exists as  $x$  is a cluster point of  $S$ ).

$x_n$  is within  $1/n$  of  $x$ :  $|x - x_n| < 1/n$ .

$\{1/n\}_{n=1}^{\infty}$  converges to zero,  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

On the other hand,

suppose  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{x\}$  converging to  $x$ .

$\Rightarrow$  for every  $\epsilon > 0$   $\exists M$  such that  $|x_M - x| < \epsilon$ .



## Proposition

Let  $S \subset \mathbb{R}$ . Then  $x \in \mathbb{R}$  is a cluster point of  $S$  if and only if there exists a convergent sequence of numbers  $\{x_n\}_{n=1}^{\infty}$  such that  $x_n \neq x$  and  $x_n \in S$  for all  $n$ , and  $\lim_{n \rightarrow \infty} x_n = x$ .

**Proof:** Suppose  $x$  is a cluster point of  $S$ .

For every  $n \in \mathbb{N}$ , pick  $x_n$  to be an arbitrary point of  $(x - 1/n, x + 1/n) \cap S \setminus \{x\}$ .

( $x_n$  exists as  $x$  is a cluster point of  $S$ ).

$x_n$  is within  $1/n$  of  $x$ :  $|x - x_n| < 1/n$ .

$\{1/n\}_{n=1}^{\infty}$  converges to zero,  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

On the other hand,

suppose  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{x\}$  converging to  $x$ .

$\Rightarrow$  for every  $\epsilon > 0$   $\exists M$  such that  $|x_M - x| < \epsilon$ .

$\Rightarrow x_M \in (x - \epsilon, x + \epsilon) \cap S \setminus \{x\}$ .



## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

We say  $L$  is the *limit* of  $f(x)$  as  $x$  goes to  $c$ .

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

We say  $L$  is the *limit* of  $f(x)$  as  $x$  goes to  $c$ .

We write  $\lim_{x \rightarrow c} f(x) := L$ , or  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

We say  $L$  is the *limit* of  $f(x)$  as  $x$  goes to  $c$ .

We write  $\lim_{x \rightarrow c} f(x) := L$ , or  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

If no such  $L$  exists, we say the limit does not exist or that  $f$  *diverges* at  $c$ .

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

We say  $L$  is the *limit* of  $f(x)$  as  $x$  goes to  $c$ .

We write  $\lim_{x \rightarrow c} f(x) := L$ , or  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

If no such  $L$  exists, we say the limit does not exist or that  $f$  *diverges* at  $c$ .

**Cheating again:** The notation assumes the limit is unique,  
we'll prove that momentarily.



## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

We say  $L$  is the *limit* of  $f(x)$  as  $x$  goes to  $c$ .

We write  $\lim_{x \rightarrow c} f(x) := L$ , or  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

If no such  $L$  exists, we say the limit does not exist or that  $f$  *diverges* at  $c$ .

**Cheating again:** The notation assumes the limit is unique, we'll prove that momentarily.

**Remark:** It is irrelevant whether  $f(c)$  is defined or not.

## Definition

Consider  $f: S \rightarrow \mathbb{R}$  where  $c$  is a cluster point of  $S \subset \mathbb{R}$ .

Suppose  $\exists L \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists \delta > 0$  such that whenever  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , we have  $|f(x) - L| < \epsilon$ .

We then say  $f(x)$  *converges* to  $L$  as  $x$  goes to  $c$ .

We say  $L$  is the *limit* of  $f(x)$  as  $x$  goes to  $c$ .

We write  $\lim_{x \rightarrow c} f(x) := L$ , or  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

If no such  $L$  exists, we say the limit does not exist or that  $f$  *diverges* at  $c$ .

**Cheating again:** The notation assumes the limit is unique, we'll prove that momentarily.

**Remark:** It is irrelevant whether  $f(c)$  is defined or not. The limit may not equal  $f(c)$ , even if  $f(c)$  is defined.

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ .*

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.  
Let  $\epsilon > 0$  be given.

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .



## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

Suppose  $x \in S$ ,  $|x - c| < \delta$ , and  $x \neq c$ .

(such an  $x$  exists as  $c$  is a cluster point).

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

Suppose  $x \in S$ ,  $|x - c| < \delta$ , and  $x \neq c$ .

(such an  $x$  exists as  $c$  is a cluster point).

$$|L_1 - L_2|$$

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

Suppose  $x \in S$ ,  $|x - c| < \delta$ , and  $x \neq c$ .

(such an  $x$  exists as  $c$  is a cluster point).

$$|L_1 - L_2| = |L_1 - f(x) + f(x) - L_2|$$

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

Suppose  $x \in S$ ,  $|x - c| < \delta$ , and  $x \neq c$ .

(such an  $x$  exists as  $c$  is a cluster point).

$$|L_1 - L_2| = |L_1 - f(x) + f(x) - L_2| \leq |L_1 - f(x)| + |f(x) - L_2|$$

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

Suppose  $x \in S$ ,  $|x - c| < \delta$ , and  $x \neq c$ .

(such an  $x$  exists as  $c$  is a cluster point).

$$|L_1 - L_2| = |L_1 - f(x) + f(x) - L_2| \leq |L_1 - f(x)| + |f(x) - L_2| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

## Proposition

*Let  $c$  be a cluster point of  $S \subset \mathbb{R}$  and let  $f: S \rightarrow \mathbb{R}$  be such that  $f(x)$  converges as  $x$  goes to  $c$ . Then the limit of  $f(x)$  as  $x$  goes to  $c$  is unique.*

**Proof:** Let both  $L_1$  and  $L_2$  satisfy the definition.

Let  $\epsilon > 0$  be given.

Find  $\delta_1 > 0$  such that  $|f(x) - L_1| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_1$ .

Find  $\delta_2 > 0$  such that  $|f(x) - L_2| < \epsilon/2$  for all  $x \in S \setminus \{c\}$  with  $|x - c| < \delta_2$ .

Put  $\delta := \min\{\delta_1, \delta_2\}$ .

Suppose  $x \in S$ ,  $|x - c| < \delta$ , and  $x \neq c$ .

(such an  $x$  exists as  $c$  is a cluster point).

$$|L_1 - L_2| = |L_1 - f(x) + f(x) - L_2| \leq |L_1 - f(x)| + |f(x) - L_2| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

$$\Rightarrow L_1 = L_2.$$



**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$



**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

Let  $\delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}$ .

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

Let  $\delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}$ .

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$|f(x) - c^2|$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$|f(x) - c^2| = |x^2 - c^2|$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$|f(x) - c^2| = |x^2 - c^2| = |(x + c)(x - c)|$$



**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$|f(x) - c^2| = |x^2 - c^2| = |(x + c)(x - c)| = |x + c| |x - c|$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$\begin{aligned} |f(x) - c^2| &= |x^2 - c^2| = |(x + c)(x - c)| = |x + c| |x - c| \\ &\leq (|x| + |c|) |x - c| \end{aligned}$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$\begin{aligned} |f(x) - c^2| &= |x^2 - c^2| = |(x + c)(x - c)| = |x + c| |x - c| \\ &\leq (|x| + |c|) |x - c| < (2|c| + 1) |x - c| \end{aligned}$$

**Example:** Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) := x^2$ . Then for any  $c \in \mathbb{R}$ ,

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} x^2 = c^2.$$

**Proof:** Fix  $c \in \mathbb{R}$ , and let  $\epsilon > 0$  is given.

$$\text{Let } \delta := \min \left\{ 1, \frac{\epsilon}{2|c| + 1} \right\}.$$

Take  $x \neq c$  such that  $|x - c| < \delta$  (in particular,  $|x - c| < 1$ ).

By reverse triangle inequality,  $|x| - |c| \leq |x - c| < 1$ .

$$\Rightarrow |x| + |c| < 2|c| + 1.$$

Then

$$\begin{aligned} |f(x) - c^2| &= |x^2 - c^2| = |(x + c)(x - c)| = |x + c| |x - c| \\ &\leq (|x| + |c|) |x - c| < (2|c| + 1) |x - c| < (2|c| + 1) \frac{\epsilon}{2|c| + 1} = \epsilon. \end{aligned}$$

□


**Example:** Define  $f: [0, 1) \rightarrow \mathbb{R}$  by  $f(x) := \begin{cases} x & \text{if } x > 0, \\ 1 & \text{if } x = 0. \end{cases}$

**Example:** Define  $f: [0, 1) \rightarrow \mathbb{R}$  by  $f(x) := \begin{cases} x & \text{if } x > 0, \\ 1 & \text{if } x = 0. \end{cases}$

Then  $\lim_{x \rightarrow 0} f(x) = 0$ , even though  $f(0) = 1$ .


**Example:** Define  $f: [0, 1) \rightarrow \mathbb{R}$  by  $f(x) := \begin{cases} x & \text{if } x > 0, \\ 1 & \text{if } x = 0. \end{cases}$

Then  $\lim_{x \rightarrow 0} f(x) = 0$ , even though  $f(0) = 1$ .



**Example:** Define  $f: [0, 1) \rightarrow \mathbb{R}$  by  $f(x) := \begin{cases} x & \text{if } x > 0, \\ 1 & \text{if } x = 0. \end{cases}$

Then  $\lim_{x \rightarrow 0} f(x) = 0$ , even though  $f(0) = 1$ .




**Proof:** Let  $\epsilon > 0$  be given.



**Example:** Define  $f: [0, 1) \rightarrow \mathbb{R}$  by  $f(x) := \begin{cases} x & \text{if } x > 0, \\ 1 & \text{if } x = 0. \end{cases}$


Then  $\lim_{x \rightarrow 0} f(x) = 0$ , even though  $f(0) = 1$ .



**Proof:** Let  $\epsilon > 0$  be given.  
Let  $\delta := \epsilon$ .

**Example:** Define  $f: [0, 1) \rightarrow \mathbb{R}$  by  $f(x) := \begin{cases} x & \text{if } x > 0, \\ 1 & \text{if } x = 0. \end{cases}$

Then  $\lim_{x \rightarrow 0} f(x) = 0$ , even though  $f(0) = 1$ .



**Proof:** Let  $\epsilon > 0$  be given.

Let  $\delta := \epsilon$ .

For  $x \in [0, 1)$ ,  $x \neq 0$ , and  $|x - 0| < \delta$ , we get  $|f(x) - 0| = |x| < \delta = \epsilon$ .



## Lemma

*Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .*

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .



## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon$

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

Now suppose it is not true that  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

Now suppose it is not true that  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

(i.e.,  $\exists \epsilon > 0$  s.t.  $\forall \delta > 0 \exists x \in S \setminus \{c\}$  where  $|x - c| < \delta$  and  $|f(x) - L| \geq \epsilon$ .)

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .

Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

Now suppose it is not true that  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

(i.e.,  $\exists \epsilon > 0$  s.t.  $\forall \delta > 0 \exists x \in S \setminus \{c\}$  where  $|x - c| < \delta$  and  $|f(x) - L| \geq \epsilon$ .)

$\Rightarrow \exists \epsilon > 0$  s.t.  $\forall n \in \mathbb{N}, \exists x_n \in S \setminus \{c\}$ , where  $|x_n - c| < 1/n$  and  $|f(x_n) - L| \geq \epsilon$ .

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .  
Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

Now suppose it is not true that  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

(i.e.,  $\exists \epsilon > 0$  s.t.  $\forall \delta > 0 \exists x \in S \setminus \{c\}$  where  $|x - c| < \delta$  and  $|f(x) - L| \geq \epsilon$ .)

$\Rightarrow \exists \epsilon > 0$  s.t.  $\forall n \in \mathbb{N}, \exists x_n \in S \setminus \{c\}$ , where  $|x_n - c| < 1/n$  and  $|f(x_n) - L| \geq \epsilon$ .

$\{x_n\}_{n=1}^{\infty}$  converges to  $c$ , but  $\{f(x_n)\}_{n=1}^{\infty}$  does not converge to  $L$ . □

## Lemma

Let  $S \subset \mathbb{R}$ , let  $c$  be a cluster point of  $S$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .  
Then  $f(x) \rightarrow L$  as  $x \rightarrow c$  if and only if for every sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S \setminus \{c\}$  such that  $\lim_{n \rightarrow \infty} x_n = c$ , we have that the sequence  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

**Proof:** Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ , and  $\{x_n\}_{n=1}^{\infty}$  is a seq. in  $S \setminus \{c\}$  and  $\lim_{n \rightarrow \infty} x_n = c$ .

Let  $\epsilon > 0$  be given.

Find  $\delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Find an  $M$  such that for  $n \geq M$ , we have  $|x_n - c| < \delta$ .

$\Rightarrow$  for  $n \geq M$ ,  $|f(x_n) - L| < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$  converges to  $L$ .

Now suppose it is not true that  $f(x) \rightarrow L$  as  $x \rightarrow c$ .

(i.e.,  $\exists \epsilon > 0$  s.t.  $\forall \delta > 0 \exists x \in S \setminus \{c\}$  where  $|x - c| < \delta$  and  $|f(x) - L| \geq \epsilon$ .)

$\Rightarrow \exists \epsilon > 0$  s.t.  $\forall n \in \mathbb{N}, \exists x_n \in S \setminus \{c\}$ , where  $|x_n - c| < 1/n$  and  $|f(x_n) - L| \geq \epsilon$ .

$\{x_n\}_{n=1}^{\infty}$  converges to  $c$ , but  $\{f(x_n)\}_{n=1}^{\infty}$  does not converge to  $L$ . □


**Exercise:** It is possible to strengthen the  $\Leftarrow$ : It is enough to suppose that  $\{f(x_n)\}_{n=1}^{\infty}$  converges for every  $\{x_n\}_{n=1}^{\infty}$  without requiring a specific limit.

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .




**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .


Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .


Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n)$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .


Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2)$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .


Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .


Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .


Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :




**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .



**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :




**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R}$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :




**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$$|\sin(t)| \leq 1 \text{ for all } t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)|$$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :




**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.


Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

$x_n \rightarrow 0$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.


Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

$x_n \rightarrow 0 \Rightarrow |x_n| \rightarrow 0$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.


Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

$x_n \rightarrow 0 \Rightarrow |x_n| \rightarrow 0 \Rightarrow x_n \sin(1/x_n) \rightarrow 0$

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .


$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

$x_n \rightarrow 0 \Rightarrow |x_n| \rightarrow 0 \Rightarrow x_n \sin(1/x_n) \rightarrow 0 \Rightarrow \lim_{x \rightarrow 0} x \sin(1/x) = 0.$

□

**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

$x_n \rightarrow 0 \Rightarrow |x_n| \rightarrow 0 \Rightarrow x_n \sin(1/x_n) \rightarrow 0 \Rightarrow \lim_{x \rightarrow 0} x \sin(1/x) = 0.$


□

**Remark:** Keep in mind the “for every sequence”:



**Example:**  $\lim_{x \rightarrow 0} \sin(1/x)$  does not exist, but  $\lim_{x \rightarrow 0} x \sin(1/x) = 0$ .

Graphs of  $\sin(1/x)$  and  $x \sin(1/x)$ :



**Proof:** Define  $x_n := \frac{1}{\pi n + \pi/2}$ , so that  $x_n \rightarrow 0 \Rightarrow \sin(1/x_n) = \sin(\pi n + \pi/2) = (-1)^n$   
 $\Rightarrow \{\sin(1/x_n)\}_{n=1}^{\infty}$  does not converge  $\Rightarrow \lim_{x \rightarrow 0} \sin(1/x)$  does not exist.

Let  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $\mathbb{R} \setminus \{0\}$  such that  $\lim_{n \rightarrow \infty} x_n = 0$ .

$|\sin(t)| \leq 1$  for all  $t \in \mathbb{R} \Rightarrow |x_n \sin(1/x_n) - 0| = |x_n| |\sin(1/x_n)| \leq |x_n|$ .

$x_n \rightarrow 0 \Rightarrow |x_n| \rightarrow 0 \Rightarrow x_n \sin(1/x_n) \rightarrow 0 \Rightarrow \lim_{x \rightarrow 0} x \sin(1/x) = 0$ . □

**Remark:** Keep in mind the “for every sequence”:

If  $x_n := 1/\pi n$ , then  $\{\sin(1/x_n)\}_{n=1}^{\infty} = \{0\}_{n=1}^{\infty}$ , but  $\lim_{x \rightarrow 0} \sin(1/x)$  DNE.

## Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and*

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

Then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

Then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$

**Proof:** Take  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $S \setminus \{c\}$  converging to  $c$ .

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

Then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$

**Proof:** Take  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $S \setminus \{c\}$  converging to  $c$ .

Let  $L_1 := \lim_{x \rightarrow c} f(x)$  and  $L_2 := \lim_{x \rightarrow c} g(x)$ .

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

Then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$

**Proof:** Take  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $S \setminus \{c\}$  converging to  $c$ .

Let  $L_1 := \lim_{x \rightarrow c} f(x)$  and  $L_2 := \lim_{x \rightarrow c} g(x)$ .

By the lemma above,  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L_1$  and  $\{g(x_n)\}_{n=1}^{\infty}$  converges to  $L_2$ .

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

Then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$

**Proof:** Take  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $S \setminus \{c\}$  converging to  $c$ .

Let  $L_1 := \lim_{x \rightarrow c} f(x)$  and  $L_2 := \lim_{x \rightarrow c} g(x)$ .

By the lemma above,  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L_1$  and  $\{g(x_n)\}_{n=1}^{\infty}$  converges to  $L_2$ .

We also have  $f(x_n) \leq g(x_n)$  for all  $n \in \mathbb{N}$ .

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are functions such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist, and

$$f(x) \leq g(x) \quad \text{for all } x \in S \setminus \{c\}.$$

Then

$$\lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} g(x).$$

**Proof:** Take  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $S \setminus \{c\}$  converging to  $c$ .

Let  $L_1 := \lim_{x \rightarrow c} f(x)$  and  $L_2 := \lim_{x \rightarrow c} g(x)$ .

By the lemma above,  $\{f(x_n)\}_{n=1}^{\infty}$  converges to  $L_1$  and  $\{g(x_n)\}_{n=1}^{\infty}$  converges to  $L_2$ .

We also have  $f(x_n) \leq g(x_n)$  for all  $n \in \mathbb{N}$ .

$\Rightarrow L_1 \leq L_2$  (by lemma about sequence limits)





Proofs of the next corollaries are exercises.

Proofs of the next corollaries are exercises.

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that the limit of  $f(x)$  as  $x \rightarrow c$  exists. Suppose  $\exists a, b \in \mathbb{R}$  such that  $a \leq f(x) \leq b$  for all  $x \in S \setminus \{c\}$ .*

Proofs of the next corollaries are exercises.

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that the limit of  $f(x)$  as  $x \rightarrow c$  exists. Suppose  $\exists a, b \in \mathbb{R}$  such that  $a \leq f(x) \leq b$  for all  $x \in S \setminus \{c\}$ .*

*Then  $a \leq \lim_{x \rightarrow c} f(x) \leq b$ .*

Proofs of the next corollaries are exercises.

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that the limit of  $f(x)$  as  $x \rightarrow c$  exists. Suppose  $\exists a, b \in \mathbb{R}$  such that  $a \leq f(x) \leq b$  for all  $x \in S \setminus \{c\}$ .*

*Then  $a \leq \lim_{x \rightarrow c} f(x) \leq b$ .*

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$ ,  $g: S \rightarrow \mathbb{R}$ , and  $h: S \rightarrow \mathbb{R}$  are such that  $f(x) \leq g(x) \leq h(x)$  for all  $x \in S \setminus \{c\}$ .*

Proofs of the next corollaries are exercises.

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that the limit of  $f(x)$  as  $x \rightarrow c$  exists. Suppose  $\exists a, b \in \mathbb{R}$  such that  $a \leq f(x) \leq b$  for all  $x \in S \setminus \{c\}$ .*

*Then  $a \leq \lim_{x \rightarrow c} f(x) \leq b$ .*

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$ ,  $g: S \rightarrow \mathbb{R}$ , and  $h: S \rightarrow \mathbb{R}$  are such that  $f(x) \leq g(x) \leq h(x)$  for all  $x \in S \setminus \{c\}$ .*

*Suppose  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x)$  (in particular, the limits exist).*

Proofs of the next corollaries are exercises.

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that the limit of  $f(x)$  as  $x \rightarrow c$  exists. Suppose  $\exists a, b \in \mathbb{R}$  such that  $a \leq f(x) \leq b$  for all  $x \in S \setminus \{c\}$ .*

*Then  $a \leq \lim_{x \rightarrow c} f(x) \leq b$ .*

### Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$ ,  $g: S \rightarrow \mathbb{R}$ , and  $h: S \rightarrow \mathbb{R}$  are such that  $f(x) \leq g(x) \leq h(x)$  for all  $x \in S \setminus \{c\}$ .*

*Suppose  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x)$  (in particular, the limits exist).*

*Then the limit of  $g(x)$  as  $x$  goes to  $c$  exists and  $\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x)$ .*

## Corollary

*Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist.*

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist. Then

(i)  $\lim_{x \rightarrow c} (f(x) + g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) + \left( \lim_{x \rightarrow c} g(x) \right).$



## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist. Then

(i)  $\lim_{x \rightarrow c} (f(x) + g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) + \left( \lim_{x \rightarrow c} g(x) \right).$

(ii)  $\lim_{x \rightarrow c} (f(x) - g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) - \left( \lim_{x \rightarrow c} g(x) \right).$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist. Then

(i)  $\lim_{x \rightarrow c} (f(x) + g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) + \left( \lim_{x \rightarrow c} g(x) \right).$

(ii)  $\lim_{x \rightarrow c} (f(x) - g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) - \left( \lim_{x \rightarrow c} g(x) \right).$

(iii)  $\lim_{x \rightarrow c} (f(x)g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) \left( \lim_{x \rightarrow c} g(x) \right).$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist. Then

$$(i) \lim_{x \rightarrow c} (f(x) + g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) + \left( \lim_{x \rightarrow c} g(x) \right).$$

$$(ii) \lim_{x \rightarrow c} (f(x) - g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) - \left( \lim_{x \rightarrow c} g(x) \right).$$

$$(iii) \lim_{x \rightarrow c} (f(x)g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) \left( \lim_{x \rightarrow c} g(x) \right).$$

$$(iv) \text{ If } \lim_{x \rightarrow c} g(x) \neq 0, \text{ and } g(x) \neq 0 \text{ for all } x \in S \setminus \{c\}, \text{ then } \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)}.$$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist. Then

$$(i) \lim_{x \rightarrow c} (f(x) + g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) + \left( \lim_{x \rightarrow c} g(x) \right).$$

$$(ii) \lim_{x \rightarrow c} (f(x) - g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) - \left( \lim_{x \rightarrow c} g(x) \right).$$

$$(iii) \lim_{x \rightarrow c} (f(x)g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) \left( \lim_{x \rightarrow c} g(x) \right).$$

$$(iv) \text{ If } \lim_{x \rightarrow c} g(x) \neq 0, \text{ and } g(x) \neq 0 \text{ for all } x \in S \setminus \{c\}, \text{ then } \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)}.$$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that  $\lim_{x \rightarrow c} f(x)$  exists.

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  and  $g: S \rightarrow \mathbb{R}$  are such that the limits of  $f(x)$  and  $g(x)$  as  $x$  goes to  $c$  both exist. Then

- (i)  $\lim_{x \rightarrow c} (f(x) + g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) + \left( \lim_{x \rightarrow c} g(x) \right).$
- (ii)  $\lim_{x \rightarrow c} (f(x) - g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) - \left( \lim_{x \rightarrow c} g(x) \right).$
- (iii)  $\lim_{x \rightarrow c} (f(x)g(x)) = \left( \lim_{x \rightarrow c} f(x) \right) \left( \lim_{x \rightarrow c} g(x) \right).$
- (iv) If  $\lim_{x \rightarrow c} g(x) \neq 0$ , and  $g(x) \neq 0$  for all  $x \in S \setminus \{c\}$ , then  $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)}.$

## Corollary

Let  $S \subset \mathbb{R}$  and let  $c$  be a cluster point of  $S$ . Suppose  $f: S \rightarrow \mathbb{R}$  is such that  $\lim_{x \rightarrow c} f(x)$  exists.

Then  $\lim_{x \rightarrow c} |f(x)| = \left| \lim_{x \rightarrow c} f(x) \right|.$

## Definition

Let  $f: S \rightarrow \mathbb{R}$  be a function and  $A \subset S$ . Define the function  $f|_A: A \rightarrow \mathbb{R}$  by

$$f|_A(x) := f(x) \quad \text{for } x \in A.$$

## Definition

Let  $f: S \rightarrow \mathbb{R}$  be a function and  $A \subset S$ . Define the function  $f|_A: A \rightarrow \mathbb{R}$  by

$$f|_A(x) := f(x) \quad \text{for } x \in A.$$

The function  $f|_A$  is called the *restriction* of  $f$  to  $A$ .

## Definition

Let  $f: S \rightarrow \mathbb{R}$  be a function and  $A \subset S$ . Define the function  $f|_A: A \rightarrow \mathbb{R}$  by

$$f|_A(x) := f(x) \quad \text{for } x \in A.$$

The function  $f|_A$  is called the *restriction* of  $f$  to  $A$ .

The function  $f|_A$  is simply the function  $f$  taken on a smaller domain.



## Definition

Let  $f: S \rightarrow \mathbb{R}$  be a function and  $A \subset S$ . Define the function  $f|_A: A \rightarrow \mathbb{R}$  by

$$f|_A(x) := f(x) \quad \text{for } x \in A.$$

The function  $f|_A$  is called the *restriction* of  $f$  to  $A$ .

The function  $f|_A$  is simply the function  $f$  taken on a smaller domain.

Be careful, e.g.,  $f: \mathbb{R} \rightarrow \mathbb{R}$  defined by say  $f(x) := x^2$  and  $f|_{[0,1]}: [0,1] \rightarrow \mathbb{R}$  really are different functions.

## Definition

Let  $f: S \rightarrow \mathbb{R}$  be a function and  $A \subset S$ . Define the function  $f|_A: A \rightarrow \mathbb{R}$  by

$$f|_A(x) := f(x) \quad \text{for } x \in A.$$

The function  $f|_A$  is called the *restriction* of  $f$  to  $A$ .

The function  $f|_A$  is simply the function  $f$  taken on a smaller domain.

Be careful, e.g.,  $f: \mathbb{R} \rightarrow \mathbb{R}$  defined by say  $f(x) := x^2$  and  $f|_{[0,1]}: [0,1] \rightarrow \mathbb{R}$  really are different functions.

E.g., it is **not** true that  $f|_{[0,1]}(2) = 2^2$ .

## Definition

Let  $f: S \rightarrow \mathbb{R}$  be a function and  $A \subset S$ . Define the function  $f|_A: A \rightarrow \mathbb{R}$  by

$$f|_A(x) := f(x) \quad \text{for } x \in A.$$

The function  $f|_A$  is called the *restriction* of  $f$  to  $A$ .

The function  $f|_A$  is simply the function  $f$  taken on a smaller domain.

Be careful, e.g.,  $f: \mathbb{R} \rightarrow \mathbb{R}$  defined by say  $f(x) := x^2$   
and  $f|_{[0,1]}: [0,1] \rightarrow \mathbb{R}$  really are different functions.

E.g., it is **not** true that  ~~$f|_{[0,1]}(2) = 2^2$~~ .  $f|_{[0,1]}(2)$  is **not defined**. But  $f(2) = 2^2$

## Proposition

*Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function.*

## Proposition

*Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .*

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .



## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0$$

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0 \implies (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0.$$

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0 \implies (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0.$   
 $\implies c$  is a cluster point of  $S$ .

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0 \implies (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0.$   
 $\implies c$  is a cluster point of  $S$ .

Second, suppose  $c$  is a cluster point of  $S$ .

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0 \implies (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0.$   
 $\implies c$  is a cluster point of  $S$ .

Second, suppose  $c$  is a cluster point of  $S$ .

For  $\epsilon > 0$  s.t.  $\epsilon < \alpha$ , we get  $(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) = (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset$ .

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0 \implies (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0.$$

$\implies c$  is a cluster point of  $S$ .

Second, suppose  $c$  is a cluster point of  $S$ .

For  $\epsilon > 0$  s.t.  $\epsilon < \alpha$ , we get  $(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) = (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset$ .

As it is true for all  $\epsilon < \alpha$ ,  $(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset$  for all  $\epsilon > 0$ .

## Proposition

Let  $S \subset \mathbb{R}$ ,  $c \in \mathbb{R}$ , and let  $f: S \rightarrow \mathbb{R}$  be a function. Suppose  $A \subset S$  is such that there is some  $\alpha > 0$  such that  $(A \setminus \{c\}) \cap (c - \alpha, c + \alpha) = (S \setminus \{c\}) \cap (c - \alpha, c + \alpha)$ .

- (i) The point  $c$  is a cluster point of  $A$  if and only if  $c$  is a cluster point of  $S$ .
- (ii) Let  $c$  be a cluster point of  $S$ .  $f(x) \rightarrow L$  as  $x \rightarrow c \iff f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Proof:** First, let  $c$  be a cluster point of  $A$ , and note that  $A \subset S$ .

$(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0 \implies (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset \quad \forall \epsilon > 0.$   
 $\implies c$  is a cluster point of  $S$ .

Second, suppose  $c$  is a cluster point of  $S$ .

For  $\epsilon > 0$  s.t.  $\epsilon < \alpha$ , we get  $(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) = (S \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset$ .

As it is true for all  $\epsilon < \alpha$ ,  $(A \setminus \{c\}) \cap (c - \epsilon, c + \epsilon) \neq \emptyset$  for all  $\epsilon > 0$ .

$\implies c$  is a cluster point of  $A$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .



Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .



Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .

$|x - c| < \alpha \quad \Rightarrow \quad x \in A \setminus \{c\},$

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .

$|x - c| < \alpha \quad \Rightarrow \quad x \in A \setminus \{c\},$

$|x - c| < \delta' \quad \Rightarrow \quad |f(x) - L| = |f|_A(x) - L| < \epsilon.$



Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .

$|x - c| < \alpha \quad \Rightarrow \quad x \in A \setminus \{c\},$

$|x - c| < \delta' \quad \Rightarrow \quad |f(x) - L| = |f|_A(x) - L| < \epsilon.$

□

**Remark:** The hypothesis on  $A$  in the proposition is necessary.

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .

$|x - c| < \alpha \quad \Rightarrow \quad x \in A \setminus \{c\},$

$|x - c| < \delta' \quad \Rightarrow \quad |f(x) - L| = |f|_A(x) - L| < \epsilon.$

□

**Remark:** The hypothesis on  $A$  in the proposition is necessary.

Without it, we only get one implication (exercise):

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .

$|x - c| < \alpha \quad \Rightarrow \quad x \in A \setminus \{c\},$

$|x - c| < \delta' \quad \Rightarrow \quad |f(x) - L| = |f|_A(x) - L| < \epsilon.$

□

**Remark:** The hypothesis on  $A$  in the proposition is necessary.

Without it, we only get one implication (exercise):

Assume  $c$  is a cluster point of  $A$ , then

$f(x) \rightarrow L$  as  $x \rightarrow c \quad \Rightarrow \quad f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Suppose  $c$  is a cluster point of  $S$  and therefore also a cluster point of  $A$ .

Suppose  $f(x) \rightarrow L$  as  $x \rightarrow c$ :

$\forall \epsilon > 0 \quad \exists \delta > 0$  such that if  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ , then  $|f(x) - L| < \epsilon$ .

If  $x \in A \setminus \{c\}$ , then  $x \in S \setminus \{c\}$ .

$\Rightarrow f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

Finally, suppose  $f|_A(x) \rightarrow L$  as  $x \rightarrow c$  and let  $\epsilon > 0$  be given.

$\exists \delta' > 0$  such that if  $x \in A \setminus \{c\}$  and  $|x - c| < \delta'$ , then  $|f|_A(x) - L| < \epsilon$ .

Take  $\delta := \min\{\delta', \alpha\}$ .

Suppose  $x \in S \setminus \{c\}$  and  $|x - c| < \delta$ .

$|x - c| < \alpha \quad \Rightarrow \quad x \in A \setminus \{c\},$

$|x - c| < \delta' \quad \Rightarrow \quad |f(x) - L| = |f|_A(x) - L| < \epsilon.$

□

**Remark:** The hypothesis on  $A$  in the proposition is necessary.

Without it, we only get one implication (exercise):

Assume  $c$  is a cluster point of  $A$ , then

$f(x) \rightarrow L$  as  $x \rightarrow c \quad \Rightarrow \quad f|_A(x) \rightarrow L$  as  $x \rightarrow c$ .

**Notation:**  $\lim_{\substack{x \rightarrow c \\ x \in A}} f(x) := \lim_{x \rightarrow c} f|_A(x).$

## Definition (One sided limits)

Let  $f: S \rightarrow \mathbb{R}$  be function and  $c \in \mathbb{R}$ .

## Definition (One sided limits)

Let  $f: S \rightarrow \mathbb{R}$  be function and  $c \in \mathbb{R}$ .

If  $c$  is a cluster point of  $S \cap (c, \infty)$  and the limit of  $f|_{S \cap (c, \infty)}$  as  $x \rightarrow c$  exists, define

$$\lim_{x \rightarrow c^+} f(x) := \lim_{x \rightarrow c} f|_{S \cap (c, \infty)}(x).$$



## Definition (One sided limits)

Let  $f: S \rightarrow \mathbb{R}$  be function and  $c \in \mathbb{R}$ .

If  $c$  is a cluster point of  $S \cap (c, \infty)$  and the limit of  $f|_{S \cap (c, \infty)}$  as  $x \rightarrow c$  exists, define

$$\lim_{x \rightarrow c^+} f(x) := \lim_{x \rightarrow c} f|_{S \cap (c, \infty)}(x).$$

If  $c$  is a cluster point of  $S \cap (-\infty, c)$  and the limit of  $f|_{S \cap (-\infty, c)}$  as  $x \rightarrow c$  exists, define

$$\lim_{x \rightarrow c^-} f(x) := \lim_{x \rightarrow c} f|_{S \cap (-\infty, c)}(x).$$

## Definition (One sided limits)

Let  $f: S \rightarrow \mathbb{R}$  be function and  $c \in \mathbb{R}$ .

If  $c$  is a cluster point of  $S \cap (c, \infty)$  and the limit of  $f|_{S \cap (c, \infty)}$  as  $x \rightarrow c$  exists, define

$$\lim_{x \rightarrow c^+} f(x) := \lim_{x \rightarrow c} f|_{S \cap (c, \infty)}(x).$$

If  $c$  is a cluster point of  $S \cap (-\infty, c)$  and the limit of  $f|_{S \cap (-\infty, c)}$  as  $x \rightarrow c$  exists, define

$$\lim_{x \rightarrow c^-} f(x) := \lim_{x \rightarrow c} f|_{S \cap (-\infty, c)}(x).$$

Many common notations:

For  $\lim_{x \rightarrow c^-} f(x)$  one sees  $\lim_{\substack{x \rightarrow c \\ x < c}} f(x)$ ,  $\lim_{x \uparrow c} f(x)$ , or  $\lim_{x \nearrow c} f(x)$ .

For  $\lim_{x \rightarrow c^+} f(x)$  one sees  $\lim_{\substack{x \rightarrow c \\ x > c}} f(x)$ ,  $\lim_{x \downarrow c} f(x)$ , or  $\lim_{x \searrow c} f(x)$ .

The proposition above does not apply to one-sided limits.

The proposition above does not apply to one-sided limits.

**Example:** Define  $f: \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) := 1$  for  $x < 0$  and  $f(x) := 0$  for  $x \geq 0$ . Then

$$\lim_{x \rightarrow 0^-} f(x) = 1, \quad \lim_{x \rightarrow 0^+} f(x) = 0, \quad \lim_{x \rightarrow 0} f(x) \text{ does not exist.}$$

The proposition above does not apply to one-sided limits.

**Example:** Define  $f: \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) := 1$  for  $x < 0$  and  $f(x) := 0$  for  $x \geq 0$ . Then

$$\lim_{x \rightarrow 0^-} f(x) = 1, \quad \lim_{x \rightarrow 0^+} f(x) = 0, \quad \lim_{x \rightarrow 0} f(x) \text{ does not exist.}$$

But we do have:

### Proposition

*Let  $S \subset \mathbb{R}$  be such that  $c$  is a cluster point of both  $S \cap (-\infty, c)$  and  $S \cap (c, \infty)$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ .*

The proposition above does not apply to one-sided limits.

**Example:** Define  $f: \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) := 1$  for  $x < 0$  and  $f(x) := 0$  for  $x \geq 0$ . Then

$$\lim_{x \rightarrow 0^-} f(x) = 1, \quad \lim_{x \rightarrow 0^+} f(x) = 0, \quad \lim_{x \rightarrow 0} f(x) \text{ does not exist.}$$

But we do have:

### Proposition

*Let  $S \subset \mathbb{R}$  be such that  $c$  is a cluster point of both  $S \cap (-\infty, c)$  and  $S \cap (c, \infty)$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ . Then  $c$  is a cluster point of  $S$  and*

$$\lim_{x \rightarrow c} f(x) = L \quad \text{if and only if} \quad \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = L.$$

The proposition above does not apply to one-sided limits.

**Example:** Define  $f: \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) := 1$  for  $x < 0$  and  $f(x) := 0$  for  $x \geq 0$ . Then

$$\lim_{x \rightarrow 0^-} f(x) = 1, \quad \lim_{x \rightarrow 0^+} f(x) = 0, \quad \lim_{x \rightarrow 0} f(x) \text{ does not exist.}$$

But we do have:

### Proposition

*Let  $S \subset \mathbb{R}$  be such that  $c$  is a cluster point of both  $S \cap (-\infty, c)$  and  $S \cap (c, \infty)$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ . Then  $c$  is a cluster point of  $S$  and*

$$\lim_{x \rightarrow c} f(x) = L \quad \text{if and only if} \quad \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = L.$$

**Proof:** Exercise.

The proposition above does not apply to one-sided limits.

**Example:** Define  $f: \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) := 1$  for  $x < 0$  and  $f(x) := 0$  for  $x \geq 0$ . Then

$$\lim_{x \rightarrow 0^-} f(x) = 1, \quad \lim_{x \rightarrow 0^+} f(x) = 0, \quad \lim_{x \rightarrow 0} f(x) \text{ does not exist.}$$

But we do have:

### Proposition

*Let  $S \subset \mathbb{R}$  be such that  $c$  is a cluster point of both  $S \cap (-\infty, c)$  and  $S \cap (c, \infty)$ , let  $f: S \rightarrow \mathbb{R}$  be a function, and let  $L \in \mathbb{R}$ . Then  $c$  is a cluster point of  $S$  and*

$$\lim_{x \rightarrow c} f(x) = L \quad \text{if and only if} \quad \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = L.$$

**Proof:** Exercise.

Hint:  $(S \cap (-\infty, c)) \cup (S \cap (c, \infty)) = S \setminus \{c\}$ .



Composition also plays nice with limits if one is careful:

Composition also plays nice with limits if one is careful:

**Exercise:** Let  $c_1$  be a cluster point of  $A \subset \mathbb{R}$  and  $c_2$  be a cluster point of  $B \subset \mathbb{R}$ .

Composition also plays nice with limits if one is careful:

**Exercise:** Let  $c_1$  be a cluster point of  $A \subset \mathbb{R}$  and  $c_2$  be a cluster point of  $B \subset \mathbb{R}$ .

Suppose  $f: A \rightarrow B$  and  $g: B \rightarrow \mathbb{R}$  are such that  $f(x) \rightarrow c_2$  as  $x \rightarrow c_1$  and  $g(y) \rightarrow L$  as  $y \rightarrow c_2$ .

Composition also plays nice with limits if one is careful:

**Exercise:** Let  $c_1$  be a cluster point of  $A \subset \mathbb{R}$  and  $c_2$  be a cluster point of  $B \subset \mathbb{R}$ .

Suppose  $f: A \rightarrow B$  and  $g: B \rightarrow \mathbb{R}$  are such that  $f(x) \rightarrow c_2$  as  $x \rightarrow c_1$  and  $g(y) \rightarrow L$  as  $y \rightarrow c_2$ .

If  $c_2 \in B$ , also suppose that  $g(c_2) = L$  (important).

Composition also plays nice with limits if one is careful:

**Exercise:** Let  $c_1$  be a cluster point of  $A \subset \mathbb{R}$  and  $c_2$  be a cluster point of  $B \subset \mathbb{R}$ .

Suppose  $f: A \rightarrow B$  and  $g: B \rightarrow \mathbb{R}$  are such that  $f(x) \rightarrow c_2$  as  $x \rightarrow c_1$  and  $g(y) \rightarrow L$  as  $y \rightarrow c_2$ .

If  $c_2 \in B$ , also suppose that  $g(c_2) = L$  (important).

Let  $h(x) := g(f(x))$ . Show  $h(x) \rightarrow L$  as  $x \rightarrow c_1$ .

Composition also plays nice with limits if one is careful:

**Exercise:** Let  $c_1$  be a cluster point of  $A \subset \mathbb{R}$  and  $c_2$  be a cluster point of  $B \subset \mathbb{R}$ .

Suppose  $f: A \rightarrow B$  and  $g: B \rightarrow \mathbb{R}$  are such that  $f(x) \rightarrow c_2$  as  $x \rightarrow c_1$  and  $g(y) \rightarrow L$  as  $y \rightarrow c_2$ .

If  $c_2 \in B$ , also suppose that  $g(c_2) = L$  (important).

Let  $h(x) := g(f(x))$ . Show  $h(x) \rightarrow L$  as  $x \rightarrow c_1$ .

Hint: Note that  $f(x)$  could equal  $c_2$  for many  $x \in A$ .