

# BA: 2.4

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

It would be nice to check convergence without knowing the limit.

It would be nice to check convergence without knowing the limit.

### Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

It would be nice to check convergence without knowing the limit.

### Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right|$

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right|$



It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$ . □

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

Then

$$\left| (-1)^n - (-1)^k \right|$$

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

Then

$$\left| (-1)^n - (-1)^k \right| = \left| (-1)^n - (-1)^{n+1} \right|$$

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

Then

$$\left| (-1)^n - (-1)^k \right| = \left| (-1)^n - (-1)^{n+1} \right| = |1 - (-1)|$$

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

Then

$$\left| (-1)^n - (-1)^k \right| = \left| (-1)^n - (-1)^{n+1} \right| = |1 - (-1)| = 2.$$

It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

Then

$$\left| (-1)^n - (-1)^k \right| = \left| (-1)^n - (-1)^{n+1} \right| = |1 - (-1)| = 2.$$

$\Rightarrow$  for any  $\epsilon \leq 2$  the definition is not satisfied.



It would be nice to check convergence without knowing the limit.

## Definition

A sequence  $\{x_n\}_{n=1}^{\infty}$  is a *Cauchy sequence* if for every  $\epsilon > 0$  there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$  and all  $k \geq M$ , we have  $|x_n - x_k| < \epsilon$ .

**Example:**  $\{1/n\}_{n=1}^{\infty}$  is a Cauchy sequence.

**Proof:** Given  $\epsilon > 0$ ,  $\exists M$  such that  $M > 2/\epsilon$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $1/n < \epsilon/2$  and  $1/k < \epsilon/2$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $\left| \frac{1}{n} - \frac{1}{k} \right| \leq \left| \frac{1}{n} \right| + \left| \frac{1}{k} \right| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$

□

**Example:**  $\{(-1)^n\}_{n=1}^{\infty}$  is not a Cauchy sequence.

**Proof:** Given any  $M \in \mathbb{N}$ , take  $n \geq M$  to even, and  $k := n + 1$ .

Then

$$\left| (-1)^n - (-1)^k \right| = \left| (-1)^n - (-1)^{n+1} \right| = |1 - (-1)| = 2.$$

$\Rightarrow$  for any  $\epsilon \leq 2$  the definition is not satisfied.  $\Rightarrow$  The sequence is not Cauchy.

□

## Proposition

*A Cauchy sequence is bounded.*

## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Pick an  $M$  such that for all  $n, k \geq M$ , we have  $|x_n - x_k| < 1$ .

## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Pick an  $M$  such that for all  $n, k \geq M$ , we have  $|x_n - x_k| < 1$ .

In particular, for all  $n \geq M$ ,  $|x_n - x_M| < 1$ .

## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Pick an  $M$  such that for all  $n, k \geq M$ , we have  $|x_n - x_k| < 1$ .

In particular, for all  $n \geq M$ ,  $|x_n - x_M| < 1$ .

By the reverse triangle inequality,  $|x_n| - |x_M| \leq |x_n - x_M| < 1$ .

## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Pick an  $M$  such that for all  $n, k \geq M$ , we have  $|x_n - x_k| < 1$ .

In particular, for all  $n \geq M$ ,  $|x_n - x_M| < 1$ .

By the reverse triangle inequality,  $|x_n| - |x_M| \leq |x_n - x_M| < 1$ .

Hence for  $n \geq M$ ,  $|x_n| < 1 + |x_M|$ .

## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Pick an  $M$  such that for all  $n, k \geq M$ , we have  $|x_n - x_k| < 1$ .

In particular, for all  $n \geq M$ ,  $|x_n - x_M| < 1$ .

By the reverse triangle inequality,  $|x_n| - |x_M| \leq |x_n - x_M| < 1$ .

Hence for  $n \geq M$ ,  $|x_n| < 1 + |x_M|$ .

Let  $B := \max\{|x_1|, |x_2|, \dots, |x_{M-1}|, 1 + |x_M|\}$ .



## Proposition

*A Cauchy sequence is bounded.*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Pick an  $M$  such that for all  $n, k \geq M$ , we have  $|x_n - x_k| < 1$ .

In particular, for all  $n \geq M$ ,  $|x_n - x_M| < 1$ .

By the reverse triangle inequality,  $||x_n| - |x_M|| \leq |x_n - x_M| < 1$ .

Hence for  $n \geq M$ ,  $|x_n| < 1 + |x_M|$ .

Let  $B := \max\{|x_1|, |x_2|, \dots, |x_{M-1}|, 1 + |x_M|\}$ .

Then  $|x_n| \leq B$  for all  $n \in \mathbb{N}$ .



## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

**Proof:** Let  $\epsilon > 0$  be given and suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

**Proof:** Let  $\epsilon > 0$  be given and suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .  
 $\exists M$  such that for  $n \geq M$ ,  $|x_n - x| < \frac{\epsilon}{2}$ .

## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

**Proof:** Let  $\epsilon > 0$  be given and suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .  
 $\exists M$  such that for  $n \geq M$ ,  $|x_n - x| < \frac{\epsilon}{2}$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $|x_n - x_k|$

## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

**Proof:** Let  $\epsilon > 0$  be given and suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .  
 $\exists M$  such that for  $n \geq M$ ,  $|x_n - x| < \frac{\epsilon}{2}$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $|x_n - x_k| = |x_n - x + x - x_k|$

## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

**Proof:** Let  $\epsilon > 0$  be given and suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

$\exists M$  such that for  $n \geq M$ ,  $|x_n - x| < \frac{\epsilon}{2}$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $|x_n - x_k| = |x_n - x + x - x_k| \leq |x_n - x| + |x - x_k|$

## Theorem

*A sequence of real numbers is Cauchy if and only if it converges.*

**Proof:** Let  $\epsilon > 0$  be given and suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $x$ .

$\exists M$  such that for  $n \geq M$ ,  $|x_n - x| < \frac{\epsilon}{2}$ .

$\Rightarrow$  for  $n, k \geq M$ ,  $|x_n - x_k| = |x_n - x + x - x_k| \leq |x_n - x| + |x - x_k| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$ .



Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy.

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .



Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ .

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

$$|a - b|$$

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

$$|a - b| = |a - x_{n_i} + x_{n_i} - x_{m_i} + x_{m_i} - b|$$

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

$$|a - b| = |a - x_{n_i} + x_{n_i} - x_{m_i} + x_{m_i} - b| \leq |a - x_{n_i}| + |x_{n_i} - x_{m_i}| + |x_{m_i} - b|$$

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

$$|a - b| = |a - x_{n_i} + x_{n_i} - x_{m_i} + x_{m_i} - b| \leq |a - x_{n_i}| + |x_{n_i} - x_{m_i}| + |x_{m_i} - b| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

$$|a - b| = |a - x_{n_i} + x_{n_i} - x_{m_i} + x_{m_i} - b| \leq |a - x_{n_i}| + |x_{n_i} - x_{m_i}| + |x_{m_i} - b| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

$$|a - b| < \epsilon \text{ for all } \epsilon > 0 \quad \Rightarrow \quad a = b.$$



Now suppose  $\{x_n\}_{n=1}^{\infty}$  is Cauchy. Thus  $\{x_n\}_{n=1}^{\infty}$  is bounded.

Define  $a := \limsup_{n \rightarrow \infty} x_n$  and  $b := \liminf_{n \rightarrow \infty} x_n$ .

By a theorem from 2.3,  $\exists$  subsequences  $\{x_{n_i}\}_{i=1}^{\infty}$  and  $\{x_{m_i}\}_{i=1}^{\infty}$ , such that  
 $\lim_{i \rightarrow \infty} x_{n_i} = a$  and  $\lim_{i \rightarrow \infty} x_{m_i} = b$ .

Given an  $\epsilon > 0$ ,

$\exists M_1$  such that  $|x_{n_i} - a| < \epsilon/3$  for all  $i \geq M_1$ ,

$\exists M_2$  such that  $|x_{m_i} - b| < \epsilon/3$  for all  $i \geq M_2$ ,

$\exists M_3$  such that  $|x_n - x_k| < \epsilon/3$  for all  $n, k \geq M_3$ .

Let  $M := \max\{M_1, M_2, M_3\}$ .

Consider  $i \geq M$ . Then  $n_i \geq M$  and  $m_i \geq M$ .

$$|a - b| = |a - x_{n_i} + x_{n_i} - x_{m_i} + x_{m_i} - b| \leq |a - x_{n_i}| + |x_{n_i} - x_{m_i}| + |x_{m_i} - b| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

$$|a - b| < \epsilon \text{ for all } \epsilon > 0 \implies a = b.$$

So  $\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n$  and  $\{x_n\}_{n=1}^{\infty}$  converges.



**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.  
The resulting field has the least-upper-bound property.

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.  
The resulting field has the least-upper-bound property.  
This thinking generalizes to more abstract settings.

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.  
The resulting field has the least-upper-bound property.  
This thinking generalizes to more abstract settings.

**Remark:** Cauchy is stronger than  $|x_{n+1} - x_n| \rightarrow 0$  as  $n \rightarrow \infty$



**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .

Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.

We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.

A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.

The resulting field has the least-upper-bound property.

This thinking generalizes to more abstract settings.

**Remark:** Cauchy is stronger than  $|x_{n+1} - x_n| \rightarrow 0$  as  $n \rightarrow \infty$  (or  $|x_{n+j} - x_n| \rightarrow 0$  for a fixed  $j$ ).

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.  
The resulting field has the least-upper-bound property.  
This thinking generalizes to more abstract settings.

**Remark:** Cauchy is stronger than  $|x_{n+1} - x_n| \rightarrow 0$  as  $n \rightarrow \infty$  (or  $|x_{n+j} - x_n| \rightarrow 0$  for a fixed  $j$ ).  
E.g., the partial sums of the harmonic series satisfy  $x_{n+1} - x_n = 1/n$ ,  
but the sequence is not Cauchy.

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.  
The resulting field has the least-upper-bound property.  
This thinking generalizes to more abstract settings.

**Remark:** Cauchy is stronger than  $|x_{n+1} - x_n| \rightarrow 0$  as  $n \rightarrow \infty$  (or  $|x_{n+j} - x_n| \rightarrow 0$  for a fixed  $j$ ).  
E.g., the partial sums of the harmonic series satisfy  $x_{n+1} - x_n = 1/n$ ,  
but the sequence is not Cauchy.  
In fact,  $\lim_{n \rightarrow \infty} |x_{n+j} - x_n| = 0$  for every  $j \in \mathbb{N}$ .

**Remark:** A set  $X$  is *Cauchy-complete* if every Cauchy sequence converges in  $X$ .  
Can be used to define completeness of  $\mathbb{R}$  instead of the least-upper-bound property.  
We proved that for  $\mathbb{R}$ , least-upper-bound property implies Cauchy-complete.  
A way to construct  $\mathbb{R}$  is to “complete”  $\mathbb{Q}$  by “throwing in” enough points to make all Cauchy sequences converge.  
The resulting field has the least-upper-bound property.  
This thinking generalizes to more abstract settings.

**Remark:** Cauchy is stronger than  $|x_{n+1} - x_n| \rightarrow 0$  as  $n \rightarrow \infty$  (or  $|x_{n+j} - x_n| \rightarrow 0$  for a fixed  $j$ ).  
E.g., the partial sums of the harmonic series satisfy  $x_{n+1} - x_n = 1/n$ ,  
but the sequence is not Cauchy.

In fact,  $\lim_{n \rightarrow \infty} |x_{n+j} - x_n| = 0$  for every  $j \in \mathbb{N}$ .

The key point in the definition of Cauchy is that  $n$  and  $k$  vary independently and can be arbitrarily far apart.