

# BA: 7.3

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

A *sequence* in a metric space  $(X, d)$  is a function  $x: \mathbb{N} \rightarrow X$ .

## Definition

A *sequence* in a metric space  $(X, d)$  is a function  $x: \mathbb{N} \rightarrow X$ . Write  $x_n$  for the  $n$ th element.

## Definition

A *sequence* in a metric space  $(X, d)$  is a function  $x: \mathbb{N} \rightarrow X$ . Write  $x_n$  for the  $n$ th element. For the whole sequence write

$$\{x_n\}_{n=1}^{\infty}.$$

## Definition

A *sequence* in a metric space  $(X, d)$  is a function  $x: \mathbb{N} \rightarrow X$ . Write  $x_n$  for the  $n$ th element. For the whole sequence write

$$\{x_n\}_{n=1}^{\infty}.$$

$\{x_n\}_{n=1}^{\infty}$  is *bounded* if  $\exists p \in X$  and  $B \in \mathbb{R}$  such that

$$d(p, x_n) \leq B \quad \text{for all } n \in \mathbb{N}.$$

## Definition

A *sequence* in a metric space  $(X, d)$  is a function  $x: \mathbb{N} \rightarrow X$ . Write  $x_n$  for the  $n$ th element. For the whole sequence write

$$\{x_n\}_{n=1}^{\infty}.$$

$\{x_n\}_{n=1}^{\infty}$  is *bounded* if  $\exists p \in X$  and  $B \in \mathbb{R}$  such that

$$d(p, x_n) \leq B \quad \text{for all } n \in \mathbb{N}.$$

That is,  $\{x_n\}_{n=1}^{\infty}$  is bounded  $\Leftrightarrow \{x_n : n \in \mathbb{N}\}$  is bounded.

## Definition

A *sequence* in a metric space  $(X, d)$  is a function  $x: \mathbb{N} \rightarrow X$ . Write  $x_n$  for the  $n$ th element. For the whole sequence write

$$\{x_n\}_{n=1}^{\infty}.$$

$\{x_n\}_{n=1}^{\infty}$  is *bounded* if  $\exists p \in X$  and  $B \in \mathbb{R}$  such that

$$d(p, x_n) \leq B \quad \text{for all } n \in \mathbb{N}.$$

That is,  $\{x_n\}_{n=1}^{\infty}$  is bounded  $\Leftrightarrow \{x_n : n \in \mathbb{N}\}$  is bounded.

If  $\{n_k\}_{k=1}^{\infty}$  is a sequence of natural numbers such that  $n_{k+1} > n_k$  for all  $k$ , then  $\{x_{n_k}\}_{k=1}^{\infty}$  is a *subsequence* of  $\{x_n\}_{n=1}^{\infty}$ .

## Definition

$\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  is said to *converge* to  $p \in X$  if for every  $\epsilon > 0$ , there exists an  $M \in \mathbb{N}$  such that  $d(x_n, p) < \epsilon$  for all  $n \geq M$ .



## Definition

$\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  is said to *converge* to  $p \in X$  if for every  $\epsilon > 0$ , there exists an  $M \in \mathbb{N}$  such that  $d(x_n, p) < \epsilon$  for all  $n \geq M$ .

$p$  is said to be a *limit* of  $\{x_n\}_{n=1}^{\infty}$ .

## Definition

$\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  is said to *converge* to  $p \in X$  if for every  $\epsilon > 0$ , there exists an  $M \in \mathbb{N}$  such that  $d(x_n, p) < \epsilon$  for all  $n \geq M$ .

$p$  is said to be a *limit* of  $\{x_n\}_{n=1}^{\infty}$ .

If the limit is unique, write  $\lim_{n \rightarrow \infty} x_n := p$ .

## Definition

$\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  is said to *converge* to  $p \in X$  if for every  $\epsilon > 0$ , there exists an  $M \in \mathbb{N}$  such that  $d(x_n, p) < \epsilon$  for all  $n \geq M$ .

$p$  is said to be a *limit* of  $\{x_n\}_{n=1}^{\infty}$ .

If the limit is unique, write  $\lim_{n \rightarrow \infty} x_n := p$ .

A sequence that converges is *convergent*.

## Definition

$\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  is said to *converge* to  $p \in X$  if for every  $\epsilon > 0$ , there exists an  $M \in \mathbb{N}$  such that  $d(x_n, p) < \epsilon$  for all  $n \geq M$ .

$p$  is said to be a *limit* of  $\{x_n\}_{n=1}^{\infty}$ .

If the limit is unique, write  $\lim_{n \rightarrow \infty} x_n := p$ .

A sequence that converges is *convergent*.

Otherwise, the sequence is *divergent*.

## Definition


$\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  is said to *converge* to  $p \in X$  if for every  $\epsilon > 0$ , there exists an  $M \in \mathbb{N}$  such that  $d(x_n, p) < \epsilon$  for all  $n \geq M$ .

$p$  is said to be a *limit* of  $\{x_n\}_{n=1}^{\infty}$ .

If the limit is unique, write  $\lim_{n \rightarrow \infty} x_n := p$ .

A sequence that converges is *convergent*.

Otherwise, the sequence is *divergent*.



Sequence converging to  $p$ . The first 10 points are shown and  $M = 7$  for this  $\epsilon$ .

## Proposition

*A convergent sequence in a metric space has a unique limit.*

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .



## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .  
Take an arbitrary  $\epsilon > 0$ .

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

$d(y, x)$

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

$$d(y, x) \leq d(y, x_n) + d(x_n, x)$$

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

$$d(y, x) \leq d(y, x_n) + d(x_n, x) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

$$d(y, x) \leq d(y, x_n) + d(x_n, x) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

$$d(y, x) < \epsilon \quad \forall \epsilon > 0 \quad \Rightarrow \quad d(y, x) = 0$$



## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

$$d(y, x) \leq d(y, x_n) + d(x_n, x) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

$$d(y, x) < \epsilon \quad \forall \epsilon > 0 \quad \Rightarrow \quad d(y, x) = 0 \quad \Rightarrow \quad y = x.$$

## Proposition

*A convergent sequence in a metric space has a unique limit.*

**Remark:** Proof is exactly the same as for real-sequences.

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  has limits  $x$  and  $y$ .

Take an arbitrary  $\epsilon > 0$ .

Find an  $M_1$  such that for all  $n \geq M_1$ ,  $d(x_n, x) < \epsilon/2$ .

Find an  $M_2$  such that for all  $n \geq M_2$ ,  $d(x_n, y) < \epsilon/2$ .

Consider  $n$  such that  $n \geq M_1$  **and**  $n \geq M_2$ .

$$d(y, x) \leq d(y, x_n) + d(x_n, x) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

$$d(y, x) < \epsilon \quad \forall \epsilon > 0 \quad \Rightarrow \quad d(y, x) = 0 \quad \Rightarrow \quad y = x.$$

Hence the limit (if it exists) is unique.



## Proposition

*A convergent sequence in a metric space is bounded.*

## Proposition

*A convergent sequence in a metric space is bounded.*

## Proposition

*A sequence  $\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  converges to  $p \in X$  if and only if there exists a sequence  $\{a_n\}_{n=1}^{\infty}$  of real numbers such that*

$$d(x_n, p) \leq a_n \quad \text{for all } n \in \mathbb{N}, \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n = 0.$$

## Proposition

*A convergent sequence in a metric space is bounded.*

## Proposition

*A sequence  $\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  converges to  $p \in X$  if and only if there exists a sequence  $\{a_n\}_{n=1}^{\infty}$  of real numbers such that*

$$d(x_n, p) \leq a_n \quad \text{for all } n \in \mathbb{N}, \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n = 0.$$

## Proposition

*Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in a metric space  $(X, d)$ .*

## Proposition

*A convergent sequence in a metric space is bounded.*

## Proposition

*A sequence  $\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  converges to  $p \in X$  if and only if there exists a sequence  $\{a_n\}_{n=1}^{\infty}$  of real numbers such that*

$$d(x_n, p) \leq a_n \quad \text{for all } n \in \mathbb{N}, \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n = 0.$$

## Proposition

*Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in a metric space  $(X, d)$ .*

- (i) If  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$ , then every subsequence  $\{x_{n_k}\}_{k=1}^{\infty}$  converges to  $p$ .*

## Proposition

*A convergent sequence in a metric space is bounded.*

## Proposition

*A sequence  $\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  converges to  $p \in X$  if and only if there exists a sequence  $\{a_n\}_{n=1}^{\infty}$  of real numbers such that*

$$d(x_n, p) \leq a_n \quad \text{for all } n \in \mathbb{N}, \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n = 0.$$

## Proposition

*Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in a metric space  $(X, d)$ .*

- (i) If  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$ , then every subsequence  $\{x_{n_k}\}_{k=1}^{\infty}$  converges to  $p$ .*
- (ii) If for some  $K \in \mathbb{N}$  the  $K$ -tail  $\{x_n\}_{n=K+1}^{\infty}$  converges to  $p \in X$ , then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .*

## Proposition

*A convergent sequence in a metric space is bounded.*

## Proposition

*A sequence  $\{x_n\}_{n=1}^{\infty}$  in a metric space  $(X, d)$  converges to  $p \in X$  if and only if there exists a sequence  $\{a_n\}_{n=1}^{\infty}$  of real numbers such that*

$$d(x_n, p) \leq a_n \quad \text{for all } n \in \mathbb{N}, \quad \text{and} \quad \lim_{n \rightarrow \infty} a_n = 0.$$

## Proposition

*Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in a metric space  $(X, d)$ .*

- (i) If  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$ , then every subsequence  $\{x_{n_k}\}_{k=1}^{\infty}$  converges to  $p$ .*
- (ii) If for some  $K \in \mathbb{N}$  the  $K$ -tail  $\{x_n\}_{n=K+1}^{\infty}$  converges to  $p \in X$ , then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .*

**Proofs:** Exercises.



**Example:** Take  $C([a, b], \mathbb{R})$  be the metric space of continuous functions where  
 $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$  (uniform norm).

**Example:** Take  $C([a, b], \mathbb{R})$  be the metric space of continuous functions where  $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$  (uniform norm).

Convergence is identical to uniform convergence (see chapter 6).

That is,  $\{f_n\}_{n=1}^{\infty}$  converges uniformly if and only if it converges in the metric space sense.

**Example:** Take  $C([a, b], \mathbb{R})$  be the metric space of continuous functions where  $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$  (uniform norm).

Convergence is identical to uniform convergence (see chapter 6).

That is,  $\{f_n\}_{n=1}^{\infty}$  converges uniformly if and only if it converges in the metric space sense.

**Remark:** No metric on the set of functions  $f: [a, b] \rightarrow \mathbb{R}$  gives pointwise convergence.

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ .

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case 
$$\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right).$$

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case 
$$\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right).$$

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case 
$$\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right).$$

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .



## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case 
$$\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right).$$

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .

Fix some  $k = 1, 2, \dots, n$ .

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case  $\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right)$ .

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .

Fix some  $k = 1, 2, \dots, n$ .

For all  $m \geq M$ ,

$$|y_k - x_{m,k}|$$

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case  $\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right)$ .

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .

Fix some  $k = 1, 2, \dots, n$ .

For all  $m \geq M$ ,

$$|y_k - x_{m,k}| = \sqrt{(y_k - x_{m,k})^2}$$

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case  $\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right)$ .

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .

Fix some  $k = 1, 2, \dots, n$ .

For all  $m \geq M$ ,

$$|y_k - x_{m,k}| = \sqrt{(y_k - x_{m,k})^2} \leq \sqrt{\sum_{\ell=1}^n (y_{\ell} - x_{m,\ell})^2}$$

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case  $\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right)$ .

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .

Fix some  $k = 1, 2, \dots, n$ .

For all  $m \geq M$ ,

$$|y_k - x_{m,k}| = \sqrt{(y_k - x_{m,k})^2} \leq \sqrt{\sum_{\ell=1}^n (y_{\ell} - x_{m,\ell})^2} = d(y, x_m) < \epsilon.$$

## Proposition

Consider  $\{x_m\}_{m=1}^{\infty}$  in  $\mathbb{R}^n$ , where  $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n}) \in \mathbb{R}^n$ . Then,  
 $\{x_m\}_{m=1}^{\infty}$  converges  $\Leftrightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges for every  $k = 1, 2, \dots, n$ ,

in which case  $\lim_{m \rightarrow \infty} x_m = \left( \lim_{m \rightarrow \infty} x_{m,1}, \lim_{m \rightarrow \infty} x_{m,2}, \dots, \lim_{m \rightarrow \infty} x_{m,n} \right)$ .

**Proof:** Suppose  $\{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

Given  $\epsilon > 0$ ,  $\exists M$  such that  $\forall m \geq M$ ,  $d(y, x_m) < \epsilon$ .

Fix some  $k = 1, 2, \dots, n$ .

For all  $m \geq M$ ,

$$|y_k - x_{m,k}| = \sqrt{(y_k - x_{m,k})^2} \leq \sqrt{\sum_{\ell=1}^n (y_{\ell} - x_{m,\ell})^2} = d(y, x_m) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$ .

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .



Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m)$$

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2}$$

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2}$$

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \sqrt{\sum_{k=1}^n \frac{\epsilon^2}{n}}$$

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \sqrt{\sum_{k=1}^n \frac{\epsilon^2}{n}} = \epsilon.$$

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \sqrt{\sum_{k=1}^n \frac{\epsilon^2}{n}} = \epsilon.$$

$\Rightarrow \{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

□

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \sqrt{\sum_{k=1}^n \frac{\epsilon^2}{n}} = \epsilon.$$

$\Rightarrow \{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

□

**Example:** In the set of complex numbers  $\mathbb{C}$ ,  $z = x + iy$ ,

Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \sqrt{\sum_{k=1}^n \frac{\epsilon^2}{n}} = \epsilon.$$

$\Rightarrow \{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .

□

**Example:** In the set of complex numbers  $\mathbb{C}$ ,  $z = x + iy$ ,

$\{z_n\}_{n=1}^{\infty} = \{x_n + iy_n\}_{n=1}^{\infty}$  converges to  $z = x + iy$



Now suppose  $\{x_{m,k}\}_{m=1}^{\infty}$  converges to  $y_k$  for every  $k = 1, 2, \dots, n$ .

Given  $\epsilon > 0$ , pick  $M$  such that  $\forall m \geq M$  and  $\forall k = 1, \dots, n$ ,  $|y_k - x_{m,k}| < \epsilon/\sqrt{n}$ .

$$d(y, x_m) = \sqrt{\sum_{k=1}^n (y_k - x_{m,k})^2} < \sqrt{\sum_{k=1}^n \left(\frac{\epsilon}{\sqrt{n}}\right)^2} = \sqrt{\sum_{k=1}^n \frac{\epsilon^2}{n}} = \epsilon.$$

$\Rightarrow \{x_m\}_{m=1}^{\infty}$  converges to  $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$ .



**Example:** In the set of complex numbers  $\mathbb{C}$ ,  $z = x + iy$ ,

$\{z_n\}_{n=1}^{\infty} = \{x_n + iy_n\}_{n=1}^{\infty}$  converges to  $z = x + iy$

$\Leftrightarrow$

$\{x_n\}_{n=1}^{\infty}$  converges to  $x$  and  $\{y_n\}_{n=1}^{\infty}$  converges to  $y$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .  
Let  $U$  be an open neighborhood of  $p$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

Find  $M \in \mathbb{N}$  such that  $\forall n \geq M, \quad d(p, x_n) < \epsilon,$

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

Find  $M \in \mathbb{N}$  such that  $\forall n \geq M, d(p, x_n) < \epsilon$ , that is,  $x_n \in B(p, \epsilon) \subset U$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

Find  $M \in \mathbb{N}$  such that  $\forall n \geq M, d(p, x_n) < \epsilon$ , that is,  $x_n \in B(p, \epsilon) \subset U$ .

The other direction:



## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

Find  $M \in \mathbb{N}$  such that  $\forall n \geq M, d(p, x_n) < \epsilon$ , that is,  $x_n \in B(p, \epsilon) \subset U$ .

The other direction:

Given  $\epsilon > 0$ , let  $U := B(p, \epsilon)$  be the neighborhood of  $p$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

Find  $M \in \mathbb{N}$  such that  $\forall n \geq M, d(p, x_n) < \epsilon$ , that is,  $x_n \in B(p, \epsilon) \subset U$ .

The other direction:

Given  $\epsilon > 0$ , let  $U := B(p, \epsilon)$  be the neighborhood of  $p$ .

$\Rightarrow \exists M \in \mathbb{N}$  such that for  $n \geq M$ ,  $x_n \in U = B(p, \epsilon)$ ,

## Proposition

*Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Then  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if for every open neighborhood  $U$  of  $p$ , there exists an  $M \in \mathbb{N}$  such that for all  $n \geq M$ , we have  $x_n \in U$ .*

**Proof:** Suppose  $\{x_n\}_{n=1}^{\infty}$  converges to  $p$ .

Let  $U$  be an open neighborhood of  $p$ .

$\Rightarrow \exists \epsilon > 0$  such that  $B(p, \epsilon) \subset U$ .

Find  $M \in \mathbb{N}$  such that  $\forall n \geq M, d(p, x_n) < \epsilon$ , that is,  $x_n \in B(p, \epsilon) \subset U$ .

The other direction:

Given  $\epsilon > 0$ , let  $U := B(p, \epsilon)$  be the neighborhood of  $p$ .

$\Rightarrow \exists M \in \mathbb{N}$  such that for  $n \geq M$ ,  $x_n \in U = B(p, \epsilon)$ ,

that is,  $d(p, x_n) < \epsilon$ .



## Proposition

*Let  $(X, d)$  be a metric space,  $E \subset X$  a closed set, and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $E$  that converges to some  $p \in X$ . Then  $p \in E$ .*

## Proposition

*Let  $(X, d)$  be a metric space,  $E \subset X$  a closed set, and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $E$  that converges to some  $p \in X$ . Then  $p \in E$ .*

**Proof:** We prove the contrapositive.

## Proposition

*Let  $(X, d)$  be a metric space,  $E \subset X$  a closed set, and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $E$  that converges to some  $p \in X$ . Then  $p \in E$ .*

**Proof:** We prove the contrapositive.

Suppose  $\{x_n\}_{n=1}^{\infty}$  in  $X$  converges to  $p \in E^c$ .

## Proposition

*Let  $(X, d)$  be a metric space,  $E \subset X$  a closed set, and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $E$  that converges to some  $p \in X$ . Then  $p \in E$ .*

**Proof:** We prove the contrapositive.

Suppose  $\{x_n\}_{n=1}^{\infty}$  in  $X$  converges to  $p \in E^c$ .

$E^c$  is open

## Proposition

*Let  $(X, d)$  be a metric space,  $E \subset X$  a closed set, and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $E$  that converges to some  $p \in X$ . Then  $p \in E$ .*

**Proof:** We prove the contrapositive.

Suppose  $\{x_n\}_{n=1}^{\infty}$  in  $X$  converges to  $p \in E^c$ .

$E^c$  is open  $\Rightarrow \exists M$  such that  $\forall n \geq M, x_n \in E^c$ .



## Proposition

*Let  $(X, d)$  be a metric space,  $E \subset X$  a closed set, and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $E$  that converges to some  $p \in X$ . Then  $p \in E$ .*

**Proof:** We prove the contrapositive.

Suppose  $\{x_n\}_{n=1}^{\infty}$  in  $X$  converges to  $p \in E^c$ .

$E^c$  is open  $\Rightarrow \exists M$  such that  $\forall n \geq M, x_n \in E^c$ .

$\Rightarrow \{x_n\}_{n=1}^{\infty}$  is not a sequence in  $E$ .



## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $p \in \bar{A}$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  of elements in  $A$  such that  $\lim_{n \rightarrow \infty} x_n = p$ .*

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $p \in \bar{A}$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  of elements in  $A$  such that  $\lim_{n \rightarrow \infty} x_n = p$ .*

**Proof:** Let  $p \in \bar{A}$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $p \in \bar{A}$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  of elements in  $A$  such that  $\lim_{n \rightarrow \infty} x_n = p$ .*

**Proof:** Let  $p \in \bar{A}$ .

For every  $n \in \mathbb{N}$ ,  $\exists x_n \in B(p, 1/n) \cap A$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $p \in \bar{A}$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  of elements in  $A$  such that  $\lim_{n \rightarrow \infty} x_n = p$ .*

**Proof:** Let  $p \in \bar{A}$ .

For every  $n \in \mathbb{N}$ ,  $\exists x_n \in B(p, 1/n) \cap A$ .

$$d(p, x_n) < 1/n \quad \forall n$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $p \in \bar{A}$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  of elements in  $A$  such that  $\lim_{n \rightarrow \infty} x_n = p$ .

**Proof:** Let  $p \in \bar{A}$ .

For every  $n \in \mathbb{N}$ ,  $\exists x_n \in B(p, 1/n) \cap A$ .

$$d(p, x_n) < 1/n \quad \forall n \quad \Rightarrow \quad \lim_{n \rightarrow \infty} x_n = p.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $p \in \bar{A}$  if and only if there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  of elements in  $A$  such that  $\lim_{n \rightarrow \infty} x_n = p$ .

**Proof:** Let  $p \in \bar{A}$ .

For every  $n \in \mathbb{N}$ ,  $\exists x_n \in B(p, 1/n) \cap A$ .

$$d(p, x_n) < 1/n \quad \forall n \quad \Rightarrow \quad \lim_{n \rightarrow \infty} x_n = p.$$

The other direction: Exercise.



**Exercise:** Let  $(X, d)$  be a metric space where  $d$  is the discrete metric. Suppose  $\{x_n\}_{n=1}^{\infty}$  is a convergent sequence in  $X$ . Show that there exists a  $K \in \mathbb{N}$  such that for all  $n \geq K$ , we have  $x_n = x_K$ .



**Exercise:** Let  $(X, d)$  be a metric space where  $d$  is the discrete metric. Suppose  $\{x_n\}_{n=1}^{\infty}$  is a convergent sequence in  $X$ . Show that there exists a  $K \in \mathbb{N}$  such that for all  $n \geq K$ , we have  $x_n = x_K$ .

A set  $S \subset X$  is said to be *dense* in  $X$  if  $X \subset \bar{S}$  or in other words if for every  $p \in X$ , there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S$  that converges to  $p$ .

**Exercise:** Let  $(X, d)$  be a metric space where  $d$  is the discrete metric. Suppose  $\{x_n\}_{n=1}^{\infty}$  is a convergent sequence in  $X$ . Show that there exists a  $K \in \mathbb{N}$  such that for all  $n \geq K$ , we have  $x_n = x_K$ .

A set  $S \subset X$  is said to be *dense* in  $X$  if  $X \subset \bar{S}$  or in other words if for every  $p \in X$ , there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S$  that converges to  $p$ .

**Exercise:** Prove that  $\mathbb{R}^n$  contains a countable dense subset.

**Exercise:** Let  $(X, d)$  be a metric space where  $d$  is the discrete metric. Suppose  $\{x_n\}_{n=1}^{\infty}$  is a convergent sequence in  $X$ . Show that there exists a  $K \in \mathbb{N}$  such that for all  $n \geq K$ , we have  $x_n = x_K$ .

A set  $S \subset X$  is said to be *dense* in  $X$  if  $X \subset \bar{S}$  or in other words if for every  $p \in X$ , there exists a sequence  $\{x_n\}_{n=1}^{\infty}$  in  $S$  that converges to  $p$ .

**Exercise:** Prove that  $\mathbb{R}^n$  contains a countable dense subset.

**Exercise:** Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty}$  a sequence in  $X$ . Prove that  $\{x_n\}_{n=1}^{\infty}$  converges to  $p \in X$  if and only if every subsequence of  $\{x_n\}_{n=1}^{\infty}$  has a subsequence that converges to  $p$ .