

# BA: 0.3 part 1

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We start with a set theory refresher.

We start with a set theory refresher.

### Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

By “ $:=$ ” we mean definition of  $S$ .

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

By “ $:=$ ” we mean definition of  $S$ .

“ $1 \in S$ ” means “1 belongs to  $S$ .”

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

By “ $:=$ ” we mean definition of  $S$ .

“ $1 \in S$ ” means “1 belongs to  $S$ .”

“ $1, 2 \in S$ ” is shorthand for “ $1 \in S$  and  $2 \in S$ .”



We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

By “ $:=$ ” we mean definition of  $S$ .

“ $1 \in S$ ” means “1 belongs to  $S$ .”

“ $1, 2 \in S$ ” is shorthand for “ $1 \in S$  and  $2 \in S$ .”

“ $7 \notin S$ ” means “7 does not belong to  $S$ .”

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

By “ $:=$ ” we mean definition of  $S$ .

“ $1 \in S$ ” means “1 belongs to  $S$ .”

“ $1, 2 \in S$ ” is shorthand for “ $1 \in S$  and  $2 \in S$ .”

“ $7 \notin S$ ” means “7 does not belong to  $S$ .”

Elements come from a *universe*.

We start with a set theory refresher.

## Definition

A *set* is a collection of objects called *elements* or *members*. A set with no objects is called the *empty set* and is denoted by  $\emptyset$  (or sometimes by  $\{\}$ ).

Set is a “club.” An element can be a member of several clubs.  
Moreover, sets (clubs) can be members as well.

$S := \{0, 1, 2\}$  is the set containing the three elements 0, 1, and 2.

By “ $:=$ ” we mean definition of  $S$ .

“ $1 \in S$ ” means “1 belongs to  $S$ .”

“ $1, 2 \in S$ ” is shorthand for “ $1 \in S$  and  $2 \in S$ .”

“ $7 \notin S$ ” means “7 does not belong to  $S$ .”


Elements come from a *universe*.

Universe is understood from context. E.g., real numbers.

Every element of  $T := \{0, 2\}$  is also an element of  $S$ . ( $S = \{0, 1, 2\}$ )


Every element of  $T := \{0, 2\}$  is also an element of  $S$ . ( $S = \{0, 1, 2\}$ )

Then write  $T \subset S$ .



Every element of  $T := \{0, 2\}$  is also an element of  $S$ . ( $S = \{0, 1, 2\}$ )

Then write  $T \subset S$ .




## Definition

- (i) A set  $A$  is a *subset* of a set  $B$  if  $x \in A$  implies  $x \in B$ , and we write  $A \subset B$  (or  $B \supset A$ ).

Every element of  $T := \{0, 2\}$  is also an element of  $S$ . ( $S = \{0, 1, 2\}$ )

Then write  $T \subset S$ .




## Definition

- (i) A set  $A$  is a *subset* of a set  $B$  if  $x \in A$  implies  $x \in B$ , and we write  $A \subset B$  (or  $B \supset A$ ).
- (ii) Two sets  $A$  and  $B$  are *equal* if  $A \subset B$  and  $B \subset A$ . We write  $A = B$ .  
If  $A$  and  $B$  are not equal we write  $A \neq B$ .

Every element of  $T := \{0, 2\}$  is also an element of  $S$ . ( $S = \{0, 1, 2\}$ )

Then write  $T \subset S$ .




## Definition

- (i) A set  $A$  is a *subset* of a set  $B$  if  $x \in A$  implies  $x \in B$ , and we write  $A \subset B$  (or  $B \supset A$ ).
- (ii) Two sets  $A$  and  $B$  are *equal* if  $A \subset B$  and  $B \subset A$ . We write  $A = B$ .  
If  $A$  and  $B$  are not equal we write  $A \neq B$ .
- (iii) A set  $A$  is a *proper subset* of  $B$  if  $A \subset B$  and  $A \neq B$ . We write  $A \subsetneq B$ .



Every element of  $T := \{0, 2\}$  is also an element of  $S$ . ( $S = \{0, 1, 2\}$ )

Then write  $T \subset S$ .



## Definition

- (i) A set  $A$  is a *subset* of a set  $B$  if  $x \in A$  implies  $x \in B$ , and we write  $A \subset B$  (or  $B \supset A$ ).
- (ii) Two sets  $A$  and  $B$  are *equal* if  $A \subset B$  and  $B \subset A$ . We write  $A = B$ .  
If  $A$  and  $B$  are not equal we write  $A \neq B$ .
- (iii) A set  $A$  is a *proper subset* of  $B$  if  $A \subset B$  and  $A \neq B$ . We write  $A \subsetneq B$ .

E.g.,  $T \subset S$ , but  $T \neq S$  so  $T \subsetneq S$ .

*Set building notation:*  $\{x \in A : P(x)\}.$

*Set building notation:*  $\{x \in A : P(x)\}.$

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

**Example sets:**

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

### **Example sets:**

*Natural numbers*,  $\mathbb{N} := \{1, 2, 3, \dots\}$ .



*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

### **Example sets:**

*Natural numbers*,  $\mathbb{N} := \{1, 2, 3, \dots\}$ .

*Integers*,  $\mathbb{Z} := \{0, -1, 1, -2, 2, \dots\}$ .

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

### **Example sets:**

*Natural numbers*,  $\mathbb{N} := \{1, 2, 3, \dots\}$ .

*Integers*,  $\mathbb{Z} := \{0, -1, 1, -2, 2, \dots\}$ .

*Rational numbers*,  $\mathbb{Q} := \{\frac{m}{n} : m, n \in \mathbb{Z} \text{ and } n \neq 0\}$ .

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

### **Example sets:**

*Natural numbers*,  $\mathbb{N} := \{1, 2, 3, \dots\}$ .

*Integers*,  $\mathbb{Z} := \{0, -1, 1, -2, 2, \dots\}$ .

*Rational numbers*,  $\mathbb{Q} := \{\frac{m}{n} : m, n \in \mathbb{Z} \text{ and } n \neq 0\}$ .

*Even natural numbers*,  $\{2m : m \in \mathbb{N}\}$ .

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

### **Example sets:**

*Natural numbers*,  $\mathbb{N} := \{1, 2, 3, \dots\}$ .

*Integers*,  $\mathbb{Z} := \{0, -1, 1, -2, 2, \dots\}$ .

*Rational numbers*,  $\mathbb{Q} := \{\frac{m}{n} : m, n \in \mathbb{Z} \text{ and } n \neq 0\}$ .

*Even natural numbers*,  $\{2m : m \in \mathbb{N}\}$ .

The set of all real numbers is denoted by  $\mathbb{R}$ .

*Set building notation:*  $\{x \in A : P(x)\}$ .

Subset of  $A$  containing elements that satisfy property  $P(x)$ .

E.g.,  $\{x \in S : x \neq 2\} = \{0, 1\}$ .

May abbreviate as  $\{x : P(x)\}$  if  $A$  is understood from context.

" $x \in A$ " can also be replaced by a formula.

### **Example sets:**

*Natural numbers*,  $\mathbb{N} := \{1, 2, 3, \dots\}$ .

*Integers*,  $\mathbb{Z} := \{0, -1, 1, -2, 2, \dots\}$ .

*Rational numbers*,  $\mathbb{Q} := \{\frac{m}{n} : m, n \in \mathbb{Z} \text{ and } n \neq 0\}$ .

*Even natural numbers*,  $\{2m : m \in \mathbb{N}\}$ .

The set of all real numbers is denoted by  $\mathbb{R}$ .

Note:  $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$ .

## Definition

- (i) A *union* of two sets  $A$  and  $B$ :  
 $A \cup B := \{x : x \in A \text{ or } x \in B\}.$

## Definition

- (i) A *union* of two sets  $A$  and  $B$ :

$$A \cup B := \{x : x \in A \text{ or } x \in B\}.$$

- (ii) An *intersection* of two sets  $A$  and  $B$ :

$$A \cap B := \{x : x \in A \text{ and } x \in B\}.$$

## Definition

- (i) A *union* of two sets  $A$  and  $B$ :  
$$A \cup B := \{x : x \in A \text{ or } x \in B\}.$$
- (ii) An *intersection* of two sets  $A$  and  $B$ :  
$$A \cap B := \{x : x \in A \text{ and } x \in B\}.$$
- (iii) A *complement of  $B$  relative to  $A$*  (or *set-theoretic difference* of  $A$  and  $B$ ):  
$$A \setminus B := \{x : x \in A \text{ and } x \notin B\}.$$




## Definition


- (i) A *union* of two sets  $A$  and  $B$ :  
$$A \cup B := \{x : x \in A \text{ or } x \in B\}.$$
- (ii) An *intersection* of two sets  $A$  and  $B$ :  
$$A \cap B := \{x : x \in A \text{ and } x \in B\}.$$
- (iii) A *complement of  $B$  relative to  $A$*  (or *set-theoretic difference* of  $A$  and  $B$ ):  
$$A \setminus B := \{x : x \in A \text{ and } x \notin B\}.$$
- (iv) We say *complement* of  $B$  and write  $B^c$  instead of  $A \setminus B$  if  $A$  is either the universe or the obvious set containing  $B$  and understood from context.

## Definition


- (i) A *union* of two sets  $A$  and  $B$ :  
 $A \cup B := \{x : x \in A \text{ or } x \in B\}.$
- (ii) An *intersection* of two sets  $A$  and  $B$ :  
 $A \cap B := \{x : x \in A \text{ and } x \in B\}.$
- (iii) A *complement of  $B$  relative to  $A$*  (or *set-theoretic difference of  $A$  and  $B$* ):  
 $A \setminus B := \{x : x \in A \text{ and } x \notin B\}.$
- (iv) We say *complement of  $B$*  and write  $B^c$  instead of  $A \setminus B$  if  $A$  is either the universe or the obvious set containing  $B$  and understood from context.
- (v)  $A$  and  $B$  are *disjoint* if  $A \cap B = \emptyset$ .



$$A \cup B$$



$$A \cap B$$



$$A \setminus B$$



$$B^c$$

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of =).

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .



## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

Thus  $x \in (A \setminus B) \cap (A \setminus C)$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

Thus  $x \in (A \setminus B) \cap (A \setminus C)$ .

On the other hand, suppose  $x \in (A \setminus B) \cap (A \setminus C)$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

Thus  $x \in (A \setminus B) \cap (A \setminus C)$ .

On the other hand, suppose  $x \in (A \setminus B) \cap (A \setminus C)$ .

In particular,  $x \in (A \setminus B)$ , so  $x \in A$  and  $x \notin B$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

Thus  $x \in (A \setminus B) \cap (A \setminus C)$ .

On the other hand, suppose  $x \in (A \setminus B) \cap (A \setminus C)$ .

In particular,  $x \in (A \setminus B)$ , so  $x \in A$  and  $x \notin B$ .

As also  $x \in (A \setminus C)$ , then  $x \notin C$ .

## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

Thus  $x \in (A \setminus B) \cap (A \setminus C)$ .

On the other hand, suppose  $x \in (A \setminus B) \cap (A \setminus C)$ .

In particular,  $x \in (A \setminus B)$ , so  $x \in A$  and  $x \notin B$ .

As also  $x \in (A \setminus C)$ , then  $x \notin C$ .

Hence  $x \in A \setminus (B \cup C)$ .



## Theorem (DeMorgan)

Let  $A, B, C$  be sets. Then  $(B \cup C)^c = B^c \cap C^c$ ,  $(B \cap C)^c = B^c \cup C^c$ ,

or,  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ ,  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Proof:** The second statement proves the first if  $A$  is the “universe.”

Let us prove  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$  (remember the definition of  $=$ ).

First, suppose  $x \in A \setminus (B \cup C)$ .

Then  $x$  is in  $A$ , but not in  $B$  nor  $C$ .

Hence  $x$  is in  $A$  and not in  $B$ , that is,  $x \in A \setminus B$ .

Similarly  $x \in A \setminus C$ .

Thus  $x \in (A \setminus B) \cap (A \setminus C)$ .

On the other hand, suppose  $x \in (A \setminus B) \cap (A \setminus C)$ .

In particular,  $x \in (A \setminus B)$ , so  $x \in A$  and  $x \notin B$ .

As also  $x \in (A \setminus C)$ , then  $x \notin C$ .

Hence  $x \in A \setminus (B \cup C)$ .

The proof of the other equality is left as an exercise.



Results called *Theorem, Proposition, Lemma, Corollary*. . .

Results called *Theorem, Proposition, Lemma, Corollary*. . .

Often:

Theorem is a bigger result than Proposition,

Results called *Theorem, Proposition, Lemma, Corollary*. . .

Often:

Theorem is a bigger result than Proposition,

Lemma is a tool or an intermediate step,

Results called *Theorem, Proposition, Lemma, Corollary*. . .

Often:

Theorem is a bigger result than Proposition,

Lemma is a tool or an intermediate step,

and Corollary is a quick consequence of something.

Results called *Theorem, Proposition, Lemma, Corollary*. . .

Often:

Theorem is a bigger result than Proposition,

Lemma is a tool or an intermediate step,

and Corollary is a quick consequence of something.

Often it is just traditional or stylistic. Do not read into it *too* much.

Results called *Theorem, Proposition, Lemma, Corollary*. . .

Often:

Theorem is a bigger result than Proposition,  
Lemma is a tool or an intermediate step,  
and Corollary is a quick consequence of something.

Often it is just traditional or stylistic. Do not read into it *too* much.

Important to distinguish between results (as above) and an *Axiom* or a *Postulate*, which are statements assumed to be true without proof.

Results called *Theorem*, *Proposition*, *Lemma*, *Corollary*. . .

Often:

Theorem is a bigger result than Proposition,  
Lemma is a tool or an intermediate step,  
and Corollary is a quick consequence of something.

Often it is just traditional or stylistic. Do not read into it *too* much.

Important to distinguish between results (as above) and an *Axiom* or a *Postulate*, which are statements assumed to be true without proof.

We also encountered *Definitions* which simply give language or notation, but do not actually say anything is true or false.



Results called *Theorem*, *Proposition*, *Lemma*, *Corollary*. . .

Often:

Theorem is a bigger result than Proposition,  
Lemma is a tool or an intermediate step,  
and Corollary is a quick consequence of something.

Often it is just traditional or stylistic. Do not read into it *too* much.

Important to distinguish between results (as above) and an *Axiom* or a *Postulate*, which are statements assumed to be true without proof.

We also encountered *Definitions* which simply give language or notation, but do not actually say anything is true or false. (Confusing sometimes when one says “by definition we have this or that,” but that just means we write out what the definition means using other statements, not that the definition itself implies something.)

Unions/intersections can be taken over more sets.

Unions/intersections can be taken over more sets. If  $\{A_1, A_2, A_3, \dots\}$ ,

Unions/intersections can be taken over more sets. If  $\{A_1, A_2, A_3, \dots\}$ , then

$$\bigcup_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for some } n \in \mathbb{N}\},$$

$$\bigcap_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for all } n \in \mathbb{N}\}.$$

Unions/intersections can be taken over more sets. If  $\{A_1, A_2, A_3, \dots\}$ , then

$$\bigcup_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for some } n \in \mathbb{N}\},$$
$$\bigcap_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for all } n \in \mathbb{N}\}.$$

We could index by other sets such as two natural numbers:

$\{A_{1,1}, A_{1,2}, A_{2,1}, A_{1,3}, A_{2,2}, A_{3,1}, \dots\}.$

Unions/intersections can be taken over more sets. If  $\{A_1, A_2, A_3, \dots\}$ , then

$$\bigcup_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for some } n \in \mathbb{N}\},$$

$$\bigcap_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for all } n \in \mathbb{N}\}.$$

We could index by other sets such as two natural numbers:

$\{A_{1,1}, A_{1,2}, A_{2,1}, A_{1,3}, A_{2,2}, A_{3,1}, \dots\}$ .

$$\bigcup_{n=1}^{\infty} \bigcup_{m=1}^{\infty} A_{n,m} = \bigcup_{n=1}^{\infty} \left( \bigcup_{m=1}^{\infty} A_{n,m} \right).$$

Unions/intersections can be taken over more sets. If  $\{A_1, A_2, A_3, \dots\}$ , then

$$\bigcup_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for some } n \in \mathbb{N}\},$$
$$\bigcap_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for all } n \in \mathbb{N}\}.$$

We could index by other sets such as two natural numbers:  
 $\{A_{1,1}, A_{1,2}, A_{2,1}, A_{1,3}, A_{2,2}, A_{3,1}, \dots\}.$

$$\bigcup_{n=1}^{\infty} \bigcup_{m=1}^{\infty} A_{n,m} = \bigcup_{n=1}^{\infty} \left( \bigcup_{m=1}^{\infty} A_{n,m} \right).$$

Same for intersections.

Unions/intersections can be taken over more sets. If  $\{A_1, A_2, A_3, \dots\}$ , then

$$\bigcup_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for some } n \in \mathbb{N}\},$$
$$\bigcap_{n=1}^{\infty} A_n := \{x : x \in A_n \text{ for all } n \in \mathbb{N}\}.$$

We could index by other sets such as two natural numbers:

$\{A_{1,1}, A_{1,2}, A_{2,1}, A_{1,3}, A_{2,2}, A_{3,1}, \dots\}$ .

$$\bigcup_{n=1}^{\infty} \bigcup_{m=1}^{\infty} A_{n,m} = \bigcup_{n=1}^{\infty} \left( \bigcup_{m=1}^{\infty} A_{n,m} \right).$$

Same for intersections.

It is not hard to see that the order of the unions can be swapped.

Same for intersections.



But switching mixed unions/intersections is not always possible.

But switching mixed unions/intersections is not always possible.

**Example:**

$$\bigcup_{n=1}^{\infty} \bigcap_{m=1}^{\infty} \{k \in \mathbb{N} : mk < n\} = \bigcup_{n=1}^{\infty} \emptyset = \emptyset.$$

But switching mixed unions/intersections is not always possible.

**Example:**

$$\bigcup_{n=1}^{\infty} \bigcap_{m=1}^{\infty} \{k \in \mathbb{N} : mk < n\} = \bigcup_{n=1}^{\infty} \emptyset = \emptyset.$$

However,

$$\bigcap_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{k \in \mathbb{N} : mk < n\} = \bigcap_{m=1}^{\infty} \mathbb{N} = \mathbb{N}.$$

But switching mixed unions/intersections is not always possible.

**Example:**

$$\bigcup_{n=1}^{\infty} \bigcap_{m=1}^{\infty} \{k \in \mathbb{N} : mk < n\} = \bigcup_{n=1}^{\infty} \emptyset = \emptyset.$$

However,

$$\bigcap_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{k \in \mathbb{N} : mk < n\} = \bigcap_{m=1}^{\infty} \mathbb{N} = \mathbb{N}.$$

The index set could be another set than natural numbers in which case we use the following notation:

$$\bigcup_{\lambda \in I} A_{\lambda} := \{x : x \in A_{\lambda} \text{ for some } \lambda \in I\}, \quad \bigcap_{\lambda \in I} A_{\lambda} := \{x : x \in A_{\lambda} \text{ for all } \lambda \in I\}.$$