

# BA: 1.4

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

For  $a < b$ , define the intervals:

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

For  $a < b$ , define the intervals:

$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$  (closed interval)

$(a, b) := \{x \in \mathbb{R} : a < x < b\}$  (open interval)

For  $a < b$ , define the intervals:

$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$  (closed interval)

$(a, b) := \{x \in \mathbb{R} : a < x < b\}$  (open interval)

$(a, b] := \{x \in \mathbb{R} : a < x \leq b\}$  (half-open interval)

For  $a < b$ , define the intervals:

$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$  (closed interval)

$(a, b) := \{x \in \mathbb{R} : a < x < b\}$  (open interval)

$(a, b] := \{x \in \mathbb{R} : a < x \leq b\}$  (half-open interval)

$[a, b) := \{x \in \mathbb{R} : a \leq x < b\}$  (half-open interval)

For  $a < b$ , define the intervals:

$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$  (closed interval)

$(a, b) := \{x \in \mathbb{R} : a < x < b\}$  (open interval)

$(a, b] := \{x \in \mathbb{R} : a < x \leq b\}$  (half-open interval)

$[a, b) := \{x \in \mathbb{R} : a \leq x < b\}$  (half-open interval)

Those were bounded intervals. Also define unbounded intervals:

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$



For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$

$$(a, \infty) := \{x \in \mathbb{R} : a < x\}$$

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$

$$(a, \infty) := \{x \in \mathbb{R} : a < x\}$$

$$(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}$$

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$

$$(a, \infty) := \{x \in \mathbb{R} : a < x\}$$

$$(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}$$

$$(-\infty, b) := \{x \in \mathbb{R} : x < b\}$$

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$

$$(a, \infty) := \{x \in \mathbb{R} : a < x\}$$

$$(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}$$

$$(-\infty, b) := \{x \in \mathbb{R} : x < b\}$$

$$(-\infty, \infty) := \mathbb{R}$$

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$

$$(a, \infty) := \{x \in \mathbb{R} : a < x\}$$

$$(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}$$

$$(-\infty, b) := \{x \in \mathbb{R} : x < b\}$$

$$(-\infty, \infty) := \mathbb{R}$$

## Proposition

*A set  $I \subset \mathbb{R}$  is an interval if and only if  $I$  contains at least 2 points and for all  $a, c \in I$  and  $b \in \mathbb{R}$  such that  $a < b < c$ , we have  $b \in I$ .*

For  $a < b$ , define the intervals:

$$[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\} \quad (\text{closed interval})$$

$$(a, b) := \{x \in \mathbb{R} : a < x < b\} \quad (\text{open interval})$$

$$(a, b] := \{x \in \mathbb{R} : a < x \leq b\} \quad (\text{half-open interval})$$

$$[a, b) := \{x \in \mathbb{R} : a \leq x < b\} \quad (\text{half-open interval})$$

Those were bounded intervals. Also define unbounded intervals:

$$[a, \infty) := \{x \in \mathbb{R} : a \leq x\}$$

$$(a, \infty) := \{x \in \mathbb{R} : a < x\}$$

$$(-\infty, b] := \{x \in \mathbb{R} : x \leq b\}$$

$$(-\infty, b) := \{x \in \mathbb{R} : x < b\}$$

$$(-\infty, \infty) := \mathbb{R}$$

## Proposition

*A set  $I \subset \mathbb{R}$  is an interval if and only if  $I$  contains at least 2 points and for all  $a, c \in I$  and  $b \in \mathbb{R}$  such that  $a < b < c$ , we have  $b \in I$ .*

Proof is an exercise.

Set theoretically intervals have the same size.

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$



Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Bijection from  $[0, 1]$  to  $(0, 1)$  is harder but possible.

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Bijection from  $[0, 1]$  to  $(0, 1)$  is harder but possible.

We saw that uncountable sets exist: e.g.,  $\mathcal{P}(\mathbb{N})$ .

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Bijection from  $[0, 1]$  to  $(0, 1)$  is harder but possible.

We saw that uncountable sets exist: e.g.,  $\mathcal{P}(\mathbb{N})$ .

### Theorem (Cantor)

$\mathbb{R}$  is uncountable.

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Bijection from  $[0, 1]$  to  $(0, 1)$  is harder but possible.

We saw that uncountable sets exist: e.g.,  $\mathcal{P}(\mathbb{N})$ .

### Theorem (Cantor)

$\mathbb{R}$  is uncountable.

So any interval is uncountable.

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Bijection from  $[0, 1]$  to  $(0, 1)$  is harder but possible.

We saw that uncountable sets exist: e.g.,  $\mathcal{P}(\mathbb{N})$ .

### Theorem (Cantor)

$\mathbb{R}$  is *uncountable*.

So any interval is uncountable.

Also, the set  $\mathbb{R} \setminus \mathbb{Q}$  is uncountable.

Set theoretically intervals have the same size.

$f(x) := 2x$  is a bijection from  $[0, 1]$  to  $[0, 2]$

$f(x) := \tan(x)$  is a bijection from  $(-\pi/2, \pi/2)$  to  $\mathbb{R}$

Bijection from  $[0, 1]$  to  $(0, 1)$  is harder but possible.

We saw that uncountable sets exist: e.g.,  $\mathcal{P}(\mathbb{N})$ .

### Theorem (Cantor)

$\mathbb{R}$  is *uncountable*.

So any interval is uncountable.

Also, the set  $\mathbb{R} \setminus \mathbb{Q}$  is uncountable.

We'll give essentially Cantor's original 1874 proof.

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ .



**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k - 1$ ,

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .



**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .

Set  $a_k := x_n$ , where  $n$  is the smallest  $n \in \mathbb{N}$  such that  $x_n \in (a_{k-1}, b_{k-1})$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .

Set  $a_k := x_n$ , where  $n$  is the smallest  $n \in \mathbb{N}$  such that  $x_n \in (a_{k-1}, b_{k-1})$ .

$x_n$  exists by assumption on  $X$ .  $n \geq k$  by assumption on  $(a_{k-1}, b_{k-1})$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .

Set  $a_k := x_n$ , where  $n$  is the smallest  $n \in \mathbb{N}$  such that  $x_n \in (a_{k-1}, b_{k-1})$ .

$x_n$  exists by assumption on  $X$ .  $n \geq k$  by assumption on  $(a_{k-1}, b_{k-1})$ .

Define  $b_k$  to be any real number in  $(a_k, b_{k-1})$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .

Set  $a_k := x_n$ , where  $n$  is the smallest  $n \in \mathbb{N}$  such that  $x_n \in (a_{k-1}, b_{k-1})$ .

$x_n$  exists by assumption on  $X$ .  $n \geq k$  by assumption on  $(a_{k-1}, b_{k-1})$ .

Define  $b_k$  to be any real number in  $(a_k, b_{k-1})$ .

Note  $a_{k-1} < a_k < b_k < b_{k-1}$ , and

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .

Set  $a_k := x_n$ , where  $n$  is the smallest  $n \in \mathbb{N}$  such that  $x_n \in (a_{k-1}, b_{k-1})$ .

$x_n$  exists by assumption on  $X$ .  $n \geq k$  by assumption on  $(a_{k-1}, b_{k-1})$ .

Define  $b_k$  to be any real number in  $(a_k, b_{k-1})$ .

Note  $a_{k-1} < a_k < b_k < b_{k-1}$ , and

$x_k \notin (a_k, b_k)$  and hence  $x_j \notin (a_k, b_k)$  for  $x_j$  for  $j = 1, 2, \dots, k$ .

**Proof:** Suppose  $X \subset \mathbb{R}$  is countably infinite such that for every pair of real numbers  $a < b$ , there is an  $x \in X$  such that  $a < x < b$ . Were  $\mathbb{R}$  countable, we could take  $X = \mathbb{R}$ . We'll show  $X \subsetneq \mathbb{R}$ .

Write  $X = \{x_1, x_2, x_3, \dots\}$  ( $X$  is countably infinite).

We construct two sequences  $a_1, a_2, a_3, \dots$  and  $b_1, b_2, b_3, \dots$ .

Let  $a_1 := x_1$  and  $b_1 := x_1 + 1$ .

Note that  $a_1 < b_1$  and  $x_1 \notin (a_1, b_1)$ .

Suppose for some  $k > 1$ ,  $a_j$  and  $b_j$  have been defined for  $j = 1, 2, \dots, k-1$ , for each such  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ , and suppose  $a_1 < a_2 < \dots < a_{k-1} < b_{k-1} < \dots < b_2 < b_1$ .

Set  $a_k := x_n$ , where  $n$  is the smallest  $n \in \mathbb{N}$  such that  $x_n \in (a_{k-1}, b_{k-1})$ .

$x_n$  exists by assumption on  $X$ .  $n \geq k$  by assumption on  $(a_{k-1}, b_{k-1})$ .

Define  $b_k$  to be any real number in  $(a_k, b_{k-1})$ .

Note  $a_{k-1} < a_k < b_k < b_{k-1}$ , and

$x_k \notin (a_k, b_k)$  and hence  $x_j \notin (a_k, b_k)$  for  $x_j$  for  $j = 1, 2, \dots, k$ .

The two sequences are now defined.

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .



We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m$

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .  
By the claim,  $\sup A \leq \inf B$ .

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \implies a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

$y \notin B$ : Similar.



We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \Rightarrow a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

$y \notin B$ : Similar.

So  $a_n < y < b_n$  for all  $n \in \mathbb{N}$  or  $y \in (a_n, b_n)$  for all  $n \in \mathbb{N}$ .

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \Rightarrow a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

$y \notin B$ : Similar.

So  $a_n < y < b_n$  for all  $n \in \mathbb{N}$  or  $y \in (a_n, b_n)$  for all  $n \in \mathbb{N}$ .

For every  $n \in \mathbb{N}$ ,  $x_n \notin (a_n, b_n)$ , so  $y \neq x_n$ .

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \Rightarrow a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

$y \notin B$ : Similar.

So  $a_n < y < b_n$  for all  $n \in \mathbb{N}$  or  $y \in (a_n, b_n)$  for all  $n \in \mathbb{N}$ .

For every  $n \in \mathbb{N}$ ,  $x_n \notin (a_n, b_n)$ , so  $y \neq x_n$ .

$\Rightarrow y \notin X$

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \Rightarrow a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

$y \notin B$ : Similar.

So  $a_n < y < b_n$  for all  $n \in \mathbb{N}$  or  $y \in (a_n, b_n)$  for all  $n \in \mathbb{N}$ .

For every  $n \in \mathbb{N}$ ,  $x_n \notin (a_n, b_n)$ , so  $y \neq x_n$ .

$\Rightarrow y \notin X \Rightarrow X \subsetneq \mathbb{R}$

We have two sequences such that  
for each  $j$ , suppose  $x_\ell \notin (a_j, b_j)$  for  $\ell = 1, 2, \dots, j$ ,  
and  $a_1 < a_2 < \dots < a_k < b_k < \dots < b_2 < b_1$  (for every  $k$ ).

**Claim:**  $a_n < b_m$  for all  $n$  and  $m$  in  $\mathbb{N}$ .

**Proof of claim:** Suppose  $n < m \Rightarrow a_n < a_{n+1} < \dots < a_{m-1} < a_m < b_m$ .  
Similarly for  $n > m$ . The claim follows.

Let  $A := \{a_n : n \in \mathbb{N}\}$  and  $B := \{b_n : n \in \mathbb{N}\}$ .

By the claim,  $\sup A \leq \inf B$ .

Define  $y := \sup A$ .

$y \notin A$ : If  $y = a_n$  for some  $n$ , then  $y < a_{n+1}$ , which is impossible.

$y \notin B$ : Similar.

So  $a_n < y < b_n$  for all  $n \in \mathbb{N}$  or  $y \in (a_n, b_n)$  for all  $n \in \mathbb{N}$ .

For every  $n \in \mathbb{N}$ ,  $x_n \notin (a_n, b_n)$ , so  $y \neq x_n$ .

$\Rightarrow y \notin X \Rightarrow X \subsetneq \mathbb{R} \Rightarrow \mathbb{R}$  is uncountable. □

**Remark:** Cantor's paper was (perhaps oddly) about algebraic numbers:

**Remark:** Cantor's paper was (perhaps oddly) about algebraic numbers:

$x \in \mathbb{R}$  is *algebraic* if  $x$  is a root of a polynomial with integer coefficients:  
 $a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0 = 0$  where  $a_0, a_1, \dots, a_n \in \mathbb{Z}$ .

**Remark:** Cantor's paper was (perhaps oddly) about algebraic numbers:

$x \in \mathbb{R}$  is *algebraic* if  $x$  is a root of a polynomial with integer coefficients:  
 $a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0 = 0$  where  $a_0, a_1, \dots, a_n \in \mathbb{Z}$ .

There are only countably many algebraic numbers (exercise).



**Remark:** Cantor's paper was (perhaps oddly) about algebraic numbers:

$x \in \mathbb{R}$  is *algebraic* if  $x$  is a root of a polynomial with integer coefficients:  
 $a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0 = 0$  where  $a_0, a_1, \dots, a_n \in \mathbb{Z}$ .

There are only countably many algebraic numbers (exercise).

Cantor's theorem shows  $\exists$  non-algebraic (transcendental) numbers.