

BA: 5.1

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

An integral is a way to “sum” or “average” the values of a function.

An integral is a way to “sum” or “average” the values of a function.

Integral is *not* the antiderivative!

An integral is a way to “sum” or “average” the values of a function.

Integral is *not* the antiderivative!

An antiderivative can be computed via an integral,

An integral is a way to “sum” or “average” the values of a function.

Integral is *not* the antiderivative!

An antiderivative can be computed via an integral,
but an integral does a lot more.

An integral is a way to “sum” or “average” the values of a function.

Integral is *not* the antiderivative!

An antiderivative can be computed via an integral,
but an integral does a lot more.

In our setting, an integral is the “area under the curve”

An integral is a way to “sum” or “average” the values of a function.

Integral is *not* the antiderivative!

An antiderivative can be computed via an integral,
but an integral does a lot more.

In our setting, an integral is the “area under the curve”

We will follow the Darboux approach rather than Riemann approach,
they are equivalent.

Definition

A *partition* P of the interval $[a, b]$ is a finite set $\{x_0, x_1, x_2, \dots, x_n\}$ such that

$$a = x_0 < x_1 < x_2 < \cdots < x_{n-1} < x_n = b.$$

Definition

A *partition* P of the interval $[a, b]$ is a finite set $\{x_0, x_1, x_2, \dots, x_n\}$ such that

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

We write

$$\Delta x_i := x_i - x_{i-1}.$$

Definition

A *partition* P of the interval $[a, b]$ is a finite set $\{x_0, x_1, x_2, \dots, x_n\}$ such that

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

We write

$$\Delta x_i := x_i - x_{i-1}.$$

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function.

Definition

A *partition* P of the interval $[a, b]$ is a finite set $\{x_0, x_1, x_2, \dots, x_n\}$ such that

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

We write

$$\Delta x_i := x_i - x_{i-1}.$$

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let P be a partition of $[a, b]$.

Definition

A *partition* P of the interval $[a, b]$ is a finite set $\{x_0, x_1, x_2, \dots, x_n\}$ such that

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

We write

$$\Delta x_i := x_i - x_{i-1}.$$

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let P be a partition of $[a, b]$. Define

$$m_i := \inf \{f(x) : x_{i-1} \leq x \leq x_i\},$$

$$M_i := \sup \{f(x) : x_{i-1} \leq x \leq x_i\},$$

$$L(P, f) := \sum_{i=1}^n m_i \Delta x_i,$$

$$U(P, f) := \sum_{i=1}^n M_i \Delta x_i.$$

Definition

A *partition* P of the interval $[a, b]$ is a finite set $\{x_0, x_1, x_2, \dots, x_n\}$ such that

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$


We write

$$\Delta x_i := x_i - x_{i-1}.$$

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let P be a partition of $[a, b]$. Define

$$\begin{aligned} m_i &:= \inf \{f(x) : x_{i-1} \leq x \leq x_i\}, & M_i &:= \sup \{f(x) : x_{i-1} \leq x \leq x_i\}, \\ L(P, f) &:= \sum_{i=1}^n m_i \Delta x_i, & U(P, f) &:= \sum_{i=1}^n M_i \Delta x_i. \end{aligned}$$

We call $L(P, f)$ the *lower Darboux sum* and $U(P, f)$ the *upper Darboux sum*.



Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.
 $m \leq m_i$ for all i and $M_i \leq M$ for all i .

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.
 $m \leq m_i$ for all i and $M_i \leq M$ for all i .
 $m_i \leq M_i$ for all i .

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$\sum_{i=1}^n \Delta x_i = (b - a)$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b - a).$$

$$\Rightarrow m(b - a)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b - a).$$

$$\Rightarrow m(b - a) = m \left(\sum_{i=1}^n \Delta x_i \right)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b - a).$$

$$\Rightarrow m(b - a) = m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b - a).$$

$$\Rightarrow m(b - a) = m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b - a) \leq L(P, f) \leq U(P, f) \leq M(b - a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b - a).$$

$$\begin{aligned} \Rightarrow m(b - a) &= m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i \\ &\leq \sum_{i=1}^n M_i \Delta x_i \end{aligned}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b-a).$$

$$\begin{aligned} \Rightarrow m(b-a) &= m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i \\ &\leq \sum_{i=1}^n M_i \Delta x_i \leq \sum_{i=1}^n M \Delta x_i \end{aligned}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b-a).$$

$$\Rightarrow m(b-a) = m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i$$

$$\leq \sum_{i=1}^n M_i \Delta x_i \leq \sum_{i=1}^n M \Delta x_i = M \left(\sum_{i=1}^n \Delta x_i \right)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b-a).$$

$$\Rightarrow m(b-a) = m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i$$

$$\leq \sum_{i=1}^n M_i \Delta x_i \leq \sum_{i=1}^n M \Delta x_i = M \left(\sum_{i=1}^n \Delta x_i \right) = M(b-a). \quad \square$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then for every partition P of $[a, b]$,

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

Proof: Let P be a partition of $[a, b]$.

$m \leq m_i$ for all i and $M_i \leq M$ for all i .

$m_i \leq M_i$ for all i .

$$\sum_{i=1}^n \Delta x_i = (b-a).$$

$$\Rightarrow m(b-a) = m \left(\sum_{i=1}^n \Delta x_i \right) = \sum_{i=1}^n m \Delta x_i \leq \sum_{i=1}^n m_i \Delta x_i$$

$$\leq \sum_{i=1}^n M_i \Delta x_i \leq \sum_{i=1}^n M \Delta x_i = M \left(\sum_{i=1}^n \Delta x_i \right) = M(b-a). \quad \square$$

In particular, the set of lower and upper sums are bounded sets.

Definition

As the sets of lower and upper Darboux sums are bounded, we define

Definition

As the sets of lower and upper Darboux sums are bounded, we define

$$\underline{\int_a^b} f(x) dx := \sup \{L(P, f) : P \text{ a partition of } [a, b]\},$$

Definition

As the sets of lower and upper Darboux sums are bounded, we define

$$\underline{\int_a^b} f(x) dx := \sup \{L(P, f) : P \text{ a partition of } [a, b]\},$$

$$\overline{\int_a^b} f(x) dx := \inf \{U(P, f) : P \text{ a partition of } [a, b]\}.$$

Definition

As the sets of lower and upper Darboux sums are bounded, we define

$$\underline{\int_a^b} f(x) dx := \sup \{L(P, f) : P \text{ a partition of } [a, b]\},$$

$$\overline{\int_a^b} f(x) dx := \inf \{U(P, f) : P \text{ a partition of } [a, b]\}.$$

We call $\underline{\int}$ the *lower Darboux integral* and $\overline{\int}$ the *upper Darboux integral*.

Definition

As the sets of lower and upper Darboux sums are bounded, we define

$$\underline{\int_a^b} f(x) dx := \sup \{L(P, f) : P \text{ a partition of } [a, b]\},$$

$$\overline{\int_a^b} f(x) dx := \inf \{U(P, f) : P \text{ a partition of } [a, b]\}.$$

We call $\underline{\int}$ the *lower Darboux integral* and $\overline{\int}$ the *upper Darboux integral*. To avoid worrying about the variable of integration, we often simply write

$$\underline{\int_a^b} f := \underline{\int_a^b} f(x) dx \quad \text{and} \quad \overline{\int_a^b} f := \overline{\int_a^b} f(x) dx.$$

Definition

As the sets of lower and upper Darboux sums are bounded, we define

$$\underline{\int_a^b} f(x) dx := \sup \{L(P, f) : P \text{ a partition of } [a, b]\},$$

$$\overline{\int_a^b} f(x) dx := \inf \{U(P, f) : P \text{ a partition of } [a, b]\}.$$

We call $\underline{\int}$ the *lower Darboux integral* and $\overline{\int}$ the *upper Darboux integral*. To avoid worrying about the variable of integration, we often simply write

$$\underline{\int_a^b} f := \underline{\int_a^b} f(x) dx \quad \text{and} \quad \overline{\int_a^b} f := \overline{\int_a^b} f(x) dx.$$

Remark: If $f: S \rightarrow \mathbb{R}$ and $[a, b] \subset S$, use $f|_{[a, b]}$. But we'll just write $\underline{\int_a^b} f$ and $\overline{\int_a^b} f$.

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$.

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$. Then

$$\underline{\int_0^1} f = 0 \quad \text{and} \quad \overline{\int_0^1} f = 1.$$

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$. Then

$$\underline{\int_0^1 f} = 0 \quad \text{and} \quad \overline{\int_0^1 f} = 1.$$

Proof: Take any partition P .

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$. Then

$$\underline{\int_0^1} f = 0 \quad \text{and} \quad \overline{\int_0^1} f = 1.$$

Proof: Take any partition P . For every i ,
 $m_i = \inf\{f(x) : x \in [x_{i-1}, x_i]\} = 0$

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$. Then

$$\underline{\int_0^1} f = 0 \quad \text{and} \quad \overline{\int_0^1} f = 1.$$

Proof: Take any partition P . For every i ,
 $m_i = \inf\{f(x) : x \in [x_{i-1}, x_i]\} = 0$ and $M_i = \sup\{f(x) : x \in [x_{i-1}, x_i]\} = 1$.

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$. Then

$$\underline{\int_0^1} f = 0 \quad \text{and} \quad \overline{\int_0^1} f = 1.$$

Proof: Take any partition P . For every i ,
 $m_i = \inf\{f(x) : x \in [x_{i-1}, x_i]\} = 0$ and $M_i = \sup\{f(x) : x \in [x_{i-1}, x_i]\} = 1$.

$$\Rightarrow L(P, f) = \sum_{i=1}^n 0 \cdot \Delta x_i = 0,$$

Example: Take the Dirichlet function $f: [0, 1] \rightarrow \mathbb{R}$, where $f(x) := 1$ if $x \in \mathbb{Q}$ and $f(x) := 0$ if $x \notin \mathbb{Q}$. Then

$$\underline{\int_0^1 f} = 0 \quad \text{and} \quad \overline{\int_0^1 f} = 1.$$

Proof: Take any partition P . For every i ,
 $m_i = \inf\{f(x) : x \in [x_{i-1}, x_i]\} = 0$ and $M_i = \sup\{f(x) : x \in [x_{i-1}, x_i]\} = 1$.

$$\Rightarrow L(P, f) = \sum_{i=1}^n 0 \cdot \Delta x_i = 0, \quad \text{and} \quad U(P, f) = \sum_{i=1}^n 1 \cdot \Delta x_i = \sum_{i=1}^n \Delta x_i = 1.$$

□

Definition

Let $P = \{x_0, x_1, \dots, x_n\}$ and $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ be partitions of $[a, b]$.

Definition

Let $P = \{x_0, x_1, \dots, x_n\}$ and $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ be partitions of $[a, b]$.
We say \tilde{P} is a *refinement* of P if as sets $P \subset \tilde{P}$.

Definition

Let $P = \{x_0, x_1, \dots, x_n\}$ and $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ be partitions of $[a, b]$.
We say \tilde{P} is a *refinement* of P if as sets $P \subset \tilde{P}$.

Example: $\{0, 0.5, 1, 2\}$ is a partition of $[0, 2]$
and $\{0, 0.2, 0.5, 1, 1.5, 1.75, 2\}$ is its refinement.

Definition

Let $P = \{x_0, x_1, \dots, x_n\}$ and $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ be partitions of $[a, b]$.
We say \tilde{P} is a *refinement* of P if as sets $P \subset \tilde{P}$.

Example: $\{0, 0.5, 1, 2\}$ is a partition of $[0, 2]$
and $\{0, 0.2, 0.5, 1, 1.5, 1.75, 2\}$ is its refinement.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P .

Definition

Let $P = \{x_0, x_1, \dots, x_n\}$ and $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ be partitions of $[a, b]$.
We say \tilde{P} is a *refinement* of P if as sets $P \subset \tilde{P}$.

Example: $\{0, 0.5, 1, 2\}$ is a partition of $[0, 2]$
and $\{0, 0.2, 0.5, 1, 1.5, 1.75, 2\}$ is its refinement.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.
 $\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.

$\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.

$\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then


$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.

$\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.



Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then


$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.

$\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.



Δx_i

Proposition


Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.
 $\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.



$$\Delta x_i = x_i - x_{i-1}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then


$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.

$\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.



$$\Delta x_i = x_i - x_{i-1} = \tilde{x}_{k_i} - \tilde{x}_{k_{i-1}}$$

Proposition


Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$

Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.
 $\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.

There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.



$$\Delta x_i = x_i - x_{i-1} = \tilde{x}_{k_i} - \tilde{x}_{k_{i-1}} = \sum_{q=k_{i-1}+1}^{k_i} \tilde{x}_q - \tilde{x}_{q-1}$$

Proposition


Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded, P a partition of $[a, b]$, \tilde{P} a refinement of P . Then

$$L(P, f) \leq L(\tilde{P}, f) \quad \text{and} \quad U(\tilde{P}, f) \leq U(P, f).$$


Proof: Suppose $\tilde{P} = \{\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_\ell\}$ is a refinement of $P = \{x_0, x_1, \dots, x_n\}$.
 $\Rightarrow x_0 = \tilde{x}_0$ and $x_n = \tilde{x}_\ell$.


There are integers $k_0 < k_1 < \dots < k_n$ such that $x_i = \tilde{x}_{k_i}$ for $i = 0, 1, 2, \dots, n$.

Let $\Delta x_i := x_i - x_{i-1}$ and $\Delta \tilde{x}_q := \tilde{x}_q - \tilde{x}_{q-1}$.




$$\Delta x_i = x_i - x_{i-1} = \tilde{x}_{k_i} - \tilde{x}_{k_{i-1}} = \sum_{q=k_{i-1}+1}^{k_i} \tilde{x}_q - \tilde{x}_{q-1} = \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q.$$






Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.



Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.


$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.



Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.


$$\Rightarrow m_i \Delta x_i$$



Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.


$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q$$



Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.


$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q$$



Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$




Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$

$$\Rightarrow L(P, f)$$




Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$

$$\Rightarrow L(P, f) = \sum_{i=1}^n m_i \Delta x_i$$




Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$

$$\Rightarrow L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q$$




Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$

$$\Rightarrow L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q = \sum_{q=1}^{\ell} \tilde{m}_q \Delta \tilde{x}_q$$




Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$

$$\Rightarrow L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q = \sum_{q=1}^{\ell} \tilde{m}_q \Delta \tilde{x}_q = L(\tilde{P}, f).$$



Let $m_i := \inf\{f(x) : x_{i-1} \leq x \leq x_i\}$ and $\tilde{m}_q := \inf\{f(x) : \tilde{x}_{q-1} \leq x \leq \tilde{x}_q\}$.

$m_i \leq \tilde{m}_q$ for $k_{i-1} < q \leq k_i$.

$$\Rightarrow m_i \Delta x_i = m_i \sum_{q=k_{i-1}+1}^{k_i} \Delta \tilde{x}_q = \sum_{q=k_{i-1}+1}^{k_i} m_i \Delta \tilde{x}_q \leq \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q.$$

$$\Rightarrow L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \sum_{q=k_{i-1}+1}^{k_i} \tilde{m}_q \Delta \tilde{x}_q = \sum_{q=1}^{\ell} \tilde{m}_q \Delta \tilde{x}_q = L(\tilde{P}, f).$$

$U(\tilde{P}, f) \leq U(P, f)$ is left as an exercise.

□

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f.$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b} f \leq M(b-a).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \int_a^b f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b f} \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \int_a^b f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b f} \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \int_a^b f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b f} \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b} f \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \int_a^b f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b f} \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f) \leq U(\tilde{P}, f)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \int_a^b f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b f} \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f) \leq U(\tilde{P}, f) \leq U(P_2, f).$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b} f \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f) \leq U(\tilde{P}, f) \leq U(P_2, f).$$

$$\Rightarrow \underline{\int_a^b} f$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b} f \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f) \leq U(\tilde{P}, f) \leq U(P_2, f).$$

$$\Rightarrow \underline{\int_a^b} f = \sup \{L(P, f) : P \text{ a partition of } [a, b]\}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b} f \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f) \leq U(\tilde{P}, f) \leq U(P_2, f).$$

$$\begin{aligned} \Rightarrow \underline{\int_a^b} f &= \sup \{L(P, f) : P \text{ a partition of } [a, b]\} \\ &\leq \inf \{U(P, f) : P \text{ a partition of } [a, b]\} \end{aligned}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Let $m, M \in \mathbb{R}$ be such that for all $x \in [a, b]$, we have $m \leq f(x) \leq M$. Then

$$m(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq M(b-a).$$

Proof: For any partition P , we proved

$$m(b-a) \leq L(P, f) \leq U(P, f) \leq M(b-a).$$

$$m(b-a) \leq L(P, f) \Rightarrow m(b-a) \leq \underline{\int_a^b} f. \quad U(P, f) \leq M(b-a) \Rightarrow \overline{\int_a^b} f \leq M(b-a).$$

Let P_1, P_2 be partitions of $[a, b]$. Define $\tilde{P} := P_1 \cup P_2$.

\tilde{P} is a refinement of both P_1 and P_2

$$\Rightarrow L(P_1, f) \leq L(\tilde{P}, f) \leq U(\tilde{P}, f) \leq U(P_2, f).$$

$$\begin{aligned} \Rightarrow \underline{\int_a^b} f &= \sup \{L(P, f) : P \text{ a partition of } [a, b]\} \\ &\leq \inf \{U(P, f) : P \text{ a partition of } [a, b]\} = \overline{\int_a^b} f. \end{aligned}$$

□

Definition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded such that

$$\underline{\int_a^b} f(x) dx = \overline{\int_a^b} f(x) dx.$$

Definition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded such that

$$\underline{\int_a^b} f(x) dx = \overline{\int_a^b} f(x) dx.$$

Then f is said to be *Riemann integrable*.

Definition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded such that

$$\underline{\int_a^b} f(x) dx = \overline{\int_a^b} f(x) dx.$$

Then f is said to be *Riemann integrable*. The set of Riemann integrable functions on $[a, b]$ is denoted by $\mathcal{R}[a, b]$.

Definition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded such that

$$\int_a^b f(x) dx = \overline{\int_a^b f(x) dx}.$$

Then f is said to be *Riemann integrable*. The set of Riemann integrable functions on $[a, b]$ is denoted by $\mathcal{R}[a, b]$. When $f \in \mathcal{R}[a, b]$, we define

$$\int_a^b f(x) dx := \int_a^b f(x) dx = \overline{\int_a^b f(x) dx}.$$

Definition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded such that

$$\int_a^b f(x) dx = \overline{\int_a^b f(x) dx}.$$

Then f is said to be *Riemann integrable*. The set of Riemann integrable functions on $[a, b]$ is denoted by $\mathcal{R}[a, b]$. When $f \in \mathcal{R}[a, b]$, we define

$$\int_a^b f(x) dx := \int_a^b f(x) dx = \overline{\int_a^b f(x) dx}.$$

We often write

$$\int_a^b f := \int_a^b f(x) dx.$$

Definition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded such that

$$\int_a^b f(x) dx = \overline{\int_a^b f(x) dx}.$$

Then f is said to be *Riemann integrable*. The set of Riemann integrable functions on $[a, b]$ is denoted by $\mathcal{R}[a, b]$. When $f \in \mathcal{R}[a, b]$, we define

$$\int_a^b f(x) dx := \int_a^b f(x) dx = \overline{\int_a^b f(x) dx}.$$

We often write

$$\int_a^b f := \int_a^b f(x) dx.$$

$\int_a^b f$ is called the *Riemann integral* of f , or simply the *integral* of f .

We immediately get:

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be Riemann integrable. Let $m, M \in \mathbb{R}$ be such that $m \leq f(x) \leq M$ for all $x \in [a, b]$. Then


$$m(b - a) \leq \int_a^b f \leq M(b - a).$$

We immediately get:

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be Riemann integrable. Let $m, M \in \mathbb{R}$ be such that $m \leq f(x) \leq M$ for all $x \in [a, b]$. Then

$$m(b - a) \leq \int_a^b f \leq M(b - a).$$




We immediately get:

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be Riemann integrable. Let $m, M \in \mathbb{R}$ be such that $m \leq f(x) \leq M$ for all $x \in [a, b]$. Then

$$m(b-a) \leq \int_a^b f \leq M(b-a).$$



Useful version: If $|f(x)| \leq M$ for all $x \in [a, b]$, then $\left| \int_a^b f \right| \leq M(b-a).$

Example: Constant function are integrable:

Example: Constant function are integrable:

If $f(x) = c$ for some constant c , then c is an upper and lower bound,

Example: Constant function are integrable:

If $f(x) = c$ for some constant c , then c is an upper and lower bound, so

$$c(b-a) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq c(b-a).$$

Example: Constant functions are integrable:

If $f(x) = c$ for some constant c , then c is an upper and lower bound, so

$$c(b-a) \leq \int_a^b f \leq \overline{\int_a^b f} \leq c(b-a).$$

So f is integrable on any interval $[a, b]$ and

$$\int_a^b f = c(b-a).$$

Example: Let $f: [0, 2] \rightarrow \mathbb{R}$ be defined by

$$f(x) := \begin{cases} 1 & \text{if } x < 1, \\ 1/2 & \text{if } x = 1, \\ 0 & \text{if } x > 1. \end{cases}$$

Example: Let $f: [0, 2] \rightarrow \mathbb{R}$ be defined by

$$f(x) := \begin{cases} 1 & \text{if } x < 1, \\ 1/2 & \text{if } x = 1, \\ 0 & \text{if } x > 1. \end{cases}$$

We claim f is Riemann integrable and $\int_0^2 f = 1$.

Example: Let $f: [0, 2] \rightarrow \mathbb{R}$ be defined by

$$f(x) := \begin{cases} 1 & \text{if } x < 1, \\ 1/2 & \text{if } x = 1, \\ 0 & \text{if } x > 1. \end{cases}$$

We claim f is Riemann integrable and $\int_0^2 f = 1$.

Proof: Let $0 < \epsilon < 1$ be arbitrary.

Example: Let $f: [0, 2] \rightarrow \mathbb{R}$ be defined by

$$f(x) := \begin{cases} 1 & \text{if } x < 1, \\ 1/2 & \text{if } x = 1, \\ 0 & \text{if } x > 1. \end{cases}$$

We claim f is Riemann integrable and $\int_0^2 f = 1$.

Proof: Let $0 < \epsilon < 1$ be arbitrary.

Let $P := \{0, 1 - \epsilon, 1 + \epsilon, 2\}$ be a partition.

Example: Let $f: [0, 2] \rightarrow \mathbb{R}$ be defined by

$$f(x) := \begin{cases} 1 & \text{if } x < 1, \\ 1/2 & \text{if } x = 1, \\ 0 & \text{if } x > 1. \end{cases}$$

We claim f is Riemann integrable and $\int_0^2 f = 1$.

Proof: Let $0 < \epsilon < 1$ be arbitrary.

Let $P := \{0, 1 - \epsilon, 1 + \epsilon, 2\}$ be a partition.

Using the notation from before:

$$m_1 = \inf \{f(x) : x \in [0, 1 - \epsilon]\} = 1,$$

$$m_2 = \inf \{f(x) : x \in [1 - \epsilon, 1 + \epsilon]\} = 0,$$

$$m_3 = \inf \{f(x) : x \in [1 + \epsilon, 2]\} = 0,$$

$$M_1 = \sup \{f(x) : x \in [0, 1 - \epsilon]\} = 1,$$

$$M_2 = \sup \{f(x) : x \in [1 - \epsilon, 1 + \epsilon]\} = 1,$$

$$M_3 = \sup \{f(x) : x \in [1 + \epsilon, 2]\} = 0.$$

Example: Let $f: [0, 2] \rightarrow \mathbb{R}$ be defined by

$$f(x) := \begin{cases} 1 & \text{if } x < 1, \\ 1/2 & \text{if } x = 1, \\ 0 & \text{if } x > 1. \end{cases}$$

We claim f is Riemann integrable and $\int_0^2 f = 1$.

Proof: Let $0 < \epsilon < 1$ be arbitrary.

Let $P := \{0, 1 - \epsilon, 1 + \epsilon, 2\}$ be a partition.

Using the notation from before:

$$m_1 = \inf \{f(x) : x \in [0, 1 - \epsilon]\} = 1,$$

$$M_1 = \sup \{f(x) : x \in [0, 1 - \epsilon]\} = 1,$$

$$m_2 = \inf \{f(x) : x \in [1 - \epsilon, 1 + \epsilon]\} = 0,$$

$$M_2 = \sup \{f(x) : x \in [1 - \epsilon, 1 + \epsilon]\} = 1,$$


$$m_3 = \inf \{f(x) : x \in [1 + \epsilon, 2]\} = 0,$$


$$M_3 = \sup \{f(x) : x \in [1 + \epsilon, 2]\} = 0.$$

Furthermore, $\Delta x_1 = 1 - \epsilon$, $\Delta x_2 = 2\epsilon$, and $\Delta x_3 = 1 - \epsilon$.


$$M_1 = M_2 = m_1 = 1$$

$$M_3 = m_2 = m_3 = 0$$







$$L(P, f)$$




$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i$$



$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon)$$




$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$




$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$

$$U(P, f)$$




$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$

$$U(P, f) = \sum_{i=1}^3 M_i \Delta x_i$$




$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$

$$U(P, f) = \sum_{i=1}^3 M_i \Delta x_i = 1 \cdot (1 - \epsilon) + 1 \cdot 2\epsilon + 0 \cdot (1 - \epsilon)$$



$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$


$$U(P, f) = \sum_{i=1}^3 M_i \Delta x_i = 1 \cdot (1 - \epsilon) + 1 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 + \epsilon.$$



$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$

$$U(P, f) = \sum_{i=1}^3 M_i \Delta x_i = 1 \cdot (1 - \epsilon) + 1 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 + \epsilon.$$


$$\Rightarrow \overline{\int_0^2 f} - \underline{\int_0^2 f}$$



$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$

$$U(P, f) = \sum_{i=1}^3 M_i \Delta x_i = 1 \cdot (1 - \epsilon) + 1 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 + \epsilon.$$

$$\Rightarrow \overline{\int_0^2 f} - \underline{\int_0^2 f} \leq U(P, f) - L(P, f)$$



$$L(P, f) = \sum_{i=1}^3 m_i \Delta x_i = 1 \cdot (1 - \epsilon) + 0 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 - \epsilon,$$

$$U(P, f) = \sum_{i=1}^3 M_i \Delta x_i = 1 \cdot (1 - \epsilon) + 1 \cdot 2\epsilon + 0 \cdot (1 - \epsilon) = 1 + \epsilon.$$

$$\Rightarrow \overline{\int_0^2 f} - \underline{\int_0^2 f} \leq U(P, f) - L(P, f) = (1 + \epsilon) - (1 - \epsilon) = 2\epsilon.$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

As ϵ was arbitrary

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \quad \Rightarrow \quad \overline{\int_0^2 f} = \underline{\int_0^2 f}$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f)$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f) \leq \int_0^2 f$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f) \leq \int_0^2 f \leq U(P, f)$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f) \leq \int_0^2 f \leq U(P, f) = 1 + \epsilon$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f) \leq \int_0^2 f \leq U(P, f) = 1 + \epsilon \Rightarrow \left| \int_0^2 f - 1 \right| \leq \epsilon.$$

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f) \leq \int_0^2 f \leq U(P, f) = 1 + \epsilon \Rightarrow \left| \int_0^2 f - 1 \right| \leq \epsilon.$$

As ϵ was arbitrary

$$\overline{\int_0^2 f} - \underline{\int_0^2 f} \leq 2\epsilon \quad \text{also} \quad \underline{\int_0^2 f} \leq \overline{\int_0^2 f}$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \overline{\int_0^2 f} = \underline{\int_0^2 f} \Rightarrow f \text{ is Riemann integrable.}$$

$$1 - \epsilon = L(P, f) \leq \int_0^2 f \leq U(P, f) = 1 + \epsilon \Rightarrow \left| \int_0^2 f - 1 \right| \leq \epsilon.$$

$$\text{As } \epsilon \text{ was arbitrary} \Rightarrow \int_0^2 f = 1.$$



Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then f is Riemann integrable if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(P, f) - L(P, f) < \epsilon.$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then f is Riemann integrable if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(P, f) - L(P, f) < \epsilon.$$

Proof: If for every $\epsilon > 0$ such a P exists, then

$$0 \leq \overline{\int_a^b f} - \underline{\int_a^b f}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then f is Riemann integrable if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(P, f) - L(P, f) < \epsilon.$$

Proof: If for every $\epsilon > 0$ such a P exists, then

$$0 \leq \overline{\int_a^b} f - \underline{\int_a^b} f \leq U(P, f) - L(P, f)$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then f is Riemann integrable if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(P, f) - L(P, f) < \epsilon.$$

Proof: If for every $\epsilon > 0$ such a P exists, then

$$0 \leq \overline{\int_a^b f} - \underline{\int_a^b f} \leq U(P, f) - L(P, f) < \epsilon.$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then f is Riemann integrable if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(P, f) - L(P, f) < \epsilon.$$

Proof: If for every $\epsilon > 0$ such a P exists, then

$$0 \leq \overline{\int_a^b f} - \underline{\int_a^b f} \leq U(P, f) - L(P, f) < \epsilon.$$

$$\Rightarrow \overline{\int_a^b f} = \underline{\int_a^b f}$$

Proposition

Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded. Then f is Riemann integrable if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(P, f) - L(P, f) < \epsilon.$$

Proof: If for every $\epsilon > 0$ such a P exists, then

$$0 \leq \overline{\int_a^b f} - \underline{\int_a^b f} \leq U(P, f) - L(P, f) < \epsilon.$$

$$\Rightarrow \overline{\int_a^b f} = \underline{\int_a^b f} \Rightarrow f \text{ is integrable.}$$

□

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$,

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\Rightarrow U(P, f) - L(P, f)$$

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\Rightarrow U(P, f) - L(P, f) = \sum_{i=1}^n \Delta x_i (M_i - m_i)$$

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\Rightarrow U(P, f) - L(P, f) = \sum_{i=1}^n \Delta x_i (M_i - m_i) = \frac{b}{n} \sum_{i=1}^n \left(\frac{1}{1 + (i-1)b/n} - \frac{1}{1 + ib/n} \right)$$

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\begin{aligned} \Rightarrow U(P, f) - L(P, f) &= \sum_{i=1}^n \Delta x_i (M_i - m_i) = \frac{b}{n} \sum_{i=1}^n \left(\frac{1}{1 + (i-1)b/n} - \frac{1}{1 + ib/n} \right) \\ &= \frac{b}{n} \left(\frac{1}{1 + 0b/n} - \frac{1}{1 + nb/n} \right) \end{aligned}$$

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\begin{aligned} \Rightarrow U(P, f) - L(P, f) &= \sum_{i=1}^n \Delta x_i (M_i - m_i) = \frac{b}{n} \sum_{i=1}^n \left(\frac{1}{1 + (i-1)b/n} - \frac{1}{1 + ib/n} \right) \\ &= \frac{b}{n} \left(\frac{1}{1 + 0b/n} - \frac{1}{1 + nb/n} \right) = \frac{b^2}{n(b+1)}. \end{aligned}$$

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.

$\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,

and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\begin{aligned} \Rightarrow U(P, f) - L(P, f) &= \sum_{i=1}^n \Delta x_i (M_i - m_i) = \frac{b}{n} \sum_{i=1}^n \left(\frac{1}{1 + (i-1)b/n} - \frac{1}{1 + ib/n} \right) \\ &= \frac{b}{n} \left(\frac{1}{1 + 0b/n} - \frac{1}{1 + nb/n} \right) = \frac{b^2}{n(b+1)}. \end{aligned}$$

Pick n to be such that $\frac{b^2}{n(b+1)} < \epsilon$,

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.
 $\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,
and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\begin{aligned} \Rightarrow U(P, f) - L(P, f) &= \sum_{i=1}^n \Delta x_i (M_i - m_i) = \frac{b}{n} \sum_{i=1}^n \left(\frac{1}{1 + (i-1)b/n} - \frac{1}{1 + ib/n} \right) \\ &= \frac{b}{n} \left(\frac{1}{1 + 0b/n} - \frac{1}{1 + nb/n} \right) = \frac{b^2}{n(b+1)}. \end{aligned}$$

Pick n to be such that $\frac{b^2}{n(b+1)} < \epsilon$, the proposition is satisfied,

Example: Claim: $\frac{1}{1+x}$ is integrable on $[0, b]$ for any $b > 0$.

Proof: Let $\epsilon > 0$ be given.

Take $n \in \mathbb{N}$ and let $x_i := ib/n$, to get partition $P := \{x_0, x_1, \dots, x_n\}$ of $[0, b]$.
 $\Delta x_i = b/n$ for all i .

f is decreasing so $m_i = \inf \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_i}$,
and $M_i = \sup \left\{ \frac{1}{1+x} : x \in [x_{i-1}, x_i] \right\} = \frac{1}{1+x_{i-1}}$.

$$\begin{aligned} \Rightarrow U(P, f) - L(P, f) &= \sum_{i=1}^n \Delta x_i (M_i - m_i) = \frac{b}{n} \sum_{i=1}^n \left(\frac{1}{1 + (i-1)b/n} - \frac{1}{1 + ib/n} \right) \\ &= \frac{b}{n} \left(\frac{1}{1 + 0b/n} - \frac{1}{1 + nb/n} \right) = \frac{b^2}{n(b+1)}. \end{aligned}$$

Pick n to be such that $\frac{b^2}{n(b+1)} < \epsilon$, the proposition is satisfied,
and the function is integrable.



Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Integrals are “global”:

total distance travelled, average temperature, total charge . . .

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Integrals are “global”:

total distance travelled, average temperature, total charge . . .

Notation: If $f: S \rightarrow \mathbb{R}$ and $[a, b] \subset S$, use the restriction, but write $\int_a^b f$ and $f \in \mathcal{R}[a, b]$.

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Integrals are “global”:

total distance travelled, average temperature, total charge . . .

Notation: If $f: S \rightarrow \mathbb{R}$ and $[a, b] \subset S$, use the restriction, but write $\int_a^b f$ and $f \in \mathcal{R}[a, b]$.

We define $\int_a^b f$ even if $a \not\leq b$:

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Integrals are “global”:

total distance travelled, average temperature, total charge . . .

Notation: If $f: S \rightarrow \mathbb{R}$ and $[a, b] \subset S$, use the restriction, but write $\int_a^b f$ and $f \in \mathcal{R}[a, b]$.

We define $\int_a^b f$ even if $a \not\leq b$:

If $b < a$ and $f \in \mathcal{R}[b, a]$, then define
$$\int_a^b f := - \int_b^a f.$$

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Integrals are “global”:

total distance travelled, average temperature, total charge . . .

Notation: If $f: S \rightarrow \mathbb{R}$ and $[a, b] \subset S$, use the restriction, but write $\int_a^b f$ and $f \in \mathcal{R}[a, b]$.

We define $\int_a^b f$ even if $a \not\leq b$:

If $b < a$ and $f \in \mathcal{R}[b, a]$, then define $\int_a^b f := -\int_b^a f$. Also $\int_a^a f := 0$.

Remark: The integral adds up (integrates) local information—it sums $f(x) dx$ over all x .

Derivatives are “local”, only “at a point”:

velocity, acceleration, rate of change, slope . . .

Integrals are “global”:

total distance travelled, average temperature, total charge . . .

Notation: If $f: S \rightarrow \mathbb{R}$ and $[a, b] \subset S$, use the restriction, but write $\int_a^b f$ and $f \in \mathcal{R}[a, b]$.

We define $\int_a^b f$ even if $a \not\leq b$:

If $b < a$ and $f \in \mathcal{R}[b, a]$, then define $\int_a^b f := -\int_b^a f$. Also $\int_a^a f := 0$.

The x in dx is just a dummy variable, so, e.g., $\int_a^b f(s) ds := \int_a^b f(x) dx$.

Remark: Here's the Riemann approach to integrating $f: [a, b] \rightarrow \mathbb{R}$.

Remark: Here's the Riemann approach to integrating $f: [a, b] \rightarrow \mathbb{R}$.
Take a partition $P = \{x_0, x_1, \dots, x_n\}$

Remark: Here's the Riemann approach to integrating $f: [a, b] \rightarrow \mathbb{R}$.
Take a partition $P = \{x_0, x_1, \dots, x_n\}$
and a set of numbers (tags) $\{c_1, c_2, \dots, c_n\}$ with $c_k \in [x_{k-1}, x_k]$ for all k .

Remark: Here's the Riemann approach to integrating $f: [a, b] \rightarrow \mathbb{R}$.

Take a partition $P = \{x_0, x_1, \dots, x_n\}$

and a set of numbers (tags) $\{c_1, c_2, \dots, c_n\}$ with $c_k \in [x_{k-1}, x_k]$ for all k .

Then compute the *Riemann sum*:
$$\sum_{k=1}^n f(c_k) \Delta x_k.$$

Remark: Here's the Riemann approach to integrating $f: [a, b] \rightarrow \mathbb{R}$.

Take a partition $P = \{x_0, x_1, \dots, x_n\}$

and a set of numbers (tags) $\{c_1, c_2, \dots, c_n\}$ with $c_k \in [x_{k-1}, x_k]$ for all k .

Then compute the *Riemann sum*:
$$\sum_{k=1}^n f(c_k) \Delta x_k.$$

The normal definition of Riemann integrable is if a number I exists such that for every $\epsilon > 0$ there exists a $\delta > 0$ such that if P is a partition with $\Delta x_i < \delta$ for all i , then for any set of tags, the Riemann sum is within ϵ of I .

Remark: Here's the Riemann approach to integrating $f: [a, b] \rightarrow \mathbb{R}$.

Take a partition $P = \{x_0, x_1, \dots, x_n\}$

and a set of numbers (tags) $\{c_1, c_2, \dots, c_n\}$ with $c_k \in [x_{k-1}, x_k]$ for all k .

Then compute the *Riemann sum*:
$$\sum_{k=1}^n f(c_k) \Delta x_k.$$

The normal definition of Riemann integrable is if a number I exists such that for every $\epsilon > 0$ there exists a $\delta > 0$ such that if P is a partition with $\Delta x_i < \delta$ for all i , then for any set of tags, the Riemann sum is within ϵ of I .

Exercise: Prove that if f is Riemann integrable, then for any $\epsilon > 0$ there exists a Riemann sum that is within ϵ of the integral.