

# BA: 7.2

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

Write  $B_X(x, \delta) := B(x, \delta)$  or  $C_X(x, \delta) := C(x, \delta)$ , to emphasize the space.

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

Write  $B_X(x, \delta) := B(x, \delta)$  or  $C_X(x, \delta) := C(x, \delta)$ , to emphasize the space.

**Example:** Take  $\mathbb{R}$  with the standard metric:

$$B(x, \delta) = (x - \delta, x + \delta) \quad \text{and} \quad C(x, \delta) = [x - \delta, x + \delta].$$

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

Write  $B_X(x, \delta) := B(x, \delta)$  or  $C_X(x, \delta) := C(x, \delta)$ , to emphasize the space.

**Example:** Take  $\mathbb{R}$  with the standard metric:

$$B(x, \delta) = (x - \delta, x + \delta) \quad \text{and} \quad C(x, \delta) = [x - \delta, x + \delta].$$

**Example:** Consider  $[0, 1]$  as a subspace of  $\mathbb{R}$ . In  $[0, 1]$ ,

$$B(0, 1/2) = B_{[0,1]}(0, 1/2)$$

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

Write  $B_X(x, \delta) := B(x, \delta)$  or  $C_X(x, \delta) := C(x, \delta)$ , to emphasize the space.

**Example:** Take  $\mathbb{R}$  with the standard metric:

$$B(x, \delta) = (x - \delta, x + \delta) \quad \text{and} \quad C(x, \delta) = [x - \delta, x + \delta].$$

**Example:** Consider  $[0, 1]$  as a subspace of  $\mathbb{R}$ . In  $[0, 1]$ ,

$$B(0, 1/2) = B_{[0,1]}(0, 1/2) = \{y \in [0, 1] : |0 - y| < 1/2\}$$



## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

Write  $B_X(x, \delta) := B(x, \delta)$  or  $C_X(x, \delta) := C(x, \delta)$ , to emphasize the space.

**Example:** Take  $\mathbb{R}$  with the standard metric:

$$B(x, \delta) = (x - \delta, x + \delta) \quad \text{and} \quad C(x, \delta) = [x - \delta, x + \delta].$$

**Example:** Consider  $[0, 1]$  as a subspace of  $\mathbb{R}$ . In  $[0, 1]$ ,

$$B(0, 1/2) = B_{[0,1]}(0, 1/2) = \{y \in [0, 1] : |0 - y| < 1/2\} = [0, 1/2).$$

## Definition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ . Define the *open ball*, or *ball*, of radius  $\delta$  around  $x$  as

$$B(x, \delta) := \{y \in X : d(x, y) < \delta\}.$$

Define the *closed ball* as

$$C(x, \delta) := \{y \in X : d(x, y) \leq \delta\}.$$

Write  $B_X(x, \delta) := B(x, \delta)$  or  $C_X(x, \delta) := C(x, \delta)$ , to emphasize the space.

**Example:** Take  $\mathbb{R}$  with the standard metric:

$$B(x, \delta) = (x - \delta, x + \delta) \quad \text{and} \quad C(x, \delta) = [x - \delta, x + \delta].$$

**Example:** Consider  $[0, 1]$  as a subspace of  $\mathbb{R}$ . In  $[0, 1]$ ,

$$B(0, 1/2) = B_{[0,1]}(0, 1/2) = \{y \in [0, 1] : |0 - y| < 1/2\} = [0, 1/2).$$

Note the difference with  $B_{\mathbb{R}}(0, 1/2) = (-1/2, 1/2)$ .

## Definition

Let  $(X, d)$  be a metric space.

## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .

## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .

If  $x \in V$  and  $V$  is open, then  $V$  is an *open neighborhood* of  $x$  (or just *neighborhood*).

## Definition


Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .

If  $x \in V$  and  $V$  is open, then  $V$  is an *open neighborhood* of  $x$  (or just *neighborhood*).





## Definition

Let  $(X, d)$  be a metric space.


$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .

If  $x \in V$  and  $V$  is open, then  $V$  is an *open neighborhood* of  $x$  (or just *neighborhood*).

**Remark:**  $\delta$  depends on  $x$ .



## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .


$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .

If  $x \in V$  and  $V$  is open, then  $V$  is an *open neighborhood* of  $x$  (or just *neighborhood*).

**Remark:**  $\delta$  depends on  $x$ .

An open set  $V$  is a set that does not include its “boundary.”



## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.


When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .

If  $x \in V$  and  $V$  is open, then  $V$  is an *open neighborhood* of  $x$  (or just *neighborhood*).

**Remark:**  $\delta$  depends on  $x$ .

An open set  $V$  is a set that does not include its “boundary.”

If we are in  $V$  we are allowed to “wiggle” and we’ll stay in  $V$ .



## Definition

Let  $(X, d)$  be a metric space.

$V \subset X$  is *open* if  $\forall x \in V, \exists \delta > 0$  such that  $B(x, \delta) \subset V$ .

$E \subset X$  is *closed* if  $E^c = X \setminus E$  is open.

When  $X$  is not clear from context, we say  $V$  is *open in  $X$*  and  $E$  is *closed in  $X$* .


If  $x \in V$  and  $V$  is open, then  $V$  is an *open neighborhood* of  $x$  (or just *neighborhood*).

**Remark:**  $\delta$  depends on  $x$ .

An open set  $V$  is a set that does not include its “boundary.”

If we are in  $V$  we are allowed to “wiggle” and we’ll stay in  $V$ .

$E$  is closed if everything not in  $E$  is some distance away from  $E$ .



**Examples:**

**Examples:**

$(0, \infty) \subset \mathbb{R}$  is open:

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty)$ .



### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty)$ .

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

$\forall \delta > 0, B(0, \delta) = (-\delta, \delta)$  contains numbers not in  $[0, 1)$

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

$$\forall \delta > 0, B(0, \delta) = (-\delta, \delta) \text{ contains numbers not in } [0, 1) \Rightarrow [0, 1) \text{ is not open.}$$

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

$\forall \delta > 0, B(0, \delta) = (-\delta, \delta)$  contains numbers not in  $[0, 1)$   $\Rightarrow [0, 1)$  is not open.

$\forall \delta > 0, B(1, \delta) = (1 - \delta, 1 + \delta)$  contains numbers in  $[0, 1)$

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

$\forall \delta > 0, B(0, \delta) = (-\delta, \delta)$  contains numbers not in  $[0, 1)$   $\Rightarrow [0, 1)$  is not open.

$\forall \delta > 0, B(1, \delta) = (1 - \delta, 1 + \delta)$  contains numbers in  $[0, 1)$   $\Rightarrow [0, 1)^c$  is not open



## Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

$\forall \delta > 0, B(0, \delta) = (-\delta, \delta)$  contains numbers not in  $[0, 1)$   $\Rightarrow [0, 1)$  is not open.

$\forall \delta > 0, B(1, \delta) = (1 - \delta, 1 + \delta)$  contains numbers in  $[0, 1)$   $\Rightarrow [0, 1)^c$  is not open  
 $\Rightarrow [0, 1)$  is not closed.

### Examples:

$(0, \infty) \subset \mathbb{R}$  is open:

Given  $x \in (0, \infty)$ , let  $\delta := x$

$$\Rightarrow B(x, \delta) = (0, 2x) \subset (0, \infty).$$

$[0, \infty) \subset \mathbb{R}$  is closed:

Given  $x \in (-\infty, 0) = [0, \infty)^c$ , let  $\delta := -x$

$$\Rightarrow B(x, \delta) = (-2x, 0) \subset (-\infty, 0) = [0, \infty)^c.$$

$[0, 1) \subset \mathbb{R}$  is neither open nor closed:

$\forall \delta > 0, B(0, \delta) = (-\delta, \delta)$  contains numbers not in  $[0, 1) \Rightarrow [0, 1)$  is not open.

$\forall \delta > 0, B(1, \delta) = (1 - \delta, 1 + \delta)$  contains numbers in  $[0, 1) \Rightarrow [0, 1)^c$  is not open  
 $\Rightarrow [0, 1)$  is not closed.

In any metric space  $(X, d)$ , if  $x \in X$ , then  $\{x\}$  is closed (exercise).

## Proposition

*Let  $(X, d)$  be a metric space.*

## Proposition

Let  $(X, d)$  be a metric space.

- (i)  $\emptyset$  and  $X$  are open.

## Proposition

Let  $(X, d)$  be a metric space.

- (i)  $\emptyset$  and  $X$  are open.
- (ii) If  $V_1, V_2, \dots, V_k$  are open subsets of  $X$ , then

$$\bigcap_{j=1}^k V_j$$

is also open. That is, a finite intersection of open sets is open.

## Proposition

Let  $(X, d)$  be a metric space.

(i)  $\emptyset$  and  $X$  are open.

(ii) If  $V_1, V_2, \dots, V_k$  are open subsets of  $X$ , then

$$\bigcap_{j=1}^k V_j$$

is also open. That is, a finite intersection of open sets is open.

(iii) If  $\{V_\lambda\}_{\lambda \in I}$  is an arbitrary collection of open subsets of  $X$ , then

$$\bigcup_{\lambda \in I} V_\lambda$$

is also open. That is, a union of open sets is open.

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .



**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

$V_\lambda$  is open



**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

$V_\lambda$  is open  $\Rightarrow \exists \delta > 0$  such that  $B(x, \delta) \subset V_\lambda$ .

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

$V_\lambda$  is open  $\Rightarrow \exists \delta > 0$  such that  $B(x, \delta) \subset V_\lambda$ .

$\Rightarrow B(x, \delta) \subset \bigcup_{\lambda \in I} V_\lambda$

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

$V_\lambda$  is open  $\Rightarrow \exists \delta > 0$  such that  $B(x, \delta) \subset V_\lambda$ .

$\Rightarrow B(x, \delta) \subset \bigcup_{\lambda \in I} V_\lambda$

$\Rightarrow$  The union is open.



**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

$V_\lambda$  is open  $\Rightarrow \exists \delta > 0$  such that  $B(x, \delta) \subset V_\lambda$ .

$\Rightarrow B(x, \delta) \subset \bigcup_{\lambda \in I} V_\lambda$

$\Rightarrow$  The union is open.



**Example:** (ii) is not true for an arbitrary intersection:

**Proof:** (i):  $\emptyset$  and  $X$  are obviously open in  $X$ .

(ii): If  $x \in \bigcap_{j=1}^k V_j$ , then  $x \in V_j$  for all  $j$ .

$V_j$  are all open  $\Rightarrow \forall j \exists \delta_j > 0$  such that  $B(x, \delta_j) \subset V_j$ .

Take  $\delta := \min\{\delta_1, \delta_2, \dots, \delta_k\}$ .

Note  $\delta > 0$ .

$\Rightarrow B(x, \delta) \subset B(x, \delta_j) \subset V_j$  for every  $j$  and so  $B(x, \delta) \subset \bigcap_{j=1}^k V_j$ .

$\Rightarrow$  The intersection is open.

(iii): If  $x \in \bigcup_{\lambda \in I} V_\lambda$ , then  $x \in V_\lambda$  for some  $\lambda \in I$ .

$V_\lambda$  is open  $\Rightarrow \exists \delta > 0$  such that  $B(x, \delta) \subset V_\lambda$ .

$\Rightarrow B(x, \delta) \subset \bigcup_{\lambda \in I} V_\lambda$

$\Rightarrow$  The union is open.



**Example:** (ii) is not true for an arbitrary intersection:

$\bigcap_{n=1}^{\infty} (-1/n, 1/n) = \{0\}$ , which is not open.

## Proposition

*Let  $(X, d)$  be a metric space.*

## Proposition

Let  $(X, d)$  be a metric space.

- (i)  $\emptyset$  and  $X$  are closed.

## Proposition

Let  $(X, d)$  be a metric space.

- (i)  $\emptyset$  and  $X$  are closed.
- (ii) If  $\{E_\lambda\}_{\lambda \in I}$  is an arbitrary collection of closed subsets of  $X$ , then

$$\bigcap_{\lambda \in I} E_\lambda$$

is also closed. That is, an intersection of closed sets is closed.



## Proposition

Let  $(X, d)$  be a metric space.

(i)  $\emptyset$  and  $X$  are closed.

(ii) If  $\{E_\lambda\}_{\lambda \in I}$  is an arbitrary collection of closed subsets of  $X$ , then

$$\bigcap_{\lambda \in I} E_\lambda$$

is also closed. That is, an intersection of closed sets is closed.

(iii) If  $E_1, E_2, \dots, E_k$  are closed subsets of  $X$ , then

$$\bigcup_{j=1}^k E_j$$

is also closed. That is, a finite union of closed sets is closed.

## Proposition

Let  $(X, d)$  be a metric space.

(i)  $\emptyset$  and  $X$  are closed.

(ii) If  $\{E_\lambda\}_{\lambda \in I}$  is an arbitrary collection of closed subsets of  $X$ , then

$$\bigcap_{\lambda \in I} E_\lambda$$

is also closed. That is, an intersection of closed sets is closed.

(iii) If  $E_1, E_2, \dots, E_k$  are closed subsets of  $X$ , then

$$\bigcup_{j=1}^k E_j$$

is also closed. That is, a finite union of closed sets is closed.

**Proof:** Exercise.

## Proposition

*Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .*

## Proposition

*Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.*

## Proposition

*Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.*

**Proof:** Let  $y \in B(x, \delta)$ .

## Proposition

*Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.*

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ .

## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$d(x, z) \leq d(x, y) + d(y, z)$$



## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< d(x, y) + \alpha \end{aligned}$$

## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :


$$\begin{aligned}d(x, z) &\leq d(x, y) + d(y, z) \\&< d(x, y) + \alpha \\&= d(x, y) + \delta - d(x, y) = \delta.\end{aligned}$$

## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< d(x, y) + \alpha \\ &= d(x, y) + \delta - d(x, y) = \delta. \end{aligned}$$



## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.


**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$d(x, z) \leq d(x, y) + d(y, z)$$

$$< d(x, y) + \alpha$$

$$= d(x, y) + \delta - d(x, y) = \delta.$$

$$\Rightarrow B(y, \alpha) \subset B(x, \delta) \quad \Rightarrow B(x, \delta) \text{ is open.}$$



## Proposition


Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.

**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< d(x, y) + \alpha \\ &= d(x, y) + \delta - d(x, y) = \delta. \end{aligned}$$

$\Rightarrow B(y, \alpha) \subset B(x, \delta) \Rightarrow B(x, \delta)$  is open.

That  $C(x, \delta)$  is closed is an exercise.



□

## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.


**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< d(x, y) + \alpha \\ &= d(x, y) + \delta - d(x, y) = \delta. \end{aligned}$$

$\Rightarrow B(y, \alpha) \subset B(x, \delta) \Rightarrow B(x, \delta)$  is open.

That  $C(x, \delta)$  is closed is an exercise.

**Careful:**  $[0, 1/2)$  is an open ball in  $[0, 1]$ , and so  $[0, 1/2)$  is an open set in  $[0, 1]$ .



## Proposition

Let  $(X, d)$  be a metric space,  $x \in X$ , and  $\delta > 0$ .  
Then  $B(x, \delta)$  is open and  $C(x, \delta)$  is closed.


**Proof:** Let  $y \in B(x, \delta)$ . Let  $\alpha := \delta - d(x, y) > 0$ . Consider  $z \in B(y, \alpha)$ :

$$\begin{aligned} d(x, z) &\leq d(x, y) + d(y, z) \\ &< d(x, y) + \alpha \\ &= d(x, y) + \delta - d(x, y) = \delta. \end{aligned}$$

$\Rightarrow B(y, \alpha) \subset B(x, \delta) \Rightarrow B(x, \delta)$  is open.

That  $C(x, \delta)$  is closed is an exercise.

**Careful:**  $[0, 1/2)$  is an open ball in  $[0, 1]$ , and so  $[0, 1/2)$  is an open set in  $[0, 1]$ .  
 $[0, 1/2)$  is neither open nor closed in  $\mathbb{R}$ .



□

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ .*



## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ .*

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ .*

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .*

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .*

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .*

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .*

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open



## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .*

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

## Proposition

*Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .*

**Proof:** Exercise.

## Proposition

*Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .*

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .

## Proposition

Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .

**Proof:** Exercise.

## Proposition

Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .  $\exists \delta > 0$  such that  $B_X(x, \delta) \subset V$ .

## Proposition

Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .

**Proof:** Exercise.

## Proposition

Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .  $\exists \delta > 0$  such that  $B_X(x, \delta) \subset V$ .

Then  $B_Y(x, \delta)$

## Proposition

Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .

**Proof:** Exercise.

## Proposition

Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .  $\exists \delta > 0$  such that  $B_X(x, \delta) \subset V$ .

Then  $B_Y(x, \delta) = B_X(x, \delta) \cap Y$

## Proposition

Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .

**Proof:** Exercise.

## Proposition

Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .  $\exists \delta > 0$  such that  $B_X(x, \delta) \subset V$ .

Then  $B_Y(x, \delta) = B_X(x, \delta) \cap Y \subset V \cap Y$

## Proposition

Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .

**Proof:** Exercise.

## Proposition

Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .  $\exists \delta > 0$  such that  $B_X(x, \delta) \subset V$ .

Then  $B_Y(x, \delta) = B_X(x, \delta) \cap Y \subset V \cap Y = U$ .

## Proposition

Let  $a, b$  be two real numbers,  $a < b$ . Then  $(a, b)$ ,  $(a, \infty)$ , and  $(-\infty, b)$  are open in  $\mathbb{R}$ . Also  $[a, b]$ ,  $[a, \infty)$ , and  $(-\infty, b]$  are closed in  $\mathbb{R}$ .

**Proof:** Exercise.

## Proposition

Suppose  $(X, d)$  is a metric space, and  $Y \subset X$ . Then  $U \subset Y$  is open in  $Y$  (in the subspace topology)  $\Leftrightarrow \exists$  open set  $V \subset X$  (so open in  $X$ ), such that  $V \cap Y = U$ .

**Example:** Let  $X := \mathbb{R}$ ,  $Y := [0, 1]$ ,  $U := [0, 1/2)$ .

$U$  is an open set in  $Y$ , may take  $V := (-1/2, 1/2)$ .

**Proof of proposition:** Suppose  $V \subset X$  is open and  $V \cap Y = U$ .

Suppose  $x \in U = V \cap Y$ .  $\exists \delta > 0$  such that  $B_X(x, \delta) \subset V$ .

Then  $B_Y(x, \delta) = B_X(x, \delta) \cap Y \subset V \cap Y = U$ .

The opposite direction is an exercise.





## Proposition

*Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.*

## Proposition

*Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.*

*(i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .*

## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

- (i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .
- (ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

- (i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .
- (ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

**Proof:** (i): Suppose  $U \subset V$  is open in the subspace topology.

## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

- (i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .
- (ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

**Proof:** (i): Suppose  $U \subset V$  is open in the subspace topology.

By the previous proposition,  $\exists W \subset X$ , open in  $X$ , such that  $U = W \cap V$

## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

(i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .

(ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

**Proof:** (i): Suppose  $U \subset V$  is open in the subspace topology.

By the previous proposition,  $\exists W \subset X$ , open in  $X$ , such that  $U = W \cap V$

Intersection of open sets is open  $\Rightarrow U$  is open in  $X$ .

## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

- (i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .
- (ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

**Proof:** (i): Suppose  $U \subset V$  is open in the subspace topology.

By the previous proposition,  $\exists W \subset X$ , open in  $X$ , such that  $U = W \cap V$

Intersection of open sets is open  $\Rightarrow U$  is open in  $X$ .

Now suppose  $U$  is open in  $X$ .

## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

(i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .

(ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

**Proof:** (i): Suppose  $U \subset V$  is open in the subspace topology.

By the previous proposition,  $\exists W \subset X$ , open in  $X$ , such that  $U = W \cap V$

Intersection of open sets is open  $\Rightarrow U$  is open in  $X$ .

Now suppose  $U$  is open in  $X$ .

$\Rightarrow U = U \cap V \Rightarrow U$  is open in  $V$  (the proposition again)



## Proposition

Suppose  $(X, d)$  is a metric space,  $V \subset X$  is open, and  $E \subset X$  is closed.

(i)  $U \subset V$  is open in the subspace topology  $\Leftrightarrow U$  is open in  $X$ .

(ii)  $F \subset E$  is closed in the subspace topology  $\Leftrightarrow F$  is closed in  $X$ .

**Proof:** (i): Suppose  $U \subset V$  is open in the subspace topology.

By the previous proposition,  $\exists W \subset X$ , open in  $X$ , such that  $U = W \cap V$

Intersection of open sets is open  $\Rightarrow U$  is open in  $X$ .

Now suppose  $U$  is open in  $X$ .

$\Rightarrow U = U \cap V \Rightarrow U$  is open in  $V$  (the proposition again)

(ii): Exercise. □

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

A nonempty  $X$  is connected if  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and  $X_1$  and  $X_2$  are open, then either  $X_1 = \emptyset$  or  $X_2 = \emptyset$ .

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

A nonempty  $X$  is connected if  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and  $X_1$  and  $X_2$  are open, then either  $X_1 = \emptyset$  or  $X_2 = \emptyset$ .

To show  $X$  is disconnected, find nonempty disjoint open sets  $X_1$  and  $X_2$  whose union is  $X$ .

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

A nonempty  $X$  is connected if  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and  $X_1$  and  $X_2$  are open, then either  $X_1 = \emptyset$  or  $X_2 = \emptyset$ .

To show  $X$  is disconnected, find nonempty disjoint open sets  $X_1$  and  $X_2$  whose union is  $X$ .

**Example:**  $B(x, \delta)$  is not always connected.

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

A nonempty  $X$  is connected if  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and  $X_1$  and  $X_2$  are open, then either  $X_1 = \emptyset$  or  $X_2 = \emptyset$ .

To show  $X$  is disconnected, find nonempty disjoint open sets  $X_1$  and  $X_2$  whose union is  $X$ .

**Example:**  $B(x, \delta)$  is not always connected.

Consider  $\{a, b\}$  with the discrete metric.



## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

A nonempty  $X$  is connected if  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and  $X_1$  and  $X_2$  are open, then either  $X_1 = \emptyset$  or  $X_2 = \emptyset$ .

To show  $X$  is disconnected, find nonempty disjoint open sets  $X_1$  and  $X_2$  whose union is  $X$ .

**Example:**  $B(x, \delta)$  is not always connected.

Consider  $\{a, b\}$  with the discrete metric.

$\Rightarrow B(a, 2) = \{a, b\}$ , which is not connected:

## Definition

A nonempty metric space  $(X, d)$  is *connected* if the only subsets of  $X$  that are both open and closed (so-called *clopen* subsets) are  $\emptyset$  and  $X$  itself.

If a nonempty  $(X, d)$  is not connected we say it is *disconnected*.

A nonempty subset  $A \subset X$  is *connected* if  $A$  is connected with the subspace topology.

A nonempty  $X$  is connected if  $X = X_1 \cup X_2$  where  $X_1 \cap X_2 = \emptyset$  and  $X_1$  and  $X_2$  are open, then either  $X_1 = \emptyset$  or  $X_2 = \emptyset$ .

To show  $X$  is disconnected, find nonempty disjoint open sets  $X_1$  and  $X_2$  whose union is  $X$ .

**Example:**  $B(x, \delta)$  is not always connected.

Consider  $\{a, b\}$  with the discrete metric.

$\Rightarrow B(a, 2) = \{a, b\}$ , which is not connected:

$B(a, 1) = \{a\}$  and  $B(b, 1) = \{b\}$  are open and disjoint.

## Proposition

*Let  $(X, d)$  be a metric space.*

## Proposition


Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$

## Proposition


Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$



## Proposition


Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$


**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:

## Proposition


Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$


**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$



**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .



## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$



**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .

$\Leftarrow$ ) Suppose  $U_1$  and  $U_2$  exist.

## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$



**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .

$\Leftarrow$ ) Suppose  $U_1$  and  $U_2$  exist.  $\Rightarrow U_1 \cap S$  and  $U_2 \cap S$  are open in  $S$ .

## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$


**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .


$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .

$\Leftarrow$ ) Suppose  $U_1$  and  $U_2$  exist.  $\Rightarrow U_1 \cap S$  and  $U_2 \cap S$  are open in  $S$ .

$\Rightarrow S$  is disconnected. □

## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$


**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .


$\Leftarrow$ ) Suppose  $U_1$  and  $U_2$  exist.  $\Rightarrow U_1 \cap S$  and  $U_2 \cap S$  are open in  $S$ .

$\Rightarrow S$  is disconnected. □

**Example:**  $S \subset \mathbb{R}$  and  $\exists x, y, z$ , s.t.  $x < z < y$  with  $x, y \in S$  and  $z \notin S$ .

## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$


**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .

$\Leftarrow$ ) Suppose  $U_1$  and  $U_2$  exist.  $\Rightarrow U_1 \cap S$  and  $U_2 \cap S$  are open in  $S$ .


$\Rightarrow S$  is disconnected. □

**Example:**  $S \subset \mathbb{R}$  and  $\exists x, y, z$ , s.t.  $x < z < y$  with  $x, y \in S$  and  $z \notin S$ .

Claim:  $S$  is disconnected.

## Proposition

Let  $(X, d)$  be a metric space. A nonempty  $S \subset X$  is disconnected  $\Leftrightarrow \exists$  open sets  $U_1$  and  $U_2$  in  $X$ , such that  $U_1 \cap U_2 \cap S = \emptyset$ ,  $U_1 \cap S \neq \emptyset$ ,  $U_2 \cap S \neq \emptyset$ , and

$$S = (U_1 \cap S) \cup (U_2 \cap S).$$


**Proof:**  $\Rightarrow$ ) Suppose  $S$  is disconnected:  $\exists$  nonempty disjoint  $S_1$  and  $S_2$  that are open in  $S$  and  $S = S_1 \cup S_2$ .

$\Rightarrow \exists U_1$  and  $U_2$  that are open in  $X$  such that  $U_1 \cap S = S_1$  and  $U_2 \cap S = S_2$ .

$\Leftarrow$ ) Suppose  $U_1$  and  $U_2$  exist.  $\Rightarrow U_1 \cap S$  and  $U_2 \cap S$  are open in  $S$ .

$\Rightarrow S$  is disconnected. □

**Example:**  $S \subset \mathbb{R}$  and  $\exists x, y, z$ , s.t.  $x < z < y$  with  $x, y \in S$  and  $z \notin S$ .

Claim:  $S$  is disconnected.

Proof:  $((-\infty, z) \cap S) \cup ((z, \infty) \cap S) = S$

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.



## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

Suppose  $z \in \mathbb{R}$  is such that  $x < z < y$ .

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

Suppose  $z \in \mathbb{R}$  is such that  $x < z < y$ .

$(-\infty, z) \cap S \neq \emptyset$  and  $(z, \infty) \cap S \neq \emptyset$ .

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

Suppose  $z \in \mathbb{R}$  is such that  $x < z < y$ .

$(-\infty, z) \cap S \neq \emptyset$  and  $(z, \infty) \cap S \neq \emptyset$ .

The sets are disjoint.

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

Suppose  $z \in \mathbb{R}$  is such that  $x < z < y$ .

$(-\infty, z) \cap S \neq \emptyset$  and  $(z, \infty) \cap S \neq \emptyset$ .

The sets are disjoint.

As  $S$  is connected, their union can't be  $S$ .

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

Suppose  $z \in \mathbb{R}$  is such that  $x < z < y$ .

$(-\infty, z) \cap S \neq \emptyset$  and  $(z, \infty) \cap S \neq \emptyset$ .

The sets are disjoint.

As  $S$  is connected, their union can't be  $S$ .

$\Rightarrow z \in S$ .

## Proposition

*A nonempty set  $S \subset \mathbb{R}$  is connected  $\Leftrightarrow S$  is an interval or a single point.*

**Proof:** Suppose  $S$  is connected.

If  $S$  is a single point, then we are done.

Suppose  $x < y$  and  $x, y \in S$ .

Suppose  $z \in \mathbb{R}$  is such that  $x < z < y$ .

$(-\infty, z) \cap S \neq \emptyset$  and  $(z, \infty) \cap S \neq \emptyset$ .

The sets are disjoint.

As  $S$  is connected, their union can't be  $S$ .

$\Rightarrow z \in S$ .

$\Rightarrow S$  is an interval.



If  $S$  is a single point, it is connected,

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ .



If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

$z = \inf(U_2 \cap [x, y])$

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .

If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

$z = \inf(U_2 \cap [x, y]) \Rightarrow \exists w \in U_2 \cap [x, y]$  s.t.  $w \in [z, z + \delta) \subset B(z, \delta) \subset U_1$ .



If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .


If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

$z = \inf(U_2 \cap [x, y]) \Rightarrow \exists w \in U_2 \cap [x, y]$  s.t.  $w \in [z, z + \delta) \subset B(z, \delta) \subset U_1$ .



If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .


If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

$z = \inf(U_2 \cap [x, y]) \Rightarrow \exists w \in U_2 \cap [x, y]$  s.t.  $w \in [z, z + \delta) \subset B(z, \delta) \subset U_1$ .



$\Rightarrow w \in U_1 \cap U_2$

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .


If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

$z = \inf(U_2 \cap [x, y]) \Rightarrow \exists w \in U_2 \cap [x, y]$  s.t.  $w \in [z, z + \delta) \subset B(z, \delta) \subset U_1$ .



$\Rightarrow w \in U_1 \cap U_2 \Rightarrow (U_1 \cap S) \cap (U_2 \cap S) \neq \emptyset$

If  $S$  is a single point, it is connected, so suppose  $S$  is an interval.

Consider  $U_1$  and  $U_2$  open in  $\mathbb{R}$ , such that

$U_1 \cap S \neq \emptyset$  and  $U_2 \cap S \neq \emptyset$  and  $S = (U_1 \cap S) \cup (U_2 \cap S)$ .

Suppose  $x \in U_1 \cap S$  and  $y \in U_2 \cap S$ , WLOG  $x < y$ .

$S$  is an interval  $\Rightarrow [x, y] \subset S$ .

$U_2 \cap [x, y] \neq \emptyset$  let  $z := \inf(U_2 \cap [x, y])$ . WTS  $z \in U_1$ .

If  $z = x$ , then  $z \in U_1$ .


If  $z > x$ , then  $\forall \epsilon > 0$ ,  $B(z, \epsilon)$  has pts of  $[x, y] \setminus U_2$ , as  $z = \inf(U_2 \cap [x, y])$ .

So  $z \notin U_2$  as  $U_2$  is open.

$\Rightarrow z \in U_1$  ( $U_1$  and  $U_2$  cover  $[x, y]$ )

$U_1$  is open  $\Rightarrow B(z, \delta) \subset U_1$  for some  $\delta > 0$ .

$z = \inf(U_2 \cap [x, y]) \Rightarrow \exists w \in U_2 \cap [x, y]$  s.t.  $w \in [z, z + \delta) \subset B(z, \delta) \subset U_1$ .



$\Rightarrow w \in U_1 \cap U_2 \Rightarrow (U_1 \cap S) \cap (U_2 \cap S) \neq \emptyset \Rightarrow S$  is connected.



## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed,



## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets  $\Rightarrow \bar{A}$  is closed.

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets  $\Rightarrow \bar{A}$  is closed.

$\exists$  at least one closed set containing  $A$ , namely  $X$  itself,

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets  $\Rightarrow \bar{A}$  is closed.

$\exists$  at least one closed set containing  $A$ , namely  $X$  itself,  $\Rightarrow A \subset \bar{A}$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets  $\Rightarrow \bar{A}$  is closed.

$\exists$  at least one closed set containing  $A$ , namely  $X$  itself,  $\Rightarrow A \subset \bar{A}$ .

If  $A$  is closed, then  $A$  is a closed set that contains  $A$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets  $\Rightarrow \bar{A}$  is closed.

$\exists$  at least one closed set containing  $A$ , namely  $X$  itself,  $\Rightarrow A \subset \bar{A}$ .

If  $A$  is closed, then  $A$  is a closed set that contains  $A$ .

$\Rightarrow \bar{A} \subset A$



## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *closure* of  $A$  is the set

$$\bar{A} := \bigcap \{E \subset X : E \text{ is closed and } A \subset E\}.$$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $\bar{A}$  is closed, and  $A \subset \bar{A}$ .

Furthermore, if  $A$  is closed, then  $\bar{A} = A$ .

**Proof:**  $\bar{A}$  an intersection of closed sets  $\Rightarrow \bar{A}$  is closed.

$\exists$  at least one closed set containing  $A$ , namely  $X$  itself,  $\Rightarrow A \subset \bar{A}$ .

If  $A$  is closed, then  $A$  is a closed set that contains  $A$ .

$\Rightarrow \bar{A} \subset A \Rightarrow A = \bar{A}$ .



**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** Ambient space matters.

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** Ambient space matters.

If  $X = (0, \infty)$ , then the closure of  $(0, 1)$  in  $(0, \infty)$  is  $(0, 1]$ .



**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** Ambient space matters.

If  $X = (0, \infty)$ , then the closure of  $(0, 1)$  in  $(0, \infty)$  is  $(0, 1]$ .

$(0, 1]$  is closed in  $(0, \infty)$  (why?).

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** Ambient space matters.

If  $X = (0, \infty)$ , then the closure of  $(0, 1)$  in  $(0, \infty)$  is  $(0, 1]$ .

$(0, 1]$  is closed in  $(0, \infty)$  (why?).

Any closed set  $E$  s.t.  $(0, 1) \subset E$  must contain 1.

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** Ambient space matters.

If  $X = (0, \infty)$ , then the closure of  $(0, 1)$  in  $(0, \infty)$  is  $(0, 1]$ .

$(0, 1]$  is closed in  $(0, \infty)$  (why?).

Any closed set  $E$  s.t.  $(0, 1) \subset E$  must contain 1.

$\Rightarrow (0, 1] \subset E$

**Example:** The closure of  $(0, 1)$  in  $\mathbb{R}$  is  $[0, 1]$ .

Proof: If  $E$  is closed and  $(0, 1) \subset E$ , then  $0, 1 \in E$  (why?).  $\Rightarrow [0, 1] \subset E$ .

$[0, 1]$  is closed  $\Rightarrow \overline{(0, 1)} = [0, 1]$ .

**Example:** Ambient space matters.

If  $X = (0, \infty)$ , then the closure of  $(0, 1)$  in  $(0, \infty)$  is  $(0, 1]$ .

$(0, 1]$  is closed in  $(0, \infty)$  (why?).

Any closed set  $E$  s.t.  $(0, 1) \subset E$  must contain 1.

$\Rightarrow (0, 1] \subset E \Rightarrow \overline{(0, 1)} = (0, 1]$  in  $(0, \infty)$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ .*

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .



## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\Rightarrow \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

Now suppose  $\exists \delta > 0$ , such that  $B(x, \delta) \cap A = \emptyset$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

Now suppose  $\exists \delta > 0$ , such that  $B(x, \delta) \cap A = \emptyset$ .

$\implies A \subset B(x, \delta)^c$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

Now suppose  $\exists \delta > 0$ , such that  $B(x, \delta) \cap A = \emptyset$ .

$\implies A \subset B(x, \delta)^c$ .

$B(x, \delta)^c$  is closed,

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

Now suppose  $\exists \delta > 0$ , such that  $B(x, \delta) \cap A = \emptyset$ .

$\implies A \subset B(x, \delta)^c$ .

$B(x, \delta)^c$  is closed,  $x \notin B(x, \delta)^c$ ,



## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

Now suppose  $\exists \delta > 0$ , such that  $B(x, \delta) \cap A = \emptyset$ .

$\implies A \subset B(x, \delta)^c$ .

$B(x, \delta)^c$  is closed,  $x \notin B(x, \delta)^c$ ,  $\bar{A}$  is the intersection of closed sets containing  $A$

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .  $x \in \bar{A} \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$ .

**Proof:** We will show  $x \notin \bar{A} \iff \exists \delta > 0$  such that  $B(x, \delta) \cap A = \emptyset$ .

Suppose  $x \notin \bar{A}$ .

$\bar{A}$  is closed  $\implies \exists \delta > 0$  such that  $B(x, \delta) \subset \bar{A}^c$ .

As  $A \subset \bar{A}$ ,  $B(x, \delta) \subset \bar{A}^c \subset A^c \implies B(x, \delta) \cap A = \emptyset$ .

Now suppose  $\exists \delta > 0$ , such that  $B(x, \delta) \cap A = \emptyset$ .

$\implies A \subset B(x, \delta)^c$ .

$B(x, \delta)^c$  is closed,  $x \notin B(x, \delta)^c$ ,  $\bar{A}$  is the intersection of closed sets containing  $A$

$\implies x \notin \bar{A}$ .



## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *interior* of  $A$  is the set

$$A^\circ := \{x \in A : \text{there exists a } \delta > 0 \text{ such that } B(x, \delta) \subset A\}.$$

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *interior* of  $A$  is the set

$$A^\circ := \{x \in A : \text{there exists a } \delta > 0 \text{ such that } B(x, \delta) \subset A\}.$$

The *boundary* of  $A$  is the set

$$\partial A := \bar{A} \setminus A^\circ.$$

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *interior* of  $A$  is the set

$$A^\circ := \{x \in A : \text{there exists a } \delta > 0 \text{ such that } B(x, \delta) \subset A\}.$$

The *boundary* of  $A$  is the set

$$\partial A := \bar{A} \setminus A^\circ.$$

**Example:** Suppose  $A := (0, 1]$  and  $X := \mathbb{R}$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *interior* of  $A$  is the set

$$A^\circ := \{x \in A : \text{there exists a } \delta > 0 \text{ such that } B(x, \delta) \subset A\}.$$

The *boundary* of  $A$  is the set

$$\partial A := \bar{A} \setminus A^\circ.$$

**Example:** Suppose  $A := (0, 1]$  and  $X := \mathbb{R}$ .

$$\Rightarrow \bar{A} = [0, 1], A^\circ = (0, 1), \text{ and } \partial A = \{0, 1\}.$$

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *interior* of  $A$  is the set

$$A^\circ := \{x \in A : \text{there exists a } \delta > 0 \text{ such that } B(x, \delta) \subset A\}.$$

The *boundary* of  $A$  is the set

$$\partial A := \bar{A} \setminus A^\circ.$$

**Example:** Suppose  $A := (0, 1]$  and  $X := \mathbb{R}$ .

$\Rightarrow \bar{A} = [0, 1], A^\circ = (0, 1), \text{ and } \partial A = \{0, 1\}.$

**Example:** Consider  $X := \{a, b\}$  with the discrete metric, and  $A := \{a\}$ .

## Definition

Let  $(X, d)$  be a metric space and  $A \subset X$ . The *interior* of  $A$  is the set

$$A^\circ := \{x \in A : \text{there exists a } \delta > 0 \text{ such that } B(x, \delta) \subset A\}.$$

The *boundary* of  $A$  is the set

$$\partial A := \bar{A} \setminus A^\circ.$$

**Example:** Suppose  $A := (0, 1]$  and  $X := \mathbb{R}$ .

$$\Rightarrow \bar{A} = [0, 1], A^\circ = (0, 1), \text{ and } \partial A = \{0, 1\}.$$

**Example:** Consider  $X := \{a, b\}$  with the discrete metric, and  $A := \{a\}$ .

$$\Rightarrow \bar{A} = A^\circ = A \text{ and } \partial A = \emptyset.$$



## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ .*

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

**Proof:** Given  $x \in A^\circ$ ,  $\exists \delta > 0$  such that  $B(x, \delta) \subset A$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

**Proof:** Given  $x \in A^\circ$ ,  $\exists \delta > 0$  such that  $B(x, \delta) \subset A$ .

If  $z \in B(x, \delta)$ ,  $\exists \epsilon > 0$  such that  $B(z, \epsilon) \subset B(x, \delta) \subset A$ .

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

**Proof:** Given  $x \in A^\circ$ ,  $\exists \delta > 0$  such that  $B(x, \delta) \subset A$ .

If  $z \in B(x, \delta)$ ,  $\exists \epsilon > 0$  such that  $B(z, \epsilon) \subset B(x, \delta) \subset A$ .

$\Rightarrow z \in A^\circ$

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

**Proof:** Given  $x \in A^\circ$ ,  $\exists \delta > 0$  such that  $B(x, \delta) \subset A$ .

If  $z \in B(x, \delta)$ ,  $\exists \epsilon > 0$  such that  $B(z, \epsilon) \subset B(x, \delta) \subset A$ .

$\Rightarrow z \in A^\circ \Rightarrow B(x, \delta) \subset A^\circ$

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

**Proof:** Given  $x \in A^\circ$ ,  $\exists \delta > 0$  such that  $B(x, \delta) \subset A$ .

If  $z \in B(x, \delta)$ ,  $\exists \epsilon > 0$  such that  $B(z, \epsilon) \subset B(x, \delta) \subset A$ .

$\Rightarrow z \in A^\circ \Rightarrow B(x, \delta) \subset A^\circ \Rightarrow A^\circ$  is open.

## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $A^\circ$  is open and  $\partial A$  is closed.*

**Proof:** Given  $x \in A^\circ$ ,  $\exists \delta > 0$  such that  $B(x, \delta) \subset A$ .

If  $z \in B(x, \delta)$ ,  $\exists \epsilon > 0$  such that  $B(z, \epsilon) \subset B(x, \delta) \subset A$ .

$\Rightarrow z \in A^\circ \Rightarrow B(x, \delta) \subset A^\circ \Rightarrow A^\circ$  is open.

$A^\circ$  is open  $\Rightarrow \partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$  is closed.





## Proposition

*Let  $(X, d)$  be a metric space and  $A \subset X$ .*

## Proposition


*Let  $(X, d)$  be a metric space and  $A \subset X$ .*

*$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .*

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .




## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and




## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.




## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .



## Proposition


Let  $(X, d)$  be a metric space and  $A \subset X$ .

$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$



## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .


$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$

as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .





## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .


$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$   
as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .

Now suppose  $x \notin \partial A$



## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .

$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .


**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$

as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .

Now suppose  $x \notin \partial A \implies x \notin \bar{A}$  or  $x \in A^\circ$ .



## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .


$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$

as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .



Now suppose  $x \notin \partial A \implies x \notin \bar{A}$  or  $x \in A^\circ$ .

If  $x \notin \bar{A}$ , then  $B(x, \delta) \subset \bar{A}^c$  for some  $\delta > 0$  as  $\bar{A}$  is closed.

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .


$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$

as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .



Now suppose  $x \notin \partial A \implies x \notin \bar{A}$  or  $x \in A^\circ$ .

If  $x \notin \bar{A}$ , then  $B(x, \delta) \subset \bar{A}^c$  for some  $\delta > 0$  as  $\bar{A}$  is closed.

$\implies B(x, \delta) \cap A = \emptyset$  as  $\bar{A}^c \subset A^c$ .

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .


$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset$  and  $B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$

as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .



Now suppose  $x \notin \partial A \implies x \notin \bar{A}$  or  $x \in A^\circ$ .

If  $x \notin \bar{A}$ , then  $B(x, \delta) \subset \bar{A}^c$  for some  $\delta > 0$  as  $\bar{A}$  is closed.

$\implies B(x, \delta) \cap A = \emptyset$  as  $\bar{A}^c \subset A^c$ .

If  $x \in A^\circ$ , then  $B(x, \delta) \subset A$  for some  $\delta > 0$ , so  $B(x, \delta) \cap A^c = \emptyset$ .

□

## Proposition

Let  $(X, d)$  be a metric space and  $A \subset X$ .


$x \in \partial A \iff \forall \delta > 0, B(x, \delta) \cap A \neq \emptyset \text{ and } B(x, \delta) \cap A^c \neq \emptyset$ .

**Proof:** Suppose  $x \in \partial A = \bar{A} \setminus A^\circ$  and  $\delta > 0$  is arbitrary.

$x \in \bar{A} \implies B(x, \delta) \cap A \neq \emptyset$ .

If  $B(x, \delta) \cap A^c = \emptyset$ , then  $x \in A^\circ$

as  $x \notin A^\circ \implies B(x, \delta) \cap A^c \neq \emptyset$ .



Now suppose  $x \notin \partial A \implies x \notin \bar{A}$  or  $x \in A^\circ$ .

If  $x \notin \bar{A}$ , then  $B(x, \delta) \subset \bar{A}^c$  for some  $\delta > 0$  as  $\bar{A}$  is closed.

$\implies B(x, \delta) \cap A = \emptyset$  as  $\bar{A}^c \subset A^c$ .

If  $x \in A^\circ$ , then  $B(x, \delta) \subset A$  for some  $\delta > 0$ , so  $B(x, \delta) \cap A^c = \emptyset$ . □

## Corollary

Let  $(X, d)$  be a metric space and  $A \subset X$ . Then  $\partial A = \bar{A} \cap \bar{A}^c$ .

**Exercise:** Prove that in a metric space,

**Exercise:** Prove that in a metric space,

a)  $E$  is closed if and only if  $\partial E \subset E$ .



**Exercise:** Prove that in a metric space,

- a)  $E$  is closed if and only if  $\partial E \subset E$ .
- b)  $U$  is open if and only if  $\partial U \cap U = \emptyset$ .

**Exercise:** Prove that in a metric space,

- a)  $E$  is closed if and only if  $\partial E \subset E$ .
- b)  $U$  is open if and only if  $\partial U \cap U = \emptyset$ .

**Exercise:** Prove that in a metric space,

**Exercise:** Prove that in a metric space,

- a)  $E$  is closed if and only if  $\partial E \subset E$ .
- b)  $U$  is open if and only if  $\partial U \cap U = \emptyset$ .

**Exercise:** Prove that in a metric space,

- a)  $A$  is open if and only if  $A^\circ = A$ .

**Exercise:** Prove that in a metric space,

- a)  $E$  is closed if and only if  $\partial E \subset E$ .
- b)  $U$  is open if and only if  $\partial U \cap U = \emptyset$ .

**Exercise:** Prove that in a metric space,

- a)  $A$  is open if and only if  $A^\circ = A$ .
- b)  $U \subset A^\circ$  for every open set  $U$  such that  $U \subset A$ .

**Exercise:** Prove that in a metric space,

- a)  $E$  is closed if and only if  $\partial E \subset E$ .
- b)  $U$  is open if and only if  $\partial U \cap U = \emptyset$ .

**Exercise:** Prove that in a metric space,

- a)  $A$  is open if and only if  $A^\circ = A$ .
- b)  $U \subset A^\circ$  for every open set  $U$  such that  $U \subset A$ .

**Exercise:** Let  $(X, d)$  be a metric space and  $A \subset X$ . Show that  $A^\circ = \bigcup \{V : V \text{ is open and } V \subset A\}$ .