

Cultivating Complex Analysis: Cauchy's formula in a disc (3.2.4)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We prove a version in a disc.

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We prove a version in a disc.

Theorem (Cauchy integral formula in a disc)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $\overline{\Delta_r(p)} \subset U$. Then for all $z \in \Delta_r(p)$,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We prove a version in a disc.

Theorem (Cauchy integral formula in a disc)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $\overline{\Delta_r(p)} \subset U$. Then for all $z \in \Delta_r(p)$,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We prove a version in a disc.

Theorem (Cauchy integral formula in a disc)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $\overline{\Delta_r(p)} \subset U$. Then for all $z \in \Delta_r(p)$,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

Values inside (large set)

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We prove a version in a disc.

Theorem (Cauchy integral formula in a disc)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $\overline{\Delta_r(p)} \subset U$. Then for all $z \in \Delta_r(p)$,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

Values inside (large set)

are given in terms of values on the boundary (small set).

We get to perhaps the most fundamental theorem in complex analysis:
The Cauchy integral formula.

We prove a version in a disc.

Theorem (Cauchy integral formula in a disc)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $\overline{\Delta_r(p)} \subset U$. Then for all $z \in \Delta_r(p)$,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

Values inside (large set)
are given in terms of values on the boundary (small set).

A quick (but hardly only) application is to compute integrals of expressions such as $\frac{\cos(z^2)}{z(z-1)}$ that blow up somewhere inside the cycle.

Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.

Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.

Let $\Delta_s(z)$ be a small disc where $\overline{\Delta_s(z)} \subset \Delta_r(p)$.


Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.

Let $\Delta_s(z)$ be a small disc where $\overline{\Delta_s(z)} \subset \Delta_r(p)$. Write $\alpha = \partial\Delta_s(z)$.

Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.

Let $\Delta_s(z)$ be a small disc where $\overline{\Delta_s(z)} \subset \Delta_r(p)$. Write $\alpha = \partial\Delta_s(z)$.

Connect α to γ via two straight lines
to give two closed paths c_1 and c_2




Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.

Let $\Delta_s(z)$ be a small disc where $\overline{\Delta_s(z)} \subset \Delta_r(p)$. Write $\alpha = \partial\Delta_s(z)$.

Connect α to γ via two straight lines
to give two closed paths c_1 and c_2

As chains, $c_1 + c_2 = \gamma - \alpha$.



Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.


Let $\Delta_s(z)$ be a small disc where $\overline{\Delta_s(z)} \subset \Delta_r(p)$. Write $\alpha = \partial\Delta_s(z)$.

Connect α to γ via two straight lines to give two closed paths c_1 and c_2

As chains, $c_1 + c_2 = \gamma - \alpha$.

Each c_j lies in a star-like domain

where $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$ is holomorphic.



Proof: Fix $z \in \Delta_r(p)$, and write $\gamma = \partial\Delta_r(p)$ oriented counterclockwise.

Let $\Delta_s(z)$ be a small disc where $\overline{\Delta_s(z)} \subset \Delta_r(p)$. Write $\alpha = \partial\Delta_s(z)$.

Connect α to γ via two straight lines to give two closed paths c_1 and c_2

As chains, $c_1 + c_2 = \gamma - \alpha$.


Each c_j lies in a star-like domain

where $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$ is holomorphic. So

$$\int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta =$$

$$\int_{c_1} \frac{f(\zeta)}{\zeta - z} d\zeta + \int_{c_2} \frac{f(\zeta)}{\zeta - z} d\zeta = 0 + 0.$$

(by Cauchy's theorem)



So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over γ (independent of s) equals the integral over α for all $s > 0$ small enough,

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over γ (independent of s) equals the integral over α for all $s > 0$ small enough, so we can take the limit as $s \rightarrow 0$.

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over γ (independent of s) equals the integral over α for all $s > 0$ small enough, so we can take the limit as $s \rightarrow 0$.

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over γ (independent of s) equals the integral over α for all $s > 0$ small enough, so we can take the limit as $s \rightarrow 0$.

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over γ (independent of s) equals the integral over α for all $s > 0$ small enough, so we can take the limit as $s \rightarrow 0$.

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize α as $\alpha(t) = z + se^{it}$.

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over γ (independent of s) equals the integral over α for all $s > 0$ small enough, so we can take the limit as $s \rightarrow 0$.

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt = f(z).$$

(by continuity of f at z) □

There is a very useful consequence left as an easy exercise:

There is a very useful consequence left as an easy exercise:

Exercise: Suppose f is holomorphic in an open neighborhood of $\overline{\Delta_r(p)}$. Show that f at p is the average of the values on $\partial\Delta_r(p)$. That is, show

$$f(p) = \frac{1}{2\pi} \int_0^{2\pi} f(p + re^{it}) dt.$$