

Cultivating Complex Analysis:  
Harmonic functions  
Mean-value property (7.2.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

$f: \mathbb{R} \rightarrow \mathbb{R}$  is harmonic if

$f: \mathbb{R} \rightarrow \mathbb{R}$  is harmonic if

$$\nabla^2 f = \frac{\partial^2}{\partial x^2} f = f'' = 0.$$

$f: \mathbb{R} \rightarrow \mathbb{R}$  is harmonic if

$$\nabla^2 f = \frac{\partial^2}{\partial x^2} f = f'' = 0.$$

So  $f(x) = Ax + B$  (affine linear).

$f: \mathbb{R} \rightarrow \mathbb{R}$  is harmonic if

$$\nabla^2 f = \frac{\partial^2}{\partial x^2} f = f'' = 0.$$

So  $f(x) = Ax + B$  (affine linear).

For any interval  $[a, b]$ ,  $a < b$ ,

$$f\left(\frac{a+b}{2}\right) = \frac{f(a) + f(b)}{2}. \quad (*)$$

$f: \mathbb{R} \rightarrow \mathbb{R}$  is harmonic if

$$\nabla^2 f = \frac{\partial^2}{\partial x^2} f = f'' = 0.$$

So  $f(x) = Ax + B$  (affine linear).

For any interval  $[a, b]$ ,  $a < b$ ,

$$f\left(\frac{a+b}{2}\right) = \frac{f(a) + f(b)}{2}. \quad (*)$$

“Mean-value property”:

**Exercise:** A continuous  $f: \mathbb{R} \rightarrow \mathbb{R}$  is harmonic (affine linear)  $\Leftrightarrow (*)$  holds for all  $[a, b]$ .

### Theorem (Mean-value property)

Suppose  $U \subset \mathbb{C}$  is open. A continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if for every  $p \in U$  there is an  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$  and

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta \quad \text{for all } r < R_p.$$

Moreover, if  $f$  is harmonic we may choose any  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$ .

### Theorem (Mean-value property)

Suppose  $U \subset \mathbb{C}$  is open. A continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if for every  $p \in U$  there is an  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$  and

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta \quad \text{for all } r < R_p.$$

Moreover, if  $f$  is harmonic we may choose any  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$ .

**Proof:**

$\Rightarrow$ )



### Theorem (Mean-value property)

Suppose  $U \subset \mathbb{C}$  is open. A continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if for every  $p \in U$  there is an  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$  and

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta \quad \text{for all } r < R_p.$$

Moreover, if  $f$  is harmonic we may choose any  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$ .

**Proof:**

$\Rightarrow$ ) Suppose  $f$  is harmonic and  $\overline{\Delta_r(p)} \subset U$ .

### Theorem (Mean-value property)

Suppose  $U \subset \mathbb{C}$  is open. A continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if for every  $p \in U$  there is an  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$  and

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta \quad \text{for all } r < R_p.$$

Moreover, if  $f$  is harmonic we may choose any  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$ .

#### Proof:

$\Rightarrow$ ) Suppose  $f$  is harmonic and  $\overline{\Delta_r(p)} \subset U$ . ( $R_p$  as large as wanted)

## Theorem (Mean-value property)

Suppose  $U \subset \mathbb{C}$  is open. A continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if for every  $p \in U$  there is an  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$  and

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta \quad \text{for all } r < R_p.$$

Moreover, if  $f$  is harmonic we may choose any  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$ .

### Proof:

$\Rightarrow$ ) Suppose  $f$  is harmonic and  $\overline{\Delta_r(p)} \subset U$ . ( $R_p$  as large as wanted)

Solve the Dirichlet problem in  $\Delta_r(p)$  (at  $p$ )

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{it}) dt = P[f|_{\partial\Delta_r(p)}](p)$$

## Theorem (Mean-value property)

Suppose  $U \subset \mathbb{C}$  is open. A continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if for every  $p \in U$  there is an  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$  and

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta \quad \text{for all } r < R_p.$$

Moreover, if  $f$  is harmonic we may choose any  $R_p > 0$  such that  $\Delta_{R_p}(p) \subset U$ .

### Proof:

$\Rightarrow$ ) Suppose  $f$  is harmonic and  $\overline{\Delta_r(p)} \subset U$ . ( $R_p$  as large as wanted)

Solve the Dirichlet problem in  $\Delta_r(p)$  (at  $p$ )

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{it}) dt = P[f|_{\partial\Delta_r(p)}](p) = f(p).$$

$\Leftarrow$ )

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ .

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ .      Let  $h = P[f|_{\partial\Delta_s(q)}]$ .



$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ .      Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous,

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ ,

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .


Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),





$\Leftrightarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .


Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),

$\varphi \leq C < \varphi(p)$  (for some  $C \in \mathbb{R}$ ) on an open subset of  $\partial\Delta_r(p)$



$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .


$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),

$\varphi \leq C < \varphi(p)$  (for some  $C \in \mathbb{R}$ ) on an open subset of  $\partial\Delta_r(p)$

So  $\frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi(p + re^{i\theta}) d\theta < \varphi(p)$ .



$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .


$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),

$\varphi \leq C < \varphi(p)$  (for some  $C \in \mathbb{R}$ ) on an open subset of  $\partial\Delta_r(p)$

So  $\frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi(p + re^{i\theta}) d\theta < \varphi(p)$ . A contradiction!



$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .


$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),

$\varphi \leq C < \varphi(p)$  (for some  $C \in \mathbb{R}$ ) on an open subset of  $\partial\Delta_r(p)$

So  $\frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi(p + re^{i\theta}) d\theta < \varphi(p)$ . A contradiction! So  $\varphi \leq 0$ .



$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.


Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .

Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),

$\varphi \leq C < \varphi(p)$  (for some  $C \in \mathbb{R}$ ) on an open subset of  $\partial\Delta_r(p)$

So  $\frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi(p + re^{i\theta}) d\theta < \varphi(p)$ . A contradiction! So  $\varphi \leq 0$ .

Similarly  $\varphi \geq 0$ .



$\Leftarrow$ ) Suppose  $f$  is continuous and satisfies the mean-value property.

Suppose  $\overline{\Delta_s(q)} \subset U$ . Let  $h = P[f|_{\partial\Delta_s(q)}]$ .

Let  $\varphi = f - h$ .  $\varphi$  is continuous, identically zero on  $\partial\Delta_s(q)$ , and satisfies the mean-value property on same circles as  $f$ .

Suppose (for contradiction)  $\varphi$  attains a maximum at  $p \in \Delta_s(q)$  and  $\varphi(p) > 0$ .

$X = \{z \in \Delta_s(q) : \varphi(z) = \varphi(p)\}$  is compact.

Assume  $p$  is the point of  $X$  closest to  $\partial\Delta_s(q)$ .


Suppose  $\partial\Delta_r(p) \subset \Delta_s(q)$  (and  $r < R_p$ ),

$\varphi \leq C < \varphi(p)$  (for some  $C \in \mathbb{R}$ ) on an open subset of  $\partial\Delta_r(p)$

So  $\frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi(p + re^{i\theta}) d\theta < \varphi(p)$ . A contradiction! So  $\varphi \leq 0$ .

Similarly  $\varphi \geq 0$ .

So  $f = h$  and  $f$  is harmonic.



□

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.



For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

**Proof:** Firstly,  $f$  is continuous.

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

**Proof:** Firstly,  $f$  is continuous.

If  $\overline{\Delta_r(p)} \subset U$ , then  $\{f_n\}$  converges uniformly on  $\partial\Delta_r(p)$ . So

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

**Proof:** Firstly,  $f$  is continuous.

If  $\overline{\Delta_r(p)} \subset U$ , then  $\{f_n\}$  converges uniformly on  $\partial\Delta_r(p)$ . So

$$f(p) = \lim_{n \rightarrow \infty} f_n(p)$$

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

**Proof:** Firstly,  $f$  is continuous.

If  $\overline{\Delta_r(p)} \subset U$ , then  $\{f_n\}$  converges uniformly on  $\partial\Delta_r(p)$ . So

$$f(p) = \lim_{n \rightarrow \infty} f_n(p) = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{-\pi}^{\pi} f_n(p + re^{i\theta}) d\theta$$

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

**Proof:** Firstly,  $f$  is continuous.

If  $\overline{\Delta_r(p)} \subset U$ , then  $\{f_n\}$  converges uniformly on  $\partial\Delta_r(p)$ . So

$$f(p) = \lim_{n \rightarrow \infty} f_n(p) = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{-\pi}^{\pi} f_n(p + re^{i\theta}) d\theta = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta.$$

For  $C^2$  functions, just because  $f_n \rightarrow f$  uniformly, doesn't mean that  $\nabla^2 f_n$  goes to  $\nabla^2 f$ .

But it does for harmonic functions.

### Theorem (Harnack's first)

*Let  $U \subset \mathbb{C}$  be open, and let  $f_n: U \rightarrow \mathbb{R}$  be a sequence of harmonic functions converging uniformly on compact subsets to  $f: U \rightarrow \mathbb{R}$ . Then  $f$  is harmonic.*

**Proof:** Firstly,  $f$  is continuous.

If  $\overline{\Delta_r(p)} \subset U$ , then  $\{f_n\}$  converges uniformly on  $\partial\Delta_r(p)$ . So

$$f(p) = \lim_{n \rightarrow \infty} f_n(p) = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{-\pi}^{\pi} f_n(p + re^{i\theta}) d\theta = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + re^{i\theta}) d\theta.$$

Done by mean-value property.



**Exercise:** Prove the maximum principle for harmonic functions directly from the mean-value property.



**Exercise:** Prove the maximum principle for harmonic functions directly from the mean-value property.

**Exercise:** Let  $U \subset \mathbb{C}$  be open. Prove that a continuous  $f: U \rightarrow \mathbb{R}$  is harmonic if and only if it satisfies the *disc mean-value property* for every  $\overline{\Delta_r(p)} \subset U$ :

$$f(p) = \frac{1}{\pi r^2} \int_{\overline{\Delta_r(p)}} f(z) dA.$$