

# Cultivating Complex Analysis: The logarithm (4.1.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Consider the primitive of  $z^n$ .

Consider the primitive of  $z^n$ .

If  $n \neq -1$ , the primitive is  $\frac{z^{n+1}}{n+1}$  (not defined at  $z = 0$  if  $n + 1 < 0$ )

Consider the primitive of  $z^n$ .

If  $n \neq -1$ , the primitive is  $\frac{z^{n+1}}{n+1}$  (not defined at  $z = 0$  if  $n + 1 < 0$ )

What about  $z^{-1} = 1/z$ ?

Consider the primitive of  $z^n$ .

If  $n \neq -1$ , the primitive is  $\frac{z^{n+1}}{n+1}$  (not defined at  $z = 0$  if  $n + 1 < 0$ )

What about  $z^{-1} = 1/z$ ?

Consider the *slit plane*

$$U = \mathbb{C} \setminus (-\infty, 0] = \mathbb{C} \setminus \{z \in \mathbb{C} : \operatorname{Re} z \leq 0, \operatorname{Im} z = 0\}.$$

Consider the primitive of  $z^n$ .

If  $n \neq -1$ , the primitive is  $\frac{z^{n+1}}{n+1}$  (not defined at  $z = 0$  if  $n + 1 < 0$ )

What about  $z^{-1} = 1/z$ ?

Consider the *slit plane*

$$U = \mathbb{C} \setminus (-\infty, 0] = \mathbb{C} \setminus \{z \in \mathbb{C} : \operatorname{Re} z \leq 0, \operatorname{Im} z = 0\}.$$

$U$  is star-like  $\Rightarrow$  holomorphic functions on  $U$  have a primitive on  $U$ , including  $1/z$ .

Consider the primitive of  $z^n$ .

If  $n \neq -1$ , the primitive is  $\frac{z^{n+1}}{n+1}$  (not defined at  $z = 0$  if  $n + 1 < 0$ )

What about  $z^{-1} = 1/z$ ?

Consider the *slit plane*

$$U = \mathbb{C} \setminus (-\infty, 0] = \mathbb{C} \setminus \{z \in \mathbb{C} : \operatorname{Re} z \leq 0, \operatorname{Im} z = 0\}.$$

$U$  is star-like  $\Rightarrow$  holomorphic functions on  $U$  have a primitive on  $U$ , including  $1/z$ .

Require this primitive to be 0 at  $z = 1$  to obtain the *principal branch* of the logarithm:

$$\operatorname{Log}: U \rightarrow \mathbb{C}$$

Consider the primitive of  $z^n$ .

If  $n \neq -1$ , the primitive is  $\frac{z^{n+1}}{n+1}$  (not defined at  $z = 0$  if  $n + 1 < 0$ )

What about  $z^{-1} = 1/z$ ?

Consider the *slit plane*

$$U = \mathbb{C} \setminus (-\infty, 0] = \mathbb{C} \setminus \{z \in \mathbb{C} : \operatorname{Re} z \leq 0, \operatorname{Im} z = 0\}.$$

$U$  is star-like  $\Rightarrow$  holomorphic functions on  $U$  have a primitive on  $U$ , including  $1/z$ .

Require this primitive to be 0 at  $z = 1$  to obtain the *principal branch* of the logarithm:

$$\operatorname{Log}: U \rightarrow \mathbb{C}$$

We want to show  $\operatorname{Log} z = \log|z| + i \operatorname{Arg} z$  (principal branch of the argument).



Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z) e^{L(z)} = L'(z) z$ .

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z) e^{L(z)} = L'(z) z$ .

Et voilà!

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z) e^{L(z)} = L'(z) z$ .

Et voilà!

Using a different branch of the argument gets another antiderivative.



Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z) e^{L(z)} = L'(z) z$ .

Et voilà!

Using a different branch of the argument gets another antiderivative.

Emboldened, we define

$$\log z \stackrel{\text{def}}{=} \log|z| + i \arg z.$$

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z) e^{L(z)} = L'(z) z$ .

Et voilà!

Using a different branch of the argument gets another antiderivative.

Emboldened, we define

$$\log z \stackrel{\text{def}}{=} \log|z| + i \arg z.$$

That sounds crazy:

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z)e^{L(z)} = L'(z)z$ .

Et voilà!

Using a different branch of the argument gets another antiderivative.

Emboldened, we define

$$\log z \stackrel{\text{def}}{=} \log|z| + i \arg z.$$

That sounds crazy:

1) The log on the right (the real log) is different than the log on the left, and

Set  $L(z) = \log|z| + i \operatorname{Arg} z$  (WTS that  $L = \operatorname{Log}$ ).

$L(1) = 0 = \operatorname{Log}(1)$ , good!

$$e^{L(z)} = e^{\log|z|} e^{i \operatorname{Arg} z} = |z| e^{i \operatorname{Arg} z} = z.$$

$L$  is the inverse of the exponential  $\Rightarrow L$  is holomorphic.

Differentiate  $z = e^{L(z)}$ :  $1 = L'(z) e^{L(z)} = L'(z) z$ .

Et voilà!

Using a different branch of the argument gets another antiderivative.



Emboldened, we define

$$\log z \stackrel{\text{def}}{=} \log|z| + i \arg z.$$



That sounds crazy:

- 1) The log on the right (the real log) is different than the log on the left, and
- 2)  $\arg$  has infinitely many values.

Here are the real and imaginary parts of  $\log z = \log|z| + i \arg z$ :





Here are the real and imaginary parts of  $\log z = \log|z| + i \arg z$ :



If we travel the unit circle in the  $z$ -plane, we travel the marked path on the graph.



Here are the real and imaginary parts of  $\log z = \log|z| + i \arg z$ :



If we travel the unit circle in the  $z$ -plane, we travel the marked path on the graph.

The real part is a nice function, it is the normal real  $\log$ :  $(0, \infty) \rightarrow \mathbb{R}$  applied to  $|z|$ .

Here are the real and imaginary parts of  $\log z = \log|z| + i \arg z$ :





If we travel the unit circle in the  $z$ -plane, we travel the marked path on the graph.

The real part is a nice function, it is the normal real  $\log$ :  $(0, \infty) \rightarrow \mathbb{R}$  applied to  $|z|$ .

The imaginary part has infinitely many values.



Here are the real and imaginary parts of  $\log z = \log|z| + i \arg z$ :



If we travel the unit circle in the  $z$ -plane, we travel the marked path on the graph.

The real part is a nice function, it is the normal real  $\log$ :  $(0, \infty) \rightarrow \mathbb{R}$  applied to  $|z|$ .

The imaginary part has infinitely many values.

Nevertheless, it is the correct definition. Much more useful than the principal branch.

How do we use log? To compute line integrals:

How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz$$

How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1$$

How do we use  $\log$ ? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1 = 2\pi i.$$

How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1 = 2\pi i.$$

That's nonsense! Let's make it better:

How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1 = 2\pi i.$$

That's nonsense! Let's make it better:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz \text{ ``="} \log 1 - \log 1 \text{ ``="} 2\pi i.$$

How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1 = 2\pi i.$$

That's nonsense! Let's make it better:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz \text{ ``="} \log 1 - \log 1 \text{ ``="} 2\pi i.$$

Maybe still not quite right.



How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1 = 2\pi i.$$

That's nonsense! Let's make it better:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz \text{ “=” } \log 1 - \log 1 \text{ “=” } 2\pi i.$$

Maybe still not quite right.

It works by following one “branch” of the logarithm along the path and then subtracting.

How do we use log? To compute line integrals:

Parametrize  $\partial\mathbb{D}$  starting and ending at  $z = 1$  and compute:

$$\int_{\partial\mathbb{D}} \frac{1}{z} dz = \log 1 - \log 1 = 2\pi i.$$



That's nonsense! Let's make it better:


$$\int_{\partial\mathbb{D}} \frac{1}{z} dz \text{ ``="} \log 1 - \log 1 \text{ ``="} 2\pi i.$$

Maybe still not quite right.


It works by following one "branch" of the logarithm along the path and then subtracting.



Let's see that graph again.







What we do:





What we do:

We start with the value  $\log 1 = 0$ .



What we do:

We start with the value  $\log 1 = 0$ .

Then we follow the graph around the circle until we end at  $\log 1 = 2\pi i$ .

A *branch* of the logarithm in  $U$  is an antiderivative of  $1/z$  in  $U$  that equals one value of  $\log z$  at every point.

A *branch* of the logarithm in  $U$  is an antiderivative of  $1/z$  in  $U$  that equals one value of  $\log z$  at every point.

We *follow a branch* along a path by taking a branch in a (small enough) neighborhood,



A *branch* of the logarithm in  $U$  is an antiderivative of  $1/z$  in  $U$  that equals one value of  $\log z$  at every point.


We *follow a branch* along a path by taking a branch in a (small enough) neighborhood, then change to another branch in another neighborhood (equal at some point).

A *branch* of the logarithm in  $U$  is an antiderivative of  $1/z$  in  $U$  that equals one value of  $\log z$  at every point.

We *follow a branch* along a path by taking a branch in a (small enough) neighborhood, then change to another branch in another neighborhood (equal at some point). Etc.


A *branch* of the logarithm in  $U$  is an antiderivative of  $1/z$  in  $U$  that equals one value of  $\log z$  at every point.

We *follow a branch* along a path by taking a branch in a (small enough) neighborhood, then change to another branch in another neighborhood (equal at some point). Etc.



A *branch* of the logarithm in  $U$  is an antiderivative of  $1/z$  in  $U$  that equals one value of  $\log z$  at every point.

We *follow a branch* along a path by taking a branch in a (small enough) neighborhood, then change to another branch in another neighborhood (equal at some point). Etc.



That's what we did in the computation above.