

Cultivating Complex Analysis:
Harmonic functions
Real and imaginary parts of holomorphic functions (7.1.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Let $U \subset \mathbb{C}$ be open. A twice continuously differentiable $f: U \rightarrow \mathbb{R}$ is *harmonic* if

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0 \quad \text{on } U.$$

Definition

Let $U \subset \mathbb{C}$ be open. A twice continuously differentiable $f: U \rightarrow \mathbb{R}$ is *harmonic* if

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0 \quad \text{on } U.$$

∇^2 (sometimes written Δ) is the *Laplacian*.

Definition

Let $U \subset \mathbb{C}$ be open. A twice continuously differentiable $f: U \rightarrow \mathbb{R}$ is *harmonic* if

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0 \quad \text{on } U.$$

∇^2 (sometimes written Δ) is the *Laplacian*.

Convenient to write using Wirtinger operators:

$$4 \frac{\partial^2}{\partial \bar{z} \partial z} f = 4 \left[\frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) \right] \left[\frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \right] f$$

Definition

Let $U \subset \mathbb{C}$ be open. A twice continuously differentiable $f: U \rightarrow \mathbb{R}$ is *harmonic* if

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0 \quad \text{on } U.$$

∇^2 (sometimes written Δ) is the *Laplacian*.

Convenient to write using Wirtinger operators:

$$4 \frac{\partial^2}{\partial \bar{z} \partial z} f = 4 \left[\frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) \right] \left[\frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \right] f = \left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right] f = \nabla^2 f.$$

Definition

Let $U \subset \mathbb{C}$ be open. A twice continuously differentiable $f: U \rightarrow \mathbb{R}$ is *harmonic* if

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0 \quad \text{on } U.$$

∇^2 (sometimes written Δ) is the *Laplacian*.

Convenient to write using Wirtinger operators:

$$4 \frac{\partial^2}{\partial \bar{z} \partial z} f = 4 \left[\frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) \right] \left[\frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \right] f = \left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right] f = \nabla^2 f.$$

$$f \text{ is harmonic} \quad \Leftrightarrow \quad \frac{\partial f}{\partial \bar{z}} \text{ is holomorphic.}$$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic.

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z)$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z) \Rightarrow c$ is C^2 and $\frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z}$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z) \Rightarrow c$ is C^2 and $\frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0$.

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z) \Rightarrow c$ is C^2 and $\frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0$.

Let $h = \bar{c}$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z) \Rightarrow c$ is C^2 and $\frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0$.

Let $h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z) \Rightarrow c$ is C^2 and $\frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0$.

Let $h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h$ is holomorphic.

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

$$\text{Let } \varphi = g + h.$$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

Let $c(z) = f(z) - g(z) \Rightarrow c$ is C^2 and $\frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0$.

Let $h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h$ is holomorphic.

$\Rightarrow f(z) = g(z) + \overline{h(z)}$, g and h holomorphic.

Let $\varphi = g + h$. As f is real-valued,

$$f(z) = \operatorname{Re} f(z)$$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

Let $\varphi = g + h$. As f is real-valued,

$$f(z) = \operatorname{Re} f(z) = \frac{g(z) + \overline{h(z)} + \overline{g(z)} + h(z)}{2} = \frac{g(z) + h(z) + \overline{g(z) + h(z)}}{2}$$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

Let $\varphi = g + h$. As f is real-valued,

$$f(z) = \operatorname{Re} f(z) = \frac{g(z) + \overline{h(z)} + \overline{g(z)} + h(z)}{2} = \frac{g(z) + h(z) + \overline{g(z) + h(z)}}{2} = \operatorname{Re} \varphi(z).$$

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

Let $\varphi = g + h$. As f is real-valued,

$$f(z) = \operatorname{Re} f(z) = \frac{g(z) + \overline{h(z)} + \overline{g(z)} + h(z)}{2} = \frac{g(z) + h(z) + \overline{g(z) + h(z)}}{2} = \operatorname{Re} \varphi(z).$$

Note: This works locally, or in a simply connected U where we can find the primitive g .

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial z}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial g}{\partial z} = \frac{\partial f}{\partial z} - \frac{\partial f}{\partial z} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

Let $\varphi = g + h$. As f is real-valued,

$$f(z) = \operatorname{Re} f(z) = \frac{g(z) + \overline{h(z)} + \overline{g(z)} + h(z)}{2} = \frac{g(z) + h(z) + \overline{g(z) + h(z)}}{2} = \operatorname{Re} \varphi(z).$$

Note: This works locally, or in a simply connected U where we can find the primitive g .

Proposition

Let $U \subset \mathbb{C}$ be a simply connected domain and $f: U \rightarrow \mathbb{R}$ a harmonic function. Then there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f = \operatorname{Re} \varphi$.

Suppose $f: U \rightarrow \mathbb{R}$ is harmonic. Find (locally) a primitive g of the holomorphic $\frac{\partial f}{\partial \bar{z}}$.

$$\text{Let } c(z) = f(z) - g(z) \Rightarrow c \text{ is } C^2 \text{ and } \frac{\partial c}{\partial \bar{z}} = \frac{\partial f}{\partial \bar{z}} - \frac{\partial g}{\partial \bar{z}} = \frac{\partial f}{\partial \bar{z}} - \frac{\partial f}{\partial \bar{z}} = 0.$$

$$\text{Let } h = \bar{c} \Rightarrow \frac{\partial h}{\partial \bar{z}} = \frac{\partial \bar{c}}{\partial \bar{z}} = \overline{\frac{\partial c}{\partial z}} = 0 \Rightarrow h \text{ is holomorphic.}$$

$$\Rightarrow f(z) = g(z) + \overline{h(z)}, \quad g \text{ and } h \text{ holomorphic.}$$

Let $\varphi = g + h$. As f is real-valued,

$$f(z) = \operatorname{Re} f(z) = \frac{g(z) + \overline{h(z)} + \overline{g(z)} + h(z)}{2} = \frac{g(z) + h(z) + \overline{g(z) + h(z)}}{2} = \operatorname{Re} \varphi(z).$$

Note: This works locally, or in a simply connected U where we can find the primitive g .

Proposition

Let $U \subset \mathbb{C}$ be a simply connected domain and $f: U \rightarrow \mathbb{R}$ a harmonic function. Then there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f = \operatorname{Re} \varphi$.

Similarly f is the imaginary part of some holomorphic function.

Conversely suppose $f(z) = \operatorname{Re} \varphi(z) = \frac{1}{2}(\varphi(z) + \overline{\varphi(z)})$ (φ holomorphic).

Conversely suppose $f(z) = \operatorname{Re} \varphi(z) = \frac{1}{2}(\varphi(z) + \overline{\varphi(z)})$ (φ holomorphic).

$$\nabla^2 = 4 \frac{\partial^2}{\partial \bar{z} \partial z} = 4 \frac{\partial^2}{\partial z \partial \bar{z}}$$

Conversely suppose $f(z) = \operatorname{Re} \varphi(z) = \frac{1}{2}(\varphi(z) + \overline{\varphi(z)})$ (φ holomorphic).

$$\nabla^2 = 4 \frac{\partial^2}{\partial \bar{z} \partial z} = 4 \frac{\partial^2}{\partial z \partial \bar{z}} \Rightarrow \nabla^2 f = 4 \frac{\partial^2}{\partial \bar{z} \partial z} \left(\frac{1}{2} (\varphi(z) + \overline{\varphi(z)}) \right) = 2 \left(\frac{\partial}{\partial z} \left(\frac{\partial \varphi}{\partial \bar{z}} \right) + \frac{\partial}{\partial \bar{z}} \left(\frac{\partial \bar{\varphi}}{\partial z} \right) \right) = 0.$$

Conversely suppose $f(z) = \operatorname{Re} \varphi(z) = \frac{1}{2}(\varphi(z) + \overline{\varphi(z)})$ (φ holomorphic).

$$\nabla^2 = 4 \frac{\partial^2}{\partial \bar{z} \partial z} = 4 \frac{\partial^2}{\partial z \partial \bar{z}} \Rightarrow \nabla^2 f = 4 \frac{\partial^2}{\partial \bar{z} \partial z} \left(\frac{1}{2}(\varphi(z) + \overline{\varphi(z)}) \right) = 2 \left(\frac{\partial}{\partial z} \left(\frac{\partial \varphi}{\partial \bar{z}} \right) + \frac{\partial}{\partial \bar{z}} \left(\frac{\partial \bar{\varphi}}{\partial z} \right) \right) = 0.$$

We proved:

Proposition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ a function.

- (i) *The function f is harmonic if and only if for every $p \in U$ there exists an open neighborhood V of p and a holomorphic $\varphi: V \rightarrow \mathbb{C}$ such that $f = \operatorname{Re} \varphi$ on V .*

Conversely suppose $f(z) = \operatorname{Re} \varphi(z) = \frac{1}{2}(\varphi(z) + \overline{\varphi(z)})$ (φ holomorphic).

$$\nabla^2 = 4 \frac{\partial^2}{\partial \bar{z} \partial z} = 4 \frac{\partial^2}{\partial z \partial \bar{z}} \Rightarrow \nabla^2 f = 4 \frac{\partial^2}{\partial \bar{z} \partial z} \left(\frac{1}{2}(\varphi(z) + \overline{\varphi(z)}) \right) = 2 \left(\frac{\partial}{\partial z} \left(\frac{\partial \varphi}{\partial \bar{z}} \right) + \frac{\partial}{\partial \bar{z}} \left(\frac{\partial \bar{\varphi}}{\partial z} \right) \right) = 0.$$

We proved:

Proposition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ a function.

- (i) The function f is harmonic if and only if for every $p \in U$ there exists an open neighborhood V of p and a holomorphic $\varphi: V \rightarrow \mathbb{C}$ such that $f = \operatorname{Re} \varphi$ on V .
- (ii) The function f is harmonic if and only if for every $p \in U$, there exists a power series expansion

$$f(z) = c_0 + \sum_{n=1}^{\infty} c_n (z-p)^n + \bar{c}_n (\overline{z-p})^n$$

converging uniformly absolutely on every closed disc $\overline{\Delta_r(p)} \subset U$.

A quick corollary:

Proposition

If $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{R}$ is harmonic, then f is infinitely (real) differentiable.

Proof: Holomorphic functions are infinitely differentiable.

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ harmonic. If $g: U \rightarrow \mathbb{R}$ is harmonic and $f + ig$ is holomorphic, then g is called the *harmonic conjugate* of f .

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ harmonic. If $g: U \rightarrow \mathbb{R}$ is harmonic and $f + ig$ is holomorphic, then g is called the *harmonic conjugate* of f .

Every harmonic f on a simply connected domain has a harmonic conjugate.

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ harmonic. If $g: U \rightarrow \mathbb{R}$ is harmonic and $f + ig$ is holomorphic, then g is called the *harmonic conjugate* of f .

Every harmonic f on a simply connected domain has a harmonic conjugate.

On $\mathbb{C} \setminus \{0\}$, $z \mapsto \log|z|$ is harmonic, but fails to have a harmonic conjugate.

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ harmonic. If $g: U \rightarrow \mathbb{R}$ is harmonic and $f + ig$ is holomorphic, then g is called the *harmonic conjugate* of f .

Every harmonic f on a simply connected domain has a harmonic conjugate.

On $\mathbb{C} \setminus \{0\}$, $z \mapsto \log|z|$ is harmonic, but fails to have a harmonic conjugate.

If it did have a harmonic conjugate then \log would have a branch in $\mathbb{C} \setminus \{0\}$.

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ harmonic. If $g: U \rightarrow \mathbb{R}$ is harmonic and $f + ig$ is holomorphic, then g is called the *harmonic conjugate* of f .

Every harmonic f on a simply connected domain has a harmonic conjugate.

On $\mathbb{C} \setminus \{0\}$, $z \mapsto \log|z|$ is harmonic, but fails to have a harmonic conjugate.

If it did have a harmonic conjugate then \log would have a branch in $\mathbb{C} \setminus \{0\}$.

Which follows from:

Proposition

If $U \subset \mathbb{C}$ is a domain $f: U \rightarrow \mathbb{R}$ is harmonic and g_1 and g_2 are two harmonic conjugates of f , then $g_1 = g_2 + C$ for some $C \in \mathbb{R}$.

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{R}$ harmonic. If $g: U \rightarrow \mathbb{R}$ is harmonic and $f + ig$ is holomorphic, then g is called the *harmonic conjugate* of f .

Every harmonic f on a simply connected domain has a harmonic conjugate.

On $\mathbb{C} \setminus \{0\}$, $z \mapsto \log|z|$ is harmonic, but fails to have a harmonic conjugate.

If it did have a harmonic conjugate then \log would have a branch in $\mathbb{C} \setminus \{0\}$.

Which follows from:

Proposition

If $U \subset \mathbb{C}$ is a domain $f: U \rightarrow \mathbb{R}$ is harmonic and g_1 and g_2 are two harmonic conjugates of f , then $g_1 = g_2 + C$ for some $C \in \mathbb{R}$.

Proof: $\frac{(f + ig_1) - (f + ig_2)}{i} = g_1 - g_2$ is holomorphic, real-valued \Rightarrow constant.

The real and imaginary parts of a holomorphic function are harmonic.

The real and imaginary parts of a holomorphic function are harmonic.

The modulus $|f(z)|$ is not.

The real and imaginary parts of a holomorphic function are harmonic.

The modulus $|f(z)|$ is not.

But $\log|f(z)|$ is harmonic (where f is nonzero).

The real and imaginary parts of a holomorphic function are harmonic.

The modulus $|f(z)|$ is not.

But $\log|f(z)|$ is harmonic (where f is nonzero).

Proposition

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic and never zero. Then

$$z \mapsto \log|f(z)|$$

is harmonic.

Proof: Exercise.

Exercise: Suppose $U \subset \mathbb{C}$ is a simply connected domain and $f: U \rightarrow \mathbb{R}$ harmonic. Prove there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f(z) = \log|\varphi(z)|$.

Exercise: Suppose $U \subset \mathbb{C}$ is a simply connected domain and $f: U \rightarrow \mathbb{R}$ harmonic. Prove there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f(z) = \log|\varphi(z)|$.

Exercise: Let $U, V \subset \mathbb{C}$ be open sets and $f: U \rightarrow V$ be holomorphic. Prove:

a) If $g: V \rightarrow \mathbb{R}$ is harmonic, then $g \circ f$ is harmonic.

b) If f is a biholomorphism, then $g: V \rightarrow \mathbb{R}$ is harmonic if and only if $g \circ f$ is harmonic.

Exercise: Suppose $U \subset \mathbb{C}$ is a simply connected domain and $f: U \rightarrow \mathbb{R}$ harmonic. Prove there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f(z) = \log|\varphi(z)|$.

Exercise: Let $U, V \subset \mathbb{C}$ be open sets and $f: U \rightarrow V$ be holomorphic. Prove:

- a) If $g: V \rightarrow \mathbb{R}$ is harmonic, then $g \circ f$ is harmonic.
- b) If f is a biholomorphism, then $g: V \rightarrow \mathbb{R}$ is harmonic if and only if $g \circ f$ is harmonic.

Exercise: Prove the Liouville theorem for harmonic functions: If $f: \mathbb{C} \rightarrow \mathbb{R}$ is harmonic and nonnegative, then f is constant.

Exercise: Suppose $U \subset \mathbb{C}$ is a simply connected domain and $f: U \rightarrow \mathbb{R}$ harmonic. Prove there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f(z) = \log|\varphi(z)|$.

Exercise: Let $U, V \subset \mathbb{C}$ be open sets and $f: U \rightarrow V$ be holomorphic. Prove:

- a) If $g: V \rightarrow \mathbb{R}$ is harmonic, then $g \circ f$ is harmonic.
- b) If f is a biholomorphism, then $g: V \rightarrow \mathbb{R}$ is harmonic if and only if $g \circ f$ is harmonic.

Exercise: Prove the Liouville theorem for harmonic functions: If $f: \mathbb{C} \rightarrow \mathbb{R}$ is harmonic and nonnegative, then f is constant.

“bounded” for holomorphic functions \leftrightarrow “nonnegative” for harmonic functions:

Exercise: Suppose $U \subset \mathbb{C}$ is a simply connected domain and $f: U \rightarrow \mathbb{R}$ harmonic. Prove there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f(z) = \log|\varphi(z)|$.

Exercise: Let $U, V \subset \mathbb{C}$ be open sets and $f: U \rightarrow V$ be holomorphic. Prove:

a) If $g: V \rightarrow \mathbb{R}$ is harmonic, then $g \circ f$ is harmonic.

b) If f is a biholomorphism, then $g: V \rightarrow \mathbb{R}$ is harmonic if and only if $g \circ f$ is harmonic.

Exercise: Prove the Liouville theorem for harmonic functions: If $f: \mathbb{C} \rightarrow \mathbb{R}$ is harmonic and nonnegative, then f is constant.

“bounded” for holomorphic functions \leftrightarrow “nonnegative” for harmonic functions:

If f is bounded and holomorphic,

then $\log|f(z) + M|$ or $\operatorname{Re}f(z) + M$ is nonnegative for large enough M .

Exercise: Suppose $U \subset \mathbb{C}$ is a simply connected domain and $f: U \rightarrow \mathbb{R}$ harmonic. Prove there exists a holomorphic $\varphi: U \rightarrow \mathbb{C}$ such that $f(z) = \log|\varphi(z)|$.

Exercise: Let $U, V \subset \mathbb{C}$ be open sets and $f: U \rightarrow V$ be holomorphic. Prove:

a) If $g: V \rightarrow \mathbb{R}$ is harmonic, then $g \circ f$ is harmonic.

b) If f is a biholomorphism, then $g: V \rightarrow \mathbb{R}$ is harmonic if and only if $g \circ f$ is harmonic.

Exercise: Prove the Liouville theorem for harmonic functions: If $f: \mathbb{C} \rightarrow \mathbb{R}$ is harmonic and nonnegative, then f is constant.

“bounded” for holomorphic functions \leftrightarrow “nonnegative” for harmonic functions:

If f is bounded and holomorphic,

then $\log|f(z) + M|$ or $\operatorname{Re} f(z) + M$ is nonnegative for large enough M .

Conversely, if $\log|f(z)| \geq 0$, then $\frac{1}{f(z)}$ is bounded,

and if $\operatorname{Re} f(z) \geq 0$, then $\frac{f(z) - 1}{f(z) + 1}$ is bounded.

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

It is like the D'Alembert solution of the one-dimensional wave equation.

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

It is like the D'Alembert solution of the one-dimensional wave equation.

The wave operator is (using (x, t) for tradition's sake):

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2}$$

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

It is like the D'Alembert solution of the one-dimensional wave equation.

The wave operator is (using (x, t) for tradition's sake):

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = \left[\frac{\partial}{\partial t} - \frac{\partial}{\partial x} \right] \left[\frac{\partial}{\partial t} + \frac{\partial}{\partial x} \right].$$

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

It is like the D'Alembert solution of the one-dimensional wave equation.

The wave operator is (using (x, t) for tradition's sake):

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = \left[\frac{\partial}{\partial t} - \frac{\partial}{\partial x} \right] \left[\frac{\partial}{\partial t} + \frac{\partial}{\partial x} \right].$$

Write $\mu = x + t$ and $\eta = x - t$ (characteristic coordinates), then

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = -4 \frac{\partial^2}{\partial \eta \partial \mu}.$$

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

It is like the D'Alembert solution of the one-dimensional wave equation.

The wave operator is (using (x, t) for tradition's sake):

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = \left[\frac{\partial}{\partial t} - \frac{\partial}{\partial x} \right] \left[\frac{\partial}{\partial t} + \frac{\partial}{\partial x} \right].$$

Write $\mu = x + t$ and $\eta = x - t$ (characteristic coordinates), then

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = -4 \frac{\partial^2}{\partial \eta \partial \mu}.$$

A solution f to the wave equation is

$$f(x, t) = A(\mu) + B(\eta) = A(x + t) + B(x - t).$$

Remark: Writing $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ so that we can integrate twice may sound familiar.

It is like the D'Alembert solution of the one-dimensional wave equation.

The wave operator is (using (x, t) for tradition's sake):

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = \left[\frac{\partial}{\partial t} - \frac{\partial}{\partial x} \right] \left[\frac{\partial}{\partial t} + \frac{\partial}{\partial x} \right].$$

Write $\mu = x + t$ and $\eta = x - t$ (characteristic coordinates), then

$$\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} = -4 \frac{\partial^2}{\partial \eta \partial \mu}.$$

A solution f to the wave equation is

$$f(x, t) = A(\mu) + B(\eta) = A(x + t) + B(x - t).$$

Two waves travelling in opposite directions.