

Cultivating Complex Analysis: Linear fractional transformations (1.4 part 1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on a, b, c, d guarantees that the ratio does not simplify.

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on a, b, c, d guarantees that the ratio does not simplify.

If $c = 0$, f can be written as $f(z) = az + b$.

Define $f(\infty) = \infty$.

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on a, b, c, d guarantees that the ratio does not simplify.

If $c = 0$, f can be written as $f(z) = az + b$.

Define $f(\infty) = \infty$.

If $c \neq 0$, the expression is defined on $\mathbb{C} \setminus \{-d/c\}$.

Define $f(-d/c) = \infty$ and $f(\infty) = \frac{a}{c}$.

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on a, b, c, d guarantees that the ratio does not simplify.

If $c = 0$, f can be written as $f(z) = az + b$.

Define $f(\infty) = \infty$.

If $c \neq 0$, the expression is defined on $\mathbb{C} \setminus \{-d/c\}$.

Define $f(-d/c) = \infty$ and $f(\infty) = \frac{a}{c}$.

So takes the Riemann sphere to Riemann sphere.

It is an easy exercise that $f: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ is bijective and continuous.

Any LFT is a composition of:

Any LFT is a composition of:

translations

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

Any LFT is a composition of:

translations

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

complex dilations

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

Any LFT is a composition of:

translations

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

complex dilations

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and *inversions*

$$I(z) = \frac{1}{z}.$$

Any LFT is a composition of:

translations

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

complex dilations

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and *inversions*

$$I(z) = \frac{1}{z}.$$

Consider an LFT $f(z) = \frac{az+b}{cz+d}$. WLOG $c = 1$ or $c = 0$.

Any LFT is a composition of:

translations

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

complex dilations

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and inversions

$$I(z) = \frac{1}{z}.$$

Consider an LFT $f(z) = \frac{az+b}{cz+d}$. WLOG $c = 1$ or $c = 0$.

If $c = 1$, then

$$f(z) = \frac{az + b}{z + d} = \frac{b - ad}{z + d} + a = T_a \left(D_{b-ad} \left(I(T_d(z)) \right) \right).$$

Any LFT is a composition of:

translations

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

complex dilations

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and *inversions*

$$I(z) = \frac{1}{z}.$$

Consider an LFT $f(z) = \frac{az+b}{cz+d}$. WLOG $c = 1$ or $c = 0$.

If $c = 1$, then

$$f(z) = \frac{az + b}{z + d} = \frac{b - ad}{z + d} + a = T_a \left(D_{b-ad} \left(I(T_d(z)) \right) \right).$$

If $c = 0$, assume $d = 1$ and $f(z) = az + b$:

$$f(z) = az + b = T_b(D_a(z)).$$

A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .

A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .

Complex dilation D_a is the normal euclidean dilation (scaling) by $|a|$ and rotation by $\arg a$. It also fixes ∞ .

A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .

Complex dilation D_a is the normal euclidean dilation (scaling) by $|a|$ and rotation by $\arg a$. It also fixes ∞ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .


Complex dilation D_a is the normal euclidean dilation (scaling) by $|a|$ and rotation by $\arg a$. It also fixes ∞ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

The euclidean inversion inverts the distance:

$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.



A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .

Complex dilation D_a is the normal euclidean dilation (scaling) by $|a|$ and rotation by $\arg a$. It also fixes ∞ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.


The euclidean inversion inverts the distance:

$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.

A use for this decomposition (exercise):

LFTs take the set of straight lines and circles to the set of straight lines and circles.



A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .

Complex dilation D_a is the normal euclidean dilation (scaling) by $|a|$ and rotation by $\arg a$. It also fixes ∞ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

The euclidean inversion inverts the distance:


$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.

A use for this decomposition (exercise):

LFTs take the set of straight lines and circles to the set of straight lines and circles.

In fact, any straight line or circle can be taken to any other straight line or circle.



A translation T_a just moves everything in \mathbb{C} in the a direction and fixes ∞ .


Complex dilation D_a is the normal euclidean dilation (scaling) by $|a|$ and rotation by $\arg a$. It also fixes ∞ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

The euclidean inversion inverts the distance:

$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.



A use for this decomposition (exercise):

LFTs take the set of straight lines and circles to the set of straight lines and circles.

In fact, any straight line or circle can be taken to any other straight line or circle.

Remark: A straight line is really a very large circle through ∞ .