

# Cultivating Complex Analysis: Schwarz's lemma (3.5.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

### Lemma (Schwarz's lemma)

*Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then*

### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

- (i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and

### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

- (i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and
- (ii)  $|f'(0)| \leq 1$ .

### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

- (i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and
- (ii)  $|f'(0)| \leq 1$ .

Furthermore, if  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\}$  or  $|f'(0)| = 1$ , then there is a  $\theta \in \mathbb{R}$  such that  $f(z) = e^{i\theta} z$  for all  $z \in \mathbb{D}$ .

### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

- (i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and
- (ii)  $|f'(0)| \leq 1$ .

Furthermore, if  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\}$  or  $|f'(0)| = 1$ , then there is a  $\theta \in \mathbb{R}$  such that  $f(z) = e^{i\theta} z$  for all  $z \in \mathbb{D}$ .

This may sound very specialized, but

### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

(i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and

(ii)  $|f'(0)| \leq 1$ .

Furthermore, if  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\}$  or  $|f'(0)| = 1$ , then there is a  $\theta \in \mathbb{R}$  such that  $f(z) = e^{i\theta} z$  for all  $z \in \mathbb{D}$ .

This may sound very specialized, but

a) A disc is a basic neighborhood and any disc can be translated and rescaled into  $\mathbb{D}$ .

### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

(i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and

(ii)  $|f'(0)| \leq 1$ .

Furthermore, if  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\}$  or  $|f'(0)| = 1$ , then there is a  $\theta \in \mathbb{R}$  such that  $f(z) = e^{i\theta} z$  for all  $z \in \mathbb{D}$ .

This may sound very specialized, but

a) A disc is a basic neighborhood and any disc can be translated and rescaled into  $\mathbb{D}$ .  
The lemma is telling us about local behavior of a holomorphic function.



### Lemma (Schwarz's lemma)

Suppose  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic and  $f(0) = 0$ , then

(i)  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and

(ii)  $|f'(0)| \leq 1$ .

Furthermore, if  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\}$  or  $|f'(0)| = 1$ , then there is a  $\theta \in \mathbb{R}$  such that  $f(z) = e^{i\theta}z$  for all  $z \in \mathbb{D}$ .

This may sound very specialized, but

a) A disc is a basic neighborhood and any disc can be translated and rescaled into  $\mathbb{D}$ .

The lemma is telling us about local behavior of a holomorphic function.

b) We'll see later that every domain "without holes" (except  $\mathbb{C}$ ) is biholomorphic to  $\mathbb{D}$ , so it tells us about global behavior as well.

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1}$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|}$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ .



**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D},$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\}$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$  attains a maximum in  $\mathbb{D}$

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$  attains a maximum in  $\mathbb{D} \Rightarrow g$  is constant

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$  attains a maximum in  $\mathbb{D} \Rightarrow g$  is constant  $\Rightarrow f(z) = e^{i\theta} z$ .

**Proof:** As  $f(0) = 0$ ,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take  $0 < r < 1$ . For  $z \in \partial\Delta_r(0)$ ,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle,  $|g(z)| \leq \frac{1}{r}$  for all  $z \in \Delta_r(0)$ .

Fix any  $z \in \mathbb{D}$  and take the limit as  $r \uparrow 1$  to find  $|g(z)| \leq 1$ . So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If  $|f(z_0)| = |z_0|$  for some  $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$  attains a maximum in  $\mathbb{D} \Rightarrow g$  is constant  $\Rightarrow f(z) = e^{i\theta} z$ .

As  $g(0) = f'(0)$ , the same conclusion holds if  $|f'(0)| = 1$ .



Consider the statement for  $f(z) = z^n$  for  $n > 1$ .



Consider the statement for  $f(z) = z^n$  for  $n > 1$ .

$f$  takes  $\mathbb{D}$  to  $\mathbb{D}$  and  $f(0) = 0$ .

Consider the statement for  $f(z) = z^n$  for  $n > 1$ .

$f$  takes  $\mathbb{D}$  to  $\mathbb{D}$  and  $f(0) = 0$ .

For  $z \in \mathbb{D} \setminus \{0\}$ ,

$$|z^n| = |z|^n < |z|.$$

Consider the statement for  $f(z) = z^n$  for  $n > 1$ .

$f$  takes  $\mathbb{D}$  to  $\mathbb{D}$  and  $f(0) = 0$ .

For  $z \in \mathbb{D} \setminus \{0\}$ ,

$$|z^n| = |z|^n < |z|.$$

As  $f'(z) = nz^{n-1}$ ,  $|f'(0)| = 0 < 1$ .

Consider the statement for  $f(z) = z^n$  for  $n > 1$ .

$f$  takes  $\mathbb{D}$  to  $\mathbb{D}$  and  $f(0) = 0$ .

For  $z \in \mathbb{D} \setminus \{0\}$ ,

$$|z^n| = |z|^n < |z|.$$

As  $f'(z) = nz^{n-1}$ ,  $|f'(0)| = 0 < 1$ .

A bound on the derivative does not hold at other points:  
Picking the right  $z$  and  $n$ , can make  $|f'(z)|$  arbitrarily large.

Consider the statement for  $f(z) = z^n$  for  $n > 1$ .

$f$  takes  $\mathbb{D}$  to  $\mathbb{D}$  and  $f(0) = 0$ .

For  $z \in \mathbb{D} \setminus \{0\}$ ,

$$|z^n| = |z|^n < |z|.$$

As  $f'(z) = nz^{n-1}$ ,  $|f'(0)| = 0 < 1$ .

A bound on the derivative does not hold at other points:

Picking the right  $z$  and  $n$ , can make  $|f'(z)|$  arbitrarily large.

We can make  $|z^n|$  arbitrarily small for a fixed  $z \in \mathbb{D}$  by picking a large enough  $n$ , but we cannot make it bigger than  $|z|$ .

Consider the statement for  $f(z) = z^n$  for  $n > 1$ .

$f$  takes  $\mathbb{D}$  to  $\mathbb{D}$  and  $f(0) = 0$ .

For  $z \in \mathbb{D} \setminus \{0\}$ ,

$$|z^n| = |z|^n < |z|.$$

As  $f'(z) = nz^{n-1}$ ,  $|f'(0)| = 0 < 1$ .

A bound on the derivative does not hold at other points:

Picking the right  $z$  and  $n$ , can make  $|f'(z)|$  arbitrarily large.

We can make  $|z^n|$  arbitrarily small for a fixed  $z \in \mathbb{D}$  by picking a large enough  $n$ , but we cannot make it bigger than  $|z|$ .

Schwarz's lemma says all holomorphic functions behave this way, not just  $z^n$ .

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:



It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose  $U \subset \mathbb{C}$  is bounded,  $f: U \rightarrow U$  is holomorphic,  $p \in U$ , and  $f(p) = p$ . Then

- (i)  $|f'(p)| \leq 1$ .
- (ii) If  $f'(p) = 1$ , then  $f(z) = z$  for all  $z \in U$ .

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose  $U \subset \mathbb{C}$  is bounded,  $f: U \rightarrow U$  is holomorphic,  $p \in U$ , and  $f(p) = p$ . Then

(i)  $|f'(p)| \leq 1$ .

(ii) If  $f'(p) = 1$ , then  $f(z) = z$  for all  $z \in U$ .

Hint: WLOG  $p = 0$ ,

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose  $U \subset \mathbb{C}$  is bounded,  $f: U \rightarrow U$  is holomorphic,  $p \in U$ , and  $f(p) = p$ . Then

(i)  $|f'(p)| \leq 1$ .

(ii) If  $f'(p) = 1$ , then  $f(z) = z$  for all  $z \in U$ .

Hint: WLOG  $p = 0$ , then consider the power series expansions of  $f^\ell$ , the  $\ell^{\text{th}}$  composition of  $f$  with itself,  $f(f(f(\cdots f(z) \cdots)))$ .

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose  $U \subset \mathbb{C}$  is bounded,  $f: U \rightarrow U$  is holomorphic,  $p \in U$ , and  $f(p) = p$ . Then

(i)  $|f'(p)| \leq 1$ .

(ii) If  $f'(p) = 1$ , then  $f(z) = z$  for all  $z \in U$ .

Hint: WLOG  $p = 0$ , then consider the power series expansions of  $f^\ell$ , the  $\ell^{\text{th}}$  composition of  $f$  with itself,  $f(f(f(\cdots f(z) \cdots)))$ .

For (i) consider the linear term of  $f^\ell$ .

It is a useful exercise to rewrite the lemma for  $f: \Delta_r(p) \rightarrow \Delta_s(q)$  with  $f(p) = q$ .

Using an LFT we can find a half-plane ( $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ ) version:

If  $f: \mathbb{H} \rightarrow \mathbb{H}$  holomorphic and  $f(i) = i$ , then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose  $U \subset \mathbb{C}$  is bounded,  $f: U \rightarrow U$  is holomorphic,  $p \in U$ , and  $f(p) = p$ . Then

(i)  $|f'(p)| \leq 1$ .

(ii) If  $f'(p) = 1$ , then  $f(z) = z$  for all  $z \in U$ .

Hint: WLOG  $p = 0$ , then consider the power series expansions of  $f^\ell$ , the  $\ell^{\text{th}}$  composition of  $f$  with itself,  $f(f(f(\cdots f(z) \cdots)))$ .

For (i) consider the linear term of  $f^\ell$ .

For (ii) use Cauchy estimates on the first nonzero nonlinear term of  $f^\ell$ .