

Cultivating Complex Analysis: Automorphisms of the disc (3.5.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

We start with certain specific automorphisms.

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

We start with certain specific automorphisms.

For $a \in \mathbb{D}$, define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

We start with certain specific automorphisms.

For $a \in \mathbb{D}$, define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

Proposition

For every $a \in \mathbb{D}$,

(i) $\varphi_a(a) = 0$, $\varphi_a(0) = -a$, $\varphi'_a(0) = 1 - |a|^2$, $\varphi'_a(a) = \frac{1}{1 - |a|^2}$,

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

We start with certain specific automorphisms.

For $a \in \mathbb{D}$, define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

Proposition

For every $a \in \mathbb{D}$,

- (i) $\varphi_a(a) = 0$, $\varphi_a(0) = -a$, $\varphi'_a(0) = 1 - |a|^2$, $\varphi'_a(a) = \frac{1}{1 - |a|^2}$,
- (ii) $\varphi_a(\partial\mathbb{D}) = \partial\mathbb{D}$, and $\varphi_a(\mathbb{D}) = \mathbb{D}$,

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

We start with certain specific automorphisms.

For $a \in \mathbb{D}$, define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

Proposition

For every $a \in \mathbb{D}$,

- (i) $\varphi_a(a) = 0$, $\varphi_a(0) = -a$, $\varphi'_a(0) = 1 - |a|^2$, $\varphi'_a(a) = \frac{1}{1 - |a|^2}$,
- (ii) $\varphi_a(\partial\mathbb{D}) = \partial\mathbb{D}$, and $\varphi_a(\mathbb{D}) = \mathbb{D}$,
- (iii) φ_a restricted to \mathbb{D} is an automorphism of the disc and $\varphi_a^{-1} = \varphi_{-a}$.

Recall, an automorphism of U is a biholomorphism $f: U \rightarrow U$ (bijective such that f and f^{-1} are both holomorphic).

Let us compute the automorphism group of the disc, $\text{Aut}(\mathbb{D})$.

We start with certain specific automorphisms.

For $a \in \mathbb{D}$, define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$


Proposition

For every $a \in \mathbb{D}$,

- (i) $\varphi_a(a) = 0$, $\varphi_a(0) = -a$, $\varphi'_a(0) = 1 - |a|^2$, $\varphi'_a(a) = \frac{1}{1 - |a|^2}$,
- (ii) $\varphi_a(\partial\mathbb{D}) = \partial\mathbb{D}$, and $\varphi_a(\mathbb{D}) = \mathbb{D}$,
- (iii) φ_a restricted to \mathbb{D} is an automorphism of the disc and $\varphi_a^{-1} = \varphi_{-a}$.


Proof is an exercise.

Here is what an automorphism φ_a does ($a = -0.4$).



a and $\varphi_a(a) = 0$ are marked with dots.


Here is what an automorphism φ_a does ($a = -0.4$).



a and $\varphi_a(a) = 0$ are marked with dots.

It turns out all automorphisms are φ_a up to a rotation:

Here is what an automorphism φ_a does ($a = -0.4$).



a and $\varphi_a(a) = 0$ are marked with dots.

It turns out all automorphisms are φ_a up to a rotation:

Proposition

If $f \in \text{Aut}(\mathbb{D})$, then there exists an $a \in \mathbb{D}$ and $\theta \in \mathbb{R}$ such that

$$f(z) = e^{i\theta} \frac{z - a}{1 - \bar{a}z} = e^{i\theta} \varphi_a(z).$$

Proof: Let $a = f(0)$.

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros:

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,

and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,

and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise)

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$
 $\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

$\Rightarrow |h(z)|$ attains a minimum in \mathbb{D}

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

$\Rightarrow |h(z)|$ attains a minimum in $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$ attains a maximum in \mathbb{D}

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

$\Rightarrow |h(z)|$ attains a minimum in $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$ attains a maximum in $\mathbb{D} \Rightarrow h$ is constant

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

$\Rightarrow |h(z)|$ attains a minimum in $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$ attains a maximum in $\mathbb{D} \Rightarrow h$ is constant

$\Rightarrow g(z) = \alpha z$ for some constant α .

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

$\Rightarrow |h(z)|$ attains a minimum in $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$ attains a maximum in $\mathbb{D} \Rightarrow h$ is constant

$\Rightarrow g(z) = \alpha z$ for some constant α . As $g \in \text{Aut}(\mathbb{D})$, $|\alpha| = 1$ ($\alpha = e^{i\theta}$).

Proof: Let $a = f(0)$.

$g = \varphi_a \circ f$ is an automorphism with $g(0) = 0$ as $\varphi_a(a) = 0$.

Find a holomorphic $h(z)$ such that $g(z) = zh(z)$ (as in the proof of Schwarz).

Schwarz says that for $z \in \mathbb{D} \setminus \{0\}$, $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$.

h can have no zeros: $h(z) = \frac{g(z)}{z}$ cannot be zero for $z \neq 0$ as g is injective,
and it cannot have a zero at $z = 0$ as $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$.

g is biholomorphic $\Rightarrow g^{-1}$ is continuous $\Rightarrow g^{-1}(K)$ compact for compact $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$ as $|z| \rightarrow 1$ (exercise) $\Rightarrow |h(z)| \rightarrow 1$ as $|z| \rightarrow 1$

$\Rightarrow |h(z)|$ attains a minimum in $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$ attains a maximum in $\mathbb{D} \Rightarrow h$ is constant

$\Rightarrow g(z) = \alpha z$ for some constant α . As $g \in \text{Aut}(\mathbb{D})$, $|\alpha| = 1$ ($\alpha = e^{i\theta}$).

Apply φ_{-a} to both sides of $e^{i\theta}z = \varphi_a \circ f$ to find $f(z) = \varphi_{-a}(ze^{i\theta}) = e^{i\theta} \varphi_{-ae^{-i\theta}}(z)$.



Some good exercises to try:

Exercise: Given distinct $a, b \in \mathbb{D}$, show that there exists a unique $f \in \text{Aut}(\mathbb{D})$ such that $f(a) = b$ and $f(b) = a$.

Some good exercises to try:

Exercise: Given distinct $a, b \in \mathbb{D}$, show that there exists a unique $f \in \text{Aut}(\mathbb{D})$ such that $f(a) = b$ and $f(b) = a$.

Exercise: The automorphisms of $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ are of the form $\frac{az+b}{cz+d}$ for real numbers a, b, c, d such that $ad - bc \neq 0$.

Some good exercises to try:

Exercise: Given distinct $a, b \in \mathbb{D}$, show that there exists a unique $f \in \text{Aut}(\mathbb{D})$ such that $f(a) = b$ and $f(b) = a$.

Exercise: The automorphisms of $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ are of the form $\frac{az+b}{cz+d}$ for real numbers a, b, c, d such that $ad - bc \neq 0$.

Another fun exercise, a version of which seems to appear on many exams is the following:

Exercise: Suppose $U \subset \mathbb{C}$ is a domain, $\overline{\mathbb{D}} \subset U$, and $f: U \rightarrow \mathbb{C}$ is holomorphic such that $f(\partial\mathbb{D}) \subset \partial\mathbb{D}$. Find a formula for f .

Some good exercises to try:

Exercise: Given distinct $a, b \in \mathbb{D}$, show that there exists a unique $f \in \text{Aut}(\mathbb{D})$ such that $f(a) = b$ and $f(b) = a$.

Exercise: The automorphisms of $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ are of the form $\frac{az+b}{cz+d}$ for real numbers a, b, c, d such that $ad - bc \neq 0$.

Another fun exercise, a version of which seems to appear on many exams is the following:

Exercise: Suppose $U \subset \mathbb{C}$ is a domain, $\overline{\mathbb{D}} \subset U$, and $f: U \rightarrow \mathbb{C}$ is holomorphic such that $f(\partial\mathbb{D}) \subset \partial\mathbb{D}$. Find a formula for f .

Hint: The idea is to show that you can divide by finitely many $\varphi_a(z)$ for various a until you get something that has no zeros in \mathbb{D} and will have to be a constant.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Lemma (Schwarz–Pick)

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Lemma (Schwarz–Pick)

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \overline{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some $z \neq \zeta$, then $f \in \text{Aut}(\mathbb{D})$.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Lemma (Schwarz–Pick)

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some $z \neq \zeta$, then $f \in \text{Aut}(\mathbb{D})$.

Conversely, if $f \in \text{Aut}(\mathbb{D})$, then equality holds in both inequalities for all $z, \zeta \in \mathbb{D}$.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Lemma (Schwarz–Pick)

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some $z \neq \zeta$, then $f \in \text{Aut}(\mathbb{D})$.

Conversely, if $f \in \text{Aut}(\mathbb{D})$, then equality holds in both inequalities for all $z, \zeta \in \mathbb{D}$.

Proof is an exercise.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Lemma (Schwarz–Pick)

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \overline{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some $z \neq \zeta$, then $f \in \text{Aut}(\mathbb{D})$.

Conversely, if $f \in \text{Aut}(\mathbb{D})$, then equality holds in both inequalities for all $z, \zeta \in \mathbb{D}$.

Proof is an exercise.

The Schwarz–Pick lemma gives a bound on the derivative at all points:

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, nonconstant, and $f(a) = b$, then

$$|f'(a)| \leq \frac{1 - |b|^2}{1 - |a|^2}.$$

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Lemma (Schwarz–Pick)

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \overline{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some $z \neq \zeta$, then $f \in \text{Aut}(\mathbb{D})$.

Conversely, if $f \in \text{Aut}(\mathbb{D})$, then equality holds in both inequalities for all $z, \zeta \in \mathbb{D}$.

Proof is an exercise.

The Schwarz–Pick lemma gives a bound on the derivative at all points:

If $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic, nonconstant, and $f(a) = b$, then

$$|f'(a)| \leq \frac{1 - |b|^2}{1 - |a|^2}.$$

Equality $\Rightarrow f(z) = \varphi_{-b}(e^{i\theta} \varphi_a(z))$ for some $\theta \in \mathbb{R}$.