

# Cultivating Complex Analysis: Wirtinger operators (2.2.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Suppose  $z = x + iy$ .

Suppose  $z = x + iy$ .

Define the *Wirtinger operators*:

$$\frac{\partial}{\partial z} \stackrel{\text{def}}{=} \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} \stackrel{\text{def}}{=} \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

Suppose  $z = x + iy$ .

Define the *Wirtinger operators*:

$$\frac{\partial}{\partial z} \stackrel{\text{def}}{=} \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} \stackrel{\text{def}}{=} \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

Not really “partial derivatives” despite the notation.

Suppose  $z = x + iy$ .

Define the *Wirtinger operators*:

$$\frac{\partial}{\partial z} \stackrel{\text{def}}{=} \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} \stackrel{\text{def}}{=} \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

Not really “partial derivatives” despite the notation.

The operators are really determined by wanting

$$\frac{\partial}{\partial z} z = 1, \quad \frac{\partial}{\partial z} \bar{z} = 0, \quad \frac{\partial}{\partial \bar{z}} z = 0, \quad \frac{\partial}{\partial \bar{z}} \bar{z} = 1.$$

Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right)$$

Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right)$$

Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right)$$



Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right)$$

This expression is zero if and only if the real and imaginary parts are zero.

Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right)$$

This expression is zero if and only if the real and imaginary parts are zero. Namely,

$$\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} = 0, \quad \text{and} \quad \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} = 0.$$

Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right)$$

This expression is zero if and only if the real and imaginary parts are zero. Namely,

$$\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} = 0, \quad \text{and} \quad \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} = 0.$$

Those are the Cauchy–Riemann equations!

Let  $f = u + iv$ .

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) = \frac{1}{2} \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right)$$

This expression is zero if and only if the real and imaginary parts are zero. Namely,

$$\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} = 0, \quad \text{and} \quad \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} = 0.$$

Those are the Cauchy–Riemann equations!

### Proposition

*Let  $U \subset \mathbb{C}$  be open. Then  $f: U \rightarrow \mathbb{C}$  is holomorphic if and only if  $f$  is (real) differentiable and*

$$\frac{\partial f}{\partial \bar{z}} \equiv 0.$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial \bar{z}}$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x}$$



On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right)$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h}$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t}$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z = \frac{\partial f}{\partial x} \Big|_z,$$



On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z = \frac{\partial f}{\partial x} \Big|_z,$$

and

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h}$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z = \frac{\partial f}{\partial x} \Big|_z,$$

and

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+it) - f(z)}{it}$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z = \frac{\partial f}{\partial x} \Big|_z,$$

and

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+it) - f(z)}{it} = \frac{1}{i} \left( \frac{\partial u}{\partial y} \Big|_z + i \frac{\partial v}{\partial y} \Big|_z \right)$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z = \frac{\partial f}{\partial x} \Big|_z,$$

and

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+it) - f(z)}{it} = \frac{1}{i} \left( \frac{\partial u}{\partial y} \Big|_z + i \frac{\partial v}{\partial y} \Big|_z \right) = \frac{1}{i} \frac{\partial f}{\partial y} \Big|_z.$$

On the other hand, if  $f$  is holomorphic,

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial f}{\partial x} = \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{1}{i} \frac{\partial f}{\partial y}.$$

(In the second form think of derivative in  $iy$  not  $y$ ).

To compute  $f'(z)$  we can approach from any direction:

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+t) - f(z)}{t} = \frac{\partial u}{\partial x} \Big|_z + i \frac{\partial v}{\partial x} \Big|_z = \frac{\partial f}{\partial x} \Big|_z,$$

and

$$f'(z) = \lim_{\substack{h \rightarrow 0 \\ h \in \mathbb{C}}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{t \rightarrow 0 \\ t \in \mathbb{R}}} \frac{f(z+it) - f(z)}{it} = \frac{1}{i} \left( \frac{\partial u}{\partial y} \Big|_z + i \frac{\partial v}{\partial y} \Big|_z \right) = \frac{1}{i} \frac{\partial f}{\partial y} \Big|_z.$$

So for a holomorphic function

$$f' = \frac{\partial f}{\partial z}.$$

The complex derivative  $f'$ , sometimes written as  $\frac{df}{dz}$ , only exists for holomorphic functions.

The complex derivative  $f'$ , sometimes written as  $\frac{df}{dz}$ , only exists for holomorphic functions.

The Wirtinger operators  $\frac{\partial f}{\partial z}$  and  $\frac{\partial f}{\partial \bar{z}}$  make sense for every real differentiable function.

The complex derivative  $f'$ , sometimes written as  $\frac{df}{dz}$ , only exists for holomorphic functions.

The Wirtinger operators  $\frac{\partial f}{\partial z}$  and  $\frac{\partial f}{\partial \bar{z}}$  make sense for every real differentiable function.

For polynomials the operators work as if  $z$  and  $\bar{z}$  were separate variables. E.g. (exercise)

$$\frac{\partial}{\partial z} [z^2 \bar{z}^3 + z^{10}] = 2z \bar{z}^3 + 10z^9 \quad \text{and} \quad \frac{\partial}{\partial \bar{z}} [z^2 \bar{z}^3 + z^{10}] = z^2 (3\bar{z}^2) + 0.$$



The complex derivative  $f'$ , sometimes written as  $\frac{df}{dz}$ , only exists for holomorphic functions.

The Wirtinger operators  $\frac{\partial f}{\partial z}$  and  $\frac{\partial f}{\partial \bar{z}}$  make sense for every real differentiable function.

For polynomials the operators work as if  $z$  and  $\bar{z}$  were separate variables. E.g. (exercise)

$$\frac{\partial}{\partial z} [z^2 \bar{z}^3 + z^{10}] = 2z \bar{z}^3 + 10z^9 \quad \text{and} \quad \frac{\partial}{\partial \bar{z}} [z^2 \bar{z}^3 + z^{10}] = z^2 (3\bar{z}^2) + 0.$$

So a holomorphic function is “one that does not depend on  $\bar{z}$ .”

The complex derivative  $f'$ , sometimes written as  $\frac{df}{dz}$ , only exists for holomorphic functions.

The Wirtinger operators  $\frac{\partial f}{\partial z}$  and  $\frac{\partial f}{\partial \bar{z}}$  make sense for every real differentiable function.

For polynomials the operators work as if  $z$  and  $\bar{z}$  were separate variables. E.g. (exercise)

$$\frac{\partial}{\partial z} [z^2 \bar{z}^3 + z^{10}] = 2z \bar{z}^3 + 10z^9 \quad \text{and} \quad \frac{\partial}{\partial \bar{z}} [z^2 \bar{z}^3 + z^{10}] = z^2(3\bar{z}^2) + 0.$$

So a holomorphic function is “one that does not depend on  $\bar{z}$ .”

**Caution:** Note that  $\frac{d}{dz} [z^2 \bar{z}^3 + z^{10}]$  does not exist, while  $\frac{\partial}{\partial z} [z^2 \bar{z}^3 + z^{10}]$  does.

A few other remarks that are left as exercises.

A few other remarks that are left as exercises.

**Remark 1:** If we write the real derivative  $Df|_p$  as the map  $h \mapsto \zeta h + \xi \bar{h}$ ,

A few other remarks that are left as exercises.

**Remark 1:** If we write the real derivative  $Df|_p$  as the map  $h \mapsto \zeta h + \xi \bar{h}$ , then

$$\frac{\partial f}{\partial z}\bigg|_p = \zeta \quad \text{and} \quad \frac{\partial f}{\partial \bar{z}}\bigg|_p = \xi.$$

A few other remarks that are left as exercises.

**Remark 1:** If we write the real derivative  $Df|_p$  as the map  $h \mapsto \zeta h + \xi \bar{h}$ , then

$$\frac{\partial f}{\partial z}|_p = \zeta \quad \text{and} \quad \frac{\partial f}{\partial \bar{z}}|_p = \xi.$$

**Remark 2:** If  $\bar{f}$  is the complex conjugate of  $f$ , then

A few other remarks that are left as exercises.

**Remark 1:** If we write the real derivative  $Df|_p$  as the map  $h \mapsto \zeta h + \xi \bar{h}$ , then

$$\frac{\partial f}{\partial z}|_p = \zeta \quad \text{and} \quad \frac{\partial f}{\partial \bar{z}}|_p = \xi.$$

**Remark 2:** If  $\bar{f}$  is the complex conjugate of  $f$ , then

$$\overline{\left(\frac{\partial f}{\partial z}\right)} = \frac{\partial \bar{f}}{\partial \bar{z}} \quad \text{and} \quad \overline{\left(\frac{\partial f}{\partial \bar{z}}\right)} = \frac{\partial \bar{f}}{\partial z}.$$

A few other remarks that are left as exercises.

**Remark 1:** If we write the real derivative  $Df|_p$  as the map  $h \mapsto \zeta h + \xi \bar{h}$ , then

$$\frac{\partial f}{\partial z}\Big|_p = \zeta \quad \text{and} \quad \frac{\partial f}{\partial \bar{z}}\Big|_p = \xi.$$

**Remark 2:** If  $\bar{f}$  is the complex conjugate of  $f$ , then

$$\overline{\left(\frac{\partial f}{\partial z}\right)} = \frac{\partial \bar{f}}{\partial \bar{z}} \quad \text{and} \quad \overline{\left(\frac{\partial f}{\partial \bar{z}}\right)} = \frac{\partial \bar{f}}{\partial z}.$$

**Remark 3:** The chain rule for real differentiable functions can be written as

$$\frac{\partial(g \circ f)}{\partial z}\Big|_p = \frac{\partial g}{\partial z}\Big|_{f(p)} \frac{\partial f}{\partial z}\Big|_p + \frac{\partial g}{\partial \bar{z}}\Big|_{f(p)} \frac{\partial \bar{f}}{\partial z}\Big|_p \quad \text{and} \quad \frac{\partial(g \circ f)}{\partial \bar{z}}\Big|_p = \frac{\partial g}{\partial z}\Big|_{f(p)} \frac{\partial f}{\partial \bar{z}}\Big|_p + \frac{\partial g}{\partial \bar{z}}\Big|_{f(p)} \frac{\partial \bar{f}}{\partial \bar{z}}\Big|_p.$$