

Cultivating Complex Analysis: Montel's theorem (6.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Let $U \subset \mathbb{C}$ be open. A set \mathcal{F} of holomorphic functions $f: U \rightarrow \mathbb{C}$ is called a *normal family* if every sequence in \mathcal{F} has a subsequence that converges uniformly on compact subsets (the limit need not be in \mathcal{F}).

Definition

Let $U \subset \mathbb{C}$ be open. A set \mathcal{F} of holomorphic functions $f: U \rightarrow \mathbb{C}$ is called a *normal family* if every sequence in \mathcal{F} has a subsequence that converges uniformly on compact subsets (the limit need not be in \mathcal{F}).

A set \mathcal{F} of functions on U is *locally bounded* if for every $p \in U$, there is a disc $\Delta_r(p) \subset U$ and $M > 0$ such that $\|f\|_{\Delta_r(p)} \leq M$ for all $f \in \mathcal{F}$.

Definition

Let $U \subset \mathbb{C}$ be open. A set \mathcal{F} of holomorphic functions $f: U \rightarrow \mathbb{C}$ is called a *normal family* if every sequence in \mathcal{F} has a subsequence that converges uniformly on compact subsets (the limit need not be in \mathcal{F}).

A set \mathcal{F} of functions on U is *locally bounded* if for every $p \in U$, there is a disc $\Delta_r(p) \subset U$ and $M > 0$ such that $\|f\|_{\Delta_r(p)} \leq M$ for all $f \in \mathcal{F}$ (i.e., $|f(z)| \leq M$ for all $z \in \Delta_r(p)$ and all $f \in \mathcal{F}$).

Definition

Let $U \subset \mathbb{C}$ be open. A set \mathcal{F} of holomorphic functions $f: U \rightarrow \mathbb{C}$ is called a *normal family* if every sequence in \mathcal{F} has a subsequence that converges uniformly on compact subsets (the limit need not be in \mathcal{F}).

A set \mathcal{F} of functions on U is *locally bounded* if for every $p \in U$, there is a disc $\Delta_r(p) \subset U$ and $M > 0$ such that $\|f\|_{\Delta_r(p)} \leq M$ for all $f \in \mathcal{F}$ (i.e., $|f(z)| \leq M$ for all $z \in \Delta_r(p)$ and all $f \in \mathcal{F}$).

In more modern language:

\mathcal{F} normal family

=

\mathcal{F} precompact in the space of holomorphic functions on U with the topology of uniform convergence on compact subsets.

Definition

Let $U \subset \mathbb{C}$ be open. A set \mathcal{F} of holomorphic functions $f: U \rightarrow \mathbb{C}$ is called a *normal family* if every sequence in \mathcal{F} has a subsequence that converges uniformly on compact subsets (the limit need not be in \mathcal{F}).

A set \mathcal{F} of functions on U is *locally bounded* if for every $p \in U$, there is a disc $\Delta_r(p) \subset U$ and $M > 0$ such that $\|f\|_{\Delta_r(p)} \leq M$ for all $f \in \mathcal{F}$ (i.e., $|f(z)| \leq M$ for all $z \in \Delta_r(p)$ and all $f \in \mathcal{F}$).

In more modern language:

\mathcal{F} normal family

=

\mathcal{F} precompact in the space of holomorphic functions on U with the topology of uniform convergence on compact subsets.

Exercise: Prove that “locally bounded” means “bounded on compact subsets.”

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$.

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right|$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta|$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta|$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right|$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta|$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

$\Rightarrow \mathcal{F}$ is equicontinuous ($\frac{4M}{r}$ does not depend on f).

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

$\Rightarrow \mathcal{F}$ is equicontinuous ($\frac{4M}{r}$ does not depend on f).

\Rightarrow Arzelà–Ascoli applies to any sequence in \mathcal{F}

Theorem (Montel)

Let $U \subset \mathbb{C}$ be open and let \mathcal{F} be a locally bounded set of holomorphic functions on U . Then \mathcal{F} is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove \mathcal{F} being holomorphic means \mathcal{F} is equicontinuous.

Proof: Consider $p \in U$, $\overline{\Delta_r(p)} \subset U$, such that $\|f\|_{\overline{\Delta_r(p)}} \leq M$ for all $f \in \mathcal{F}$.

If $z \in \overline{\Delta_{r/2}(p)}$ and $\zeta \in \partial\Delta_r(p)$, then $|\zeta - z| \geq r/2$. So for $z \in \overline{\Delta_{r/2}(p)}$ and all $f \in \mathcal{F}$,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

$\Rightarrow \mathcal{F}$ is equicontinuous ($\frac{4M}{r}$ does not depend on f).

\Rightarrow Arzelà–Ascoli applies to any sequence in $\mathcal{F} \Rightarrow \mathcal{F}$ is a normal family. □

Montel's theorem is good for finding extremal functions.

Montel's theorem is good for finding extremal functions.
E.g., we will prove the Riemann mapping theorem:

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected U to \mathbb{D} is a map that maximizes the derivative at a point.

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected U to \mathbb{D} is a map that maximizes the derivative at a point.

Montel gives us a way of finding a maximizer.

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected U to \mathbb{D} is a map that maximizes the derivative at a point.

Montel gives us a way of finding a maximizer.

Another commonly used consequence of Montel is Vitali's theorem.

Theorem (Vitali)

Suppose $U \subset \mathbb{C}$ is a domain, $\{f_n\}$ is a locally bounded sequence of holomorphic functions that converges pointwise on a set $E \subset U$, and E has a limit point in U . Then $\{f_n\}$ converges uniformly on compact subsets in U .

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected U to \mathbb{D} is a map that maximizes the derivative at a point.

Montel gives us a way of finding a maximizer.

Another commonly used consequence of Montel is Vitali's theorem.

Theorem (Vitali)

Suppose $U \subset \mathbb{C}$ is a domain, $\{f_n\}$ is a locally bounded sequence of holomorphic functions that converges pointwise on a set $E \subset U$, and E has a limit point in U . Then $\{f_n\}$ converges uniformly on compact subsets in U .

Proof is an exercise.

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Let $U \subset \mathbb{C}$ be open and \mathcal{F} a normal family of holomorphic functions on U . Show that $\{f' : f \in \mathcal{F}\}$ is a normal family.

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Let $U \subset \mathbb{C}$ be open and \mathcal{F} a normal family of holomorphic functions on U . Show that $\{f' : f \in \mathcal{F}\}$ is a normal family. Note: The converse is false without an extra hypothesis.

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Let $U \subset \mathbb{C}$ be open and \mathcal{F} a normal family of holomorphic functions on U . Show that $\{f' : f \in \mathcal{F}\}$ is a normal family. Note: The converse is false without an extra hypothesis.

Exercise: Let $U \subset \mathbb{C}$ be a domain, $p \in U$, and suppose \exists a nonconstant bounded holomorphic function on U .

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Let $U \subset \mathbb{C}$ be open and \mathcal{F} a normal family of holomorphic functions on U . Show that $\{f' : f \in \mathcal{F}\}$ is a normal family. Note: The converse is false without an extra hypothesis.

Exercise: Let $U \subset \mathbb{C}$ be a domain, $p \in U$, and suppose \exists a nonconstant bounded holomorphic function on U .

a) Prove \exists a holomorphic $F: U \rightarrow \mathbb{D}$ such that $F'(p) \neq 0$, and $|f'(p)| \leq |F'(p)|$ for all holomorphic $f: U \rightarrow \mathbb{D}$.

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Let $U \subset \mathbb{C}$ be open and \mathcal{F} a normal family of holomorphic functions on U . Show that $\{f' : f \in \mathcal{F}\}$ is a normal family. Note: The converse is false without an extra hypothesis.

Exercise: Let $U \subset \mathbb{C}$ be a domain, $p \in U$, and suppose \exists a nonconstant bounded holomorphic function on U .

- a) Prove \exists a holomorphic $F: U \rightarrow \mathbb{D}$ such that $F'(p) \neq 0$, and $|f'(p)| \leq |F'(p)|$ for all holomorphic $f: U \rightarrow \mathbb{D}$.
- b) Show $F(p) = 0$.

Exercise: Prove the converse to Montel: If \mathcal{F} is a normal family of holomorphic functions on an open set $U \subset \mathbb{C}$, then \mathcal{F} is locally bounded.

Exercise: Let $U \subset \mathbb{C}$ be open and \mathcal{F} a normal family of holomorphic functions on U . Show that $\{f' : f \in \mathcal{F}\}$ is a normal family. Note: The converse is false without an extra hypothesis.

Exercise: Let $U \subset \mathbb{C}$ be a domain, $p \in U$, and suppose \exists a nonconstant bounded holomorphic function on U .

a) Prove \exists a holomorphic $F: U \rightarrow \mathbb{D}$ such that $F'(p) \neq 0$, and $|f'(p)| \leq |F'(p)|$ for all holomorphic $f: U \rightarrow \mathbb{D}$.

b) Show $F(p) = 0$.

Exercise: Show that if the partial sums of a power series centered at p are uniformly bounded on $\Delta_r(p)$ for some $r > 0$, then the power series converges in $\Delta_r(p)$.