

# Cultivating Complex Analysis: Riemann mapping theorem (6.3.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Riemann mapping theorem: The only simply connected domains in  $\mathbb{C}$  (up to biholomorphisms) are  $\mathbb{C}$  and  $\mathbb{D}$ .

Riemann mapping theorem: The only simply connected domains in  $\mathbb{C}$  (up to biholomorphisms) are  $\mathbb{C}$  and  $\mathbb{D}$ .

More precisely:

### Theorem (Riemann mapping theorem)

*Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic (conformal) map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .*

Riemann mapping theorem: The only simply connected domains in  $\mathbb{C}$  (up to biholomorphisms) are  $\mathbb{C}$  and  $\mathbb{D}$ .


More precisely:

### Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic (conformal) map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

$U =$  the upper half disk

$p = (\sqrt{2} - 1)i$



Riemann mapping theorem: The only simply connected domains in  $\mathbb{C}$  (up to biholomorphisms) are  $\mathbb{C}$  and  $\mathbb{D}$ .


More precisely:

### Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic (conformal) map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

$U =$  the upper half disk

$p = (\sqrt{2} - 1)i$



Proof is to “maximize”  $|f'(p)|$  among all maps into the disc and  $f(p) = 0$ .

## Theorem (Riemann mapping theorem)

*Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .*

## Theorem (Riemann mapping theorem)

*Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .*

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

## Theorem (Riemann mapping theorem)

*Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .*

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:



## Theorem (Riemann mapping theorem)

*Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .*

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$$g(z) = g(\zeta)$$

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2$$

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta$$

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta)$

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2$



## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g \text{ is injective.}$$

$$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta$$

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

$g(U)$  is open

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

$g(U)$  is open  $\Rightarrow -g(U)$  is open

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

$g(U)$  is open  $\Rightarrow -g(U)$  is open  $\Rightarrow \exists \Delta_r(\xi) \subset \mathbb{C} \setminus g(U)$

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

$g(U)$  is open  $\Rightarrow -g(U)$  is open  $\Rightarrow \exists \Delta_r(\xi) \subset \mathbb{C} \setminus g(U)$

$\Rightarrow z \mapsto \frac{r}{g(z) - \xi}$  takes  $U$  to  $\mathbb{D}$ .

## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

$g(U)$  is open  $\Rightarrow -g(U)$  is open  $\Rightarrow \exists \Delta_r(\xi) \subset \mathbb{C} \setminus g(U)$

$\Rightarrow z \mapsto \frac{r}{g(z) - \xi}$  takes  $U$  to  $\mathbb{D}$ .

Compose with an automorphism of  $\mathbb{D}$  to make  $p$  go to 0



## Theorem (Riemann mapping theorem)

Let  $U \subset \mathbb{C}$  be a simply connected domain such that  $U \neq \mathbb{C}$ . Let  $p \in U$  be given. Then there exists a unique biholomorphic map  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$  and  $f'(p) > 0$ .

**Proof:** Let  $\mathcal{F}$  be the set of injective holomorphic  $f: U \rightarrow \mathbb{D}$  such that  $f(p) = 0$ .

First we need to prove that  $\mathcal{F}$  is nonempty:

Let  $q \in \mathbb{C} \setminus U$ .  $\exists g: U \rightarrow \mathbb{C}$  such that  $(g(z))^2 = z - q$  ( $U$  simply connected).

$g(z) = g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow g$  is injective.

$g(z) = -g(\zeta) \Rightarrow (g(z))^2 = (g(\zeta))^2 \Rightarrow z = \zeta \Rightarrow$  contradiction ( $g$  is never zero).  
 $\Rightarrow g(U) \cap (-g(U)) = \emptyset$  where  $-g(U) = \{z \in \mathbb{C} : -z \in g(U)\}$ .

$g(U)$  is open  $\Rightarrow -g(U)$  is open  $\Rightarrow \exists \Delta_r(\xi) \subset \mathbb{C} \setminus g(U)$

$\Rightarrow z \mapsto \frac{r}{g(z) - \xi}$  takes  $U$  to  $\mathbb{D}$ .

Compose with an automorphism of  $\mathbb{D}$  to make  $p$  go to 0  $\Rightarrow \mathcal{F}$  is nonempty.

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto.

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.



Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p)$$

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p)$$

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p) = 2g(p)\varphi'_{-g(p)}(h(p))h'(p)$$

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p) = 2g(p)\varphi'_{-g(p)}(h(p))h'(p) = 2g(p)(1 - |g(p)|^2)h'(p).$$



Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p) = 2g(p)\varphi'_{-g(p)}(h(p))h'(p) = 2g(p)(1 - |g(p)|^2)h'(p).$$

$$|f'(p)| = \frac{2|g(p)|(1 - |g(p)|^2)}{1 - |q|^2}|h'(p)|$$

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p) = 2g(p)\varphi'_{-g(p)}(h(p))h'(p) = 2g(p)(1 - |g(p)|^2)h'(p).$$

$$|f'(p)| = \frac{2|g(p)|(1 - |g(p)|^2)}{1 - |q|^2}|h'(p)| = \frac{2\sqrt{|q|}}{1 + |q|}|h'(p)| \quad \text{as } (g(p))^2 = -q.$$

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p) = 2g(p)\varphi'_{-g(p)}(h(p))h'(p) = 2g(p)(1 - |g(p)|^2)h'(p).$$

$$|f'(p)| = \frac{2|g(p)|(1 - |g(p)|^2)}{1 - |q|^2}|h'(p)| = \frac{2\sqrt{|q|}}{1 + |q|}|h'(p)| \quad \text{as } (g(p))^2 = -q.$$

$$|q| < 1 \Rightarrow \frac{2\sqrt{|q|}}{1 + |q|} < 1$$

Suppose  $f: U \rightarrow \mathbb{D}$  is in  $\mathcal{F}$  but not onto. Suppose  $q \in \mathbb{D} \setminus f(U)$ .

Let  $\varphi_q(z) = \frac{z - q}{1 - \bar{q}z}$ . Note  $\varphi_q \in \text{Aut}(\mathbb{D})$ ,  $\varphi_q(q) = 0$ , and  $\varphi_q \circ f$  nonzero.

$\exists g: U \rightarrow \mathbb{C}$  s.t.  $(g(z))^2 = \varphi_q(f(z))$ .  $g(U) \subset \mathbb{D}$ .  $g(z) = g(\zeta) \Rightarrow \varphi_q(f(z)) = \varphi_q(f(\zeta))$ .  
 $\varphi_q \circ f$  injective  $\Rightarrow g$  injective.

Define  $h = \varphi_{g(p)} \circ g$ .  $h(p) = 0 \Rightarrow h \in \mathcal{F}$ . Also  $g = \varphi_{-g(p)} \circ h$ .

Differentiate  $\varphi_q \circ f = g^2$  at  $p$  (recall  $\varphi'_a(0) = 1 - |a|^2$ ):

$$(1 - |q|^2)f'(p) = \varphi'_q(f(p))f'(p) = 2g(p)g'(p) = 2g(p)\varphi'_{-g(p)}(h(p))h'(p) = 2g(p)(1 - |g(p)|^2)h'(p).$$

$$|f'(p)| = \frac{2|g(p)|(1 - |g(p)|^2)}{1 - |q|^2}|h'(p)| = \frac{2\sqrt{|q|}}{1 + |q|}|h'(p)| \quad \text{as } (g(p))^2 = -q.$$

$$|q| < 1 \Rightarrow \frac{2\sqrt{|q|}}{1 + |q|} < 1 \Rightarrow |f'(p)| < |h'(p)|.$$

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence,

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.



Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.

By taking limits:  $|f'(p)| > 0$  ( $f$  not constant),

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.

By taking limits:  $|f'(p)| > 0$  ( $f$  not constant),  $f(p) = 0$ ,

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.

By taking limits:  $|f'(p)| > 0$  ( $f$  not constant),  $f(p) = 0$ ,  $|f(z)| \leq 1$  for all  $z \in U$ .

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.

By taking limits:  $|f'(p)| > 0$  ( $f$  not constant),  $f(p) = 0$ ,  $|f(z)| \leq 1$  for all  $z \in U$ .  
Open mapping theorem  $\Rightarrow |f(z)| < 1$  for all  $z \in U$ .

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.

By taking limits:  $|f'(p)| > 0$  ( $f$  not constant),  $f(p) = 0$ ,  $|f(z)| \leq 1$  for all  $z \in U$ .  
Open mapping theorem  $\Rightarrow |f(z)| < 1$  for all  $z \in U$ .

$f$  must be onto, otherwise there would be an  $h \in \mathcal{F}$  with  $|f'(p)| < |h'(p)|$ .

Construct a sequence  $\{f_n\}$  in  $\mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} |f'_n(p)| = \sup_{f \in \mathcal{F}} |f'(p)|$$

Montel says ( $\mathcal{F}$  is uniformly bounded), there exists a convergent subsequence, WLOG  $\{f_n\}$  converges to  $f$ .

By the corollary to Hurwitz,  $f$  is injective or constant.

By taking limits:  $|f'(p)| > 0$  ( $f$  not constant),  $f(p) = 0$ ,  $|f(z)| \leq 1$  for all  $z \in U$ .  
Open mapping theorem  $\Rightarrow |f(z)| < 1$  for all  $z \in U$ .

$f$  must be onto, otherwise there would be an  $h \in \mathcal{F}$  with  $|f'(p)| < |h'(p)|$ .

Uniqueness left as an exercise.



**Remark:** An explicit map is useful, e.g., in differential equations.

**Remark:** An explicit map is useful, e.g., in differential equations.  
The theorem doesn't answer how a map is constructed.



**Remark:** An explicit map is useful, e.g., in differential equations.  
The theorem doesn't answer how a map is constructed.  
There is lots of literature on constructing the map.

**Remark:** An explicit map is useful, e.g., in differential equations.

The theorem doesn't answer how a map is constructed.

There is lots of literature on constructing the map.

E.g., if  $U$  is a polygon, there is an explicit formula: the Schwarz–Christoffel mapping.

**Remark:** An explicit map is useful, e.g., in differential equations.

The theorem doesn't answer how a map is constructed.

There is lots of literature on constructing the map.

E.g., if  $U$  is a polygon, there is an explicit formula: the Schwarz–Christoffel mapping.

**Remark:** The theorem doesn't answer how regular the map is up to the boundary.

**Remark:** An explicit map is useful, e.g., in differential equations.

The theorem doesn't answer how a map is constructed.

There is lots of literature on constructing the map.

E.g., if  $U$  is a polygon, there is an explicit formula: the Schwarz–Christoffel mapping.

**Remark:** The theorem doesn't answer how regular the map is up to the boundary.

The nicer the boundary, the nicer the map will be.

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a simply connected domain. Show that for every two points  $z, w \in U$ , there exists an automorphism  $\psi \in \text{Aut}(U)$  such that  $\psi(z) = w$ .

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a simply connected domain. Show that for every two points  $z, w \in U$ , there exists an automorphism  $\psi \in \text{Aut}(U)$  such that  $\psi(z) = w$ .

**Exercise:**

a) Suppose  $U \subset \mathbb{C}$  is a simply connected domain,  $U \neq \mathbb{C}$ ,  $p, q \in U$  are distinct points, and  $f: U \rightarrow U$  is holomorphic such that  $f(p) = p$  and  $f(q) = q$ . Prove that  $f$  is the identity.

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a simply connected domain. Show that for every two points  $z, w \in U$ , there exists an automorphism  $\psi \in \text{Aut}(U)$  such that  $\psi(z) = w$ .

**Exercise:**

- a) Suppose  $U \subset \mathbb{C}$  is a simply connected domain,  $U \neq \mathbb{C}$ ,  $p, q \in U$  are distinct points, and  $f: U \rightarrow U$  is holomorphic such that  $f(p) = p$  and  $f(q) = q$ . Prove that  $f$  is the identity.
- b) Find a counterexample if  $U = \mathbb{C}$ .

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a simply connected domain. Show that for every two points  $z, w \in U$ , there exists an automorphism  $\psi \in \text{Aut}(U)$  such that  $\psi(z) = w$ .

**Exercise:**

- a) Suppose  $U \subset \mathbb{C}$  is a simply connected domain,  $U \neq \mathbb{C}$ ,  $p, q \in U$  are distinct points, and  $f: U \rightarrow U$  is holomorphic such that  $f(p) = p$  and  $f(q) = q$ . Prove that  $f$  is the identity.
- b) Find a counterexample if  $U = \mathbb{C}$ .

**Exercise:** Show that  $\mathbb{D} \setminus \{0\}$  and the annulus  $\text{ann}(0; 1, 2)$  are not biholomorphic.



**Exercise:** Suppose  $U \subset \mathbb{C}$  is a simply connected domain. Show that for every two points  $z, w \in U$ , there exists an automorphism  $\psi \in \text{Aut}(U)$  such that  $\psi(z) = w$ .

**Exercise:**

- a) Suppose  $U \subset \mathbb{C}$  is a simply connected domain,  $U \neq \mathbb{C}$ ,  $p, q \in U$  are distinct points, and  $f: U \rightarrow U$  is holomorphic such that  $f(p) = p$  and  $f(q) = q$ . Prove that  $f$  is the identity.
- b) Find a counterexample if  $U = \mathbb{C}$ .

**Exercise:** Show that  $\mathbb{D} \setminus \{0\}$  and the annulus  $\text{ann}(0; 1, 2)$  are not biholomorphic.

**Exercise:** Suppose  $f: \mathbb{C} \rightarrow \mathbb{C}$  is entire holomorphic and injective, prove that  $f$  is onto.