

Cultivating Complex Analysis: Holomorphic functions are analytic (3.3.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

A consequence of Cauchy's formula is that holomorphic functions are analytic.

A consequence of Cauchy's formula is that holomorphic functions are analytic.

We already proved that analytic functions are holomorphic, let's prove the converse.

A consequence of Cauchy's formula is that holomorphic functions are analytic.

We already proved that analytic functions are holomorphic, let's prove the converse.

Theorem

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ holomorphic, $p \in U$, and $\Delta_R(p) \subset U$.

A consequence of Cauchy's formula is that holomorphic functions are analytic.

We already proved that analytic functions are holomorphic, let's prove the converse.

Theorem

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ holomorphic, $p \in U$, and $\Delta_R(p) \subset U$. Then there exists a power series $\sum c_n(z - p)^n$ such that for all $z \in \Delta_R(p)$,

$$f(z) = \sum_{n=0}^{\infty} c_n(z - p)^n.$$

A consequence of Cauchy's formula is that holomorphic functions are analytic.

We already proved that analytic functions are holomorphic, let's prove the converse.

Theorem

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ holomorphic, $p \in U$, and $\Delta_R(p) \subset U$. Then there exists a power series $\sum c_n(z-p)^n$ such that for all $z \in \Delta_R(p)$,

$$f(z) = \sum_{n=0}^{\infty} c_n(z-p)^n.$$

Moreover,

$$c_n = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{(z-p)^{n+1}} dz,$$

where γ is any circle of radius r , $0 < r < R$, centered at p oriented counterclockwise.

Proof: Fix an r such that $0 < r < R$.

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$.

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$. For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{r} < 1.$$

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$. For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{r} < 1.$$

So

$$\sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n = \frac{1}{1 - \frac{z-p}{\zeta-p}} = \frac{\zeta-p}{\zeta-z}.$$

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$. For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{r} < 1.$$

So

$$\sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n = \frac{1}{1 - \frac{z-p}{\zeta-p}} = \frac{\zeta-p}{\zeta-z}.$$

$$f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-z} d\zeta$$

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$. For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{r} < 1.$$

So

$$\sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n = \frac{1}{1 - \frac{z-p}{\zeta-p}} = \frac{\zeta-p}{\zeta-z}.$$

$$f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-p} \frac{\zeta-p}{\zeta-z} d\zeta$$

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$. For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{r} < 1.$$

So

$$\sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n = \frac{1}{1 - \frac{z-p}{\zeta-p}} = \frac{\zeta-p}{\zeta-z}.$$

$$\begin{aligned} f(z) &= \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-p} \frac{\zeta-p}{\zeta-z} d\zeta \\ &= \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-p} \sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n d\zeta \end{aligned}$$

Proof: Fix an r such that $0 < r < R$. Thus $\overline{\Delta_r(p)} \subset U$.

Fix a $z \in \Delta_r(p)$. For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{r} < 1.$$

So

$$\sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n = \frac{1}{1 - \frac{z-p}{\zeta-p}} = \frac{\zeta-p}{\zeta-z}.$$

$$\begin{aligned} f(z) &= \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-p} \frac{\zeta-p}{\zeta-z} d\zeta \\ &= \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta-p} \sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p} \right)^n d\zeta = \sum_{n=0}^{\infty} \underbrace{\left(\frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta-p)^{n+1}} d\zeta \right)}_{c_n} (z-p)^n. \end{aligned}$$

The last equality held because the sum converges uniformly in $\zeta \in \partial\Delta_r(p)$ (we'll justify that on the next slide).

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta - p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set),

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta - p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set), then

$$\left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right| \leq M \left(\frac{|z - p|}{r} \right)^n, \quad \text{and} \quad \frac{|z - p|}{r} < 1.$$

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta - p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set), then

$$\left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right| \leq M \left(\frac{|z - p|}{r} \right)^n, \quad \text{and} \quad \frac{|z - p|}{r} < 1.$$

Thus, $\sum \left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right|$ converges uniformly in $\zeta \in \partial\Delta_r(p)$,

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta-p} \left(\frac{z-p}{\zeta-p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta-p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set), then

$$\left| \frac{f(\zeta)}{\zeta-p} \left(\frac{z-p}{\zeta-p} \right)^n \right| \leq M \left(\frac{|z-p|}{r} \right)^n, \quad \text{and} \quad \frac{|z-p|}{r} < 1.$$

Thus, $\sum \left| \frac{f(\zeta)}{\zeta-p} \left(\frac{z-p}{\zeta-p} \right)^n \right|$ converges uniformly in $\zeta \in \partial\Delta_r(p)$,

and so $\sum \frac{f(\zeta)}{\zeta-p} \left(\frac{z-p}{\zeta-p} \right)^n$ converges uniformly absolutely (and hence uniformly).

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta - p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set), then

$$\left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right| \leq M \left(\frac{|z - p|}{r} \right)^n, \quad \text{and} \quad \frac{|z - p|}{r} < 1.$$

Thus, $\sum \left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right|$ converges uniformly in $\zeta \in \partial\Delta_r(p)$,

and so $\sum \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly absolutely (and hence uniformly).

We found a power series converging to $f(z)$ for all $z \in \Delta_r(p)$.

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta - p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set), then

$$\left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right| \leq M \left(\frac{|z - p|}{r} \right)^n, \quad \text{and} \quad \frac{|z - p|}{r} < 1.$$

Thus, $\sum \left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right|$ converges uniformly in $\zeta \in \partial\Delta_r(p)$,

and so $\sum \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly absolutely (and hence uniformly).

We found a power series converging to $f(z)$ for all $z \in \Delta_r(p)$.

By uniqueness of the power series, the c_n is independent of r .

We claim $\sum_{n=0}^{\infty} \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly in $\zeta \in \partial\Delta_r(p)$.

z is fixed and if M is the supremum of $\left| \frac{f(\zeta)}{\zeta - p} \right| = \frac{|f(\zeta)|}{r}$ on $\partial\Delta_r(p)$ (a compact set), then

$$\left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right| \leq M \left(\frac{|z - p|}{r} \right)^n, \quad \text{and} \quad \frac{|z - p|}{r} < 1.$$

Thus, $\sum \left| \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n \right|$ converges uniformly in $\zeta \in \partial\Delta_r(p)$,

and so $\sum \frac{f(\zeta)}{\zeta - p} \left(\frac{z - p}{\zeta - p} \right)^n$ converges uniformly absolutely (and hence uniformly).

We found a power series converging to $f(z)$ for all $z \in \Delta_r(p)$.

By uniqueness of the power series, the c_n is independent of r .

We get the same series for every r and it converges in $\Delta_R(p)$.



The key is writing the *Cauchy kernel* as

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - p} \frac{\zeta - p}{\zeta - z}$$

The key is writing the *Cauchy kernel* as

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - p} \frac{\zeta - p}{\zeta - z}$$

and using the geometric series.

A common technique: the integral of a function against a kernel gains properties of the kernel.


The key is writing the *Cauchy kernel* as

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - p} \frac{\zeta - p}{\zeta - z}$$

and using the geometric series.

A common technique: the integral of a function against a kernel gains properties of the kernel.

Besides analyticity, we also proved that the radius of convergence is at least R , where R is the maximum R such that $\Delta_R(p) \subset U$.



The key is writing the *Cauchy kernel* as


$$\frac{1}{\zeta - z} = \frac{1}{\zeta - p} \frac{\zeta - p}{\zeta - z}$$

and using the geometric series.

A common technique: the integral of a function against a kernel gains properties of the kernel.

Besides analyticity, we also proved that the radius of convergence is at least R , where R is the maximum R such that $\Delta_R(p) \subset U$.

Note that the radius of convergence gives bounds on the derivatives!



The key is writing the *Cauchy kernel* as

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - p} \frac{\zeta - p}{\zeta - z}$$


and using the geometric series.

A common technique: the integral of a function against a kernel gains properties of the kernel.

Besides analyticity, we also proved that the radius of convergence is at least R , where R is the maximum R such that $\Delta_R(p) \subset U$.

Note that the radius of convergence gives bounds on the derivatives!

So we know about the size of the derivatives at p just from knowing how far away from p is f still holomorphic.



The key is writing the *Cauchy kernel* as

$$\frac{1}{\zeta - z} = \frac{1}{\zeta - p} \frac{\zeta - p}{\zeta - z}$$


and using the geometric series.

A common technique: the integral of a function against a kernel gains properties of the kernel.

Besides analyticity, we also proved that the radius of convergence is at least R , where R is the maximum R such that $\Delta_R(p) \subset U$.

Note that the radius of convergence gives bounds on the derivatives!

So we know about the size of the derivatives at p just from knowing how far away from p is f still holomorphic.



Remark: Nothing like this is true for real-analytic functions such as $\varphi(x) = \frac{1}{1+x^2}$ whose radius of convergence at $x = 0$ is 1, but $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ is (real) analytic everywhere.

Let us restate the main conclusion:

Corollary

Let $U \subset \mathbb{C}$ be an open set. A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is analytic.

Let us restate the main conclusion:

Corollary

Let $U \subset \mathbb{C}$ be an open set. A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is analytic.

Results we proved for analytic functions are true for holomorphic functions and vice versa.

Let us restate the main conclusion:

Corollary

Let $U \subset \mathbb{C}$ be an open set. A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is analytic.

Results we proved for analytic functions are true for holomorphic functions and vice versa.

E.g., it is easy to show that the composition of holomorphic functions is holomorphic via the chain rule.

Let us restate the main conclusion:

Corollary

Let $U \subset \mathbb{C}$ be an open set. A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is analytic.

Results we proved for analytic functions are true for holomorphic functions and vice versa.

E.g., it is easy to show that the composition of holomorphic functions is holomorphic via the chain rule.

It is much harder to show this for analytic functions by directly manipulating power series.

Let us restate the main conclusion:

Corollary

Let $U \subset \mathbb{C}$ be an open set. A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is analytic.

Results we proved for analytic functions are true for holomorphic functions and vice versa.

E.g., it is easy to show that the composition of holomorphic functions is holomorphic via the chain rule.

It is much harder to show this for analytic functions by directly manipulating power series.

But we don't have to, analytic functions are holomorphic.

Let us restate the main conclusion:

Corollary

Let $U \subset \mathbb{C}$ be an open set. A function $f: U \rightarrow \mathbb{C}$ is holomorphic if and only if f is analytic.

Results we proved for analytic functions are true for holomorphic functions and vice versa.

E.g., it is easy to show that the composition of holomorphic functions is holomorphic via the chain rule.

It is much harder to show this for analytic functions by directly manipulating power series.

But we don't have to, analytic functions are holomorphic.

We have also finally proved the following:

A convergent power series defines an analytic function.