

Cultivating Complex Analysis: Primitives, cycles, and Cauchy for derivatives (3.2.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{C}$ a function. A holomorphic $F: U \rightarrow \mathbb{C}$ with $f = F'$ is called a (holomorphic) *primitive* (or an *antiderivative*) of f .

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{C}$ a function. A holomorphic $F: U \rightarrow \mathbb{C}$ with $f = F'$ is called a (holomorphic) *primitive* (or an *antiderivative*) of f .

Primitives do not always exist, but if they do, then they are unique up to a constant.

Proposition

Suppose $U \subset \mathbb{C}$ is a domain, and $F: U \rightarrow \mathbb{C}$ and $G: U \rightarrow \mathbb{C}$ are holomorphic such that $F' = G'$. Then $F(z) = G(z) + C$ for some $C \in \mathbb{C}$.

Definition

Let $U \subset \mathbb{C}$ be open and $f: U \rightarrow \mathbb{C}$ a function. A holomorphic $F: U \rightarrow \mathbb{C}$ with $f = F'$ is called a (holomorphic) *primitive* (or an *antiderivative*) of f .

Primitives do not always exist, but if they do, then they are unique up to a constant.

Proposition

Suppose $U \subset \mathbb{C}$ is a domain, and $F: U \rightarrow \mathbb{C}$ and $G: U \rightarrow \mathbb{C}$ are holomorphic such that $F' = G'$. Then $F(z) = G(z) + C$ for some $C \in \mathbb{C}$.

Proof is an exercise.

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$).

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz$$

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt$$

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{d}{dt} (F(\gamma(t))) dt$$

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{d}{dt} (F(\gamma(t))) dt = F(\gamma(b)) - F(\gamma(a)).$$

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{d}{dt} (F(\gamma(t))) dt = F(\gamma(b)) - F(\gamma(a)).$$

We used the chain rule for composing real differentiable and a holomorphic function,

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{d}{dt} (F(\gamma(t))) dt = F(\gamma(b)) - F(\gamma(a)).$$

We used the chain rule for composing real differentiable and a holomorphic function, and the standard (real) fundamental theorem of calculus (applied to the real and imaginary parts). □

Theorem (Fundamental theorem of calculus for line integrals)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$ (so $F' = f$). Let $\gamma: [a, b] \rightarrow U$ be a piecewise- C^1 path. Then

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

Proof:

$$\int_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t)) \gamma'(t) dt = \int_a^b \frac{d}{dt} (F(\gamma(t))) dt = F(\gamma(b)) - F(\gamma(a)).$$

We used the chain rule for composing real differentiable and a holomorphic function, and the standard (real) fundamental theorem of calculus (applied to the real and imaginary parts). □

Remark: The hypothesis that $f = F'$ is continuous is extraneous (we will prove later that f is better than continuous, it is holomorphic).

Definition

A chain Γ that is equivalent to $a_1\gamma_1 + \cdots + a_n\gamma_n$, where $\gamma_1, \dots, \gamma_n$ are closed piecewise- C^1 paths and $a_1, \dots, a_n \in \mathbb{Z}$, is called a *cycle*.

Definition

A chain Γ that is equivalent to $a_1\gamma_1 + \cdots + a_n\gamma_n$, where $\gamma_1, \dots, \gamma_n$ are closed piecewise- C^1 paths and $a_1, \dots, a_n \in \mathbb{Z}$, is called a *cycle*.

A cycle need not be a sum of closed paths, just equivalent to it.

Definition

A chain Γ that is equivalent to $a_1\gamma_1 + \cdots + a_n\gamma_n$, where $\gamma_1, \dots, \gamma_n$ are closed piecewise- C^1 paths and $a_1, \dots, a_n \in \mathbb{Z}$, is called a *cycle*.

A cycle need not be a sum of closed paths, just equivalent to it.

E.g., the unit square path we saw before is a cycle and could be written as the chain $[0, 1] + [1, 1 + i] + [1 + i, i] + [i, 0]$.

Definition

A chain Γ that is equivalent to $a_1\gamma_1 + \cdots + a_n\gamma_n$, where $\gamma_1, \dots, \gamma_n$ are closed piecewise- C^1 paths and $a_1, \dots, a_n \in \mathbb{Z}$, is called a *cycle*.

A cycle need not be a sum of closed paths, just equivalent to it.

E.g., the unit square path we saw before is a cycle and could be written as the chain $[0, 1] + [1, 1 + i] + [1 + i, i] + [i, 0]$.

The fundamental theorem has an immediate corollary, noting that for a closed path $\gamma: [a, b] \rightarrow \mathbb{C}$, $\gamma(a) = \gamma(b)$.

Definition

A chain Γ that is equivalent to $a_1\gamma_1 + \cdots + a_n\gamma_n$, where $\gamma_1, \dots, \gamma_n$ are closed piecewise- C^1 paths and $a_1, \dots, a_n \in \mathbb{Z}$, is called a *cycle*.

A cycle need not be a sum of closed paths, just equivalent to it.

E.g., the unit square path we saw before is a cycle and could be written as the chain $[0, 1] + [1, 1 + i] + [1 + i, i] + [i, 0]$.

The fundamental theorem has an immediate corollary, noting that for a closed path $\gamma: [a, b] \rightarrow \mathbb{C}$, $\gamma(a) = \gamma(b)$.

Corollary (Cauchy's theorem for derivatives)

Suppose $U \subset \mathbb{C}$ is open and $f: U \rightarrow \mathbb{C}$ is continuous with a primitive $F: U \rightarrow \mathbb{C}$. Let Γ be a cycle in U . Then

$$\int_{\Gamma} f(z) dz = 0.$$

We will prove several versions of Cauchy, though usually we will put restrictions on U or the Γ rather than the function.

We will prove several versions of Cauchy, though usually we will put restrictions on U or the Γ rather than the function.

“Cauchy’s theorem” (in general) is a sort of an independence of path result used to define a primitive, so either the function has a primitive in U or we have to restrict which Γ s are allowed.

We will prove several versions of Cauchy, though usually we will put restrictions on U or the Γ rather than the function.

“Cauchy’s theorem” (in general) is a sort of an independence of path result used to define a primitive, so either the function has a primitive in U or we have to restrict which Γ s are allowed.

We can now prove a very neat version of the theorem (exercise):

Corollary (Cauchy’s theorem for polynomials)

Suppose $P(z)$ is a polynomial and Γ is a cycle (in \mathbb{C}). Then

$$\int_{\Gamma} P(z) dz = 0.$$