

# Cultivating Complex Analysis: Rouché's theorem (5.4.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

For meromorphic functions the difference of zeros and poles is the same.

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

For meromorphic functions the difference of zeros and poles is the same.

**Example:**  $z^2$  and  $(z - \epsilon)(z + \epsilon)$  are close on  $\partial\mathbb{D}$ .

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

For meromorphic functions the difference of zeros and poles is the same.

**Example:**  $z^2$  and  $(z - \epsilon)(z + \epsilon)$  are close on  $\partial\mathbb{D}$ .

**Example:** 1 and  $\frac{z-\epsilon}{z+\epsilon}$  are close on  $\partial\mathbb{D}$ .

## Theorem (Rouché)

*Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , and  $n(\Gamma; z)$  is either 0 or 1 for all  $z \notin \Gamma$ .*

## Theorem (Rouché)

*Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , and  $n(\Gamma; z)$  is either 0 or 1 for all  $z \notin \Gamma$ . Suppose that  $f: U \rightarrow \mathbb{C}_\infty$  and  $g: U \rightarrow \mathbb{C}_\infty$  are meromorphic functions with no zeros or poles on  $\Gamma$  such that*

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

## Theorem (Rouché)

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , and  $n(\Gamma; z)$  is either 0 or 1 for all  $z \notin \Gamma$ . Suppose that  $f: U \rightarrow \mathbb{C}_\infty$  and  $g: U \rightarrow \mathbb{C}_\infty$  are meromorphic functions with no zeros or poles on  $\Gamma$  such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Let  $V = \{z \in U : n(\Gamma; z) = 1\}$ . Let  $N_f, N_g$  be the number of zeros in  $V$  and  $P_f, P_g$  the number of poles in  $V$  (both counting multiplicity) of  $f$  and  $g$  respectively.



## Theorem (Rouché)

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , and  $n(\Gamma; z)$  is either 0 or 1 for all  $z \notin \Gamma$ . Suppose that  $f: U \rightarrow \mathbb{C}_\infty$  and  $g: U \rightarrow \mathbb{C}_\infty$  are meromorphic functions with no zeros or poles on  $\Gamma$  such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Let  $V = \{z \in U : n(\Gamma; z) = 1\}$ . Let  $N_f, N_g$  be the number of zeros in  $V$  and  $P_f, P_g$  the number of poles in  $V$  (both counting multiplicity) of  $f$  and  $g$  respectively. Then

$$N_f - P_f = N_g - P_g.$$

## Theorem (Rouché)

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , and  $n(\Gamma; z)$  is either 0 or 1 for all  $z \notin \Gamma$ . Suppose that  $f: U \rightarrow \mathbb{C}_\infty$  and  $g: U \rightarrow \mathbb{C}_\infty$  are meromorphic functions with no zeros or poles on  $\Gamma$  such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Let  $V = \{z \in U : n(\Gamma; z) = 1\}$ . Let  $N_f, N_g$  be the number of zeros in  $V$  and  $P_f, P_g$  the number of poles in  $V$  (both counting multiplicity) of  $f$  and  $g$  respectively. Then

$$N_f - P_f = N_g - P_g.$$

## Corollary (Rouché)

Let  $U, \Gamma$  and  $V$  be as in the theorem. Suppose  $f: U \rightarrow \mathbb{C}$  and  $g: U \rightarrow \mathbb{C}$  are holomorphic such that  $|f(z) - g(z)| < |f(z)| + |g(z)|$  for all  $z \in \Gamma$ . Then  $f$  and  $g$  have the same number of zeros (counting multiplicity) in  $V$ .


Classical statement of the theorem uses the weaker inequality

$$|f(z) - g(z)| < |f(z)|$$

Classical statement of the theorem uses the weaker inequality

$$|f(z) - g(z)| < |f(z)|$$

It has a nice geometric interpretation:



**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

Let  $\text{Log}$  be the principal branch of  $\log$  on  $\mathbb{C} \setminus (-\infty, 0]$ .

**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

Let  $\text{Log}$  be the principal branch of  $\log$  on  $\mathbb{C} \setminus (-\infty, 0]$ .

The function  $\frac{\varphi'}{\varphi}$  has an antiderivative  $\text{Log} \circ \varphi$  on a neighborhood of  $\Gamma$ .



**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

Let  $\text{Log}$  be the principal branch of  $\log$  on  $\mathbb{C} \setminus (-\infty, 0]$ .

The function  $\frac{\varphi'}{\varphi}$  has an antiderivative  $\text{Log} \circ \varphi$  on a neighborhood of  $\Gamma$ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$0 = \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz$$

**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

Let  $\text{Log}$  be the principal branch of  $\log$  on  $\mathbb{C} \setminus (-\infty, 0]$ .

The function  $\frac{\varphi'}{\varphi}$  has an antiderivative  $\text{Log} \circ \varphi$  on a neighborhood of  $\Gamma$ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$0 = \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz = \frac{1}{2\pi i} \int_{\Gamma} \left( \frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \right) dz$$

**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

Let  $\text{Log}$  be the principal branch of  $\log$  on  $\mathbb{C} \setminus (-\infty, 0]$ .

The function  $\frac{\varphi'}{\varphi}$  has an antiderivative  $\text{Log} \circ \varphi$  on a neighborhood of  $\Gamma$ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz = \frac{1}{2\pi i} \int_{\Gamma} \left( \frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \right) dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz - \frac{1}{2\pi i} \int_{\Gamma} \frac{g'(z)}{g(z)} dz \end{aligned}$$

**Proof:**

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So  $\varphi(z) = \frac{f(z)}{g(z)}$  is never negative (and it is never zero) on  $\Gamma$  (hence on a neighborhood).

Let  $\text{Log}$  be the principal branch of  $\log$  on  $\mathbb{C} \setminus (-\infty, 0]$ .

The function  $\frac{\varphi'}{\varphi}$  has an antiderivative  $\text{Log} \circ \varphi$  on a neighborhood of  $\Gamma$ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz = \frac{1}{2\pi i} \int_{\Gamma} \left( \frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \right) dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz - \frac{1}{2\pi i} \int_{\Gamma} \frac{g'(z)}{g(z)} dz = (N_f - P_f) - (N_g - P_g). \end{aligned}$$

□

The classical hypothesis  $|f(z) - g(z)| < |f(z)|$  is often sufficient.

The classical hypothesis  $|f(z) - g(z)| < |f(z)|$  is often sufficient.

**Example:** Consider  $P(z) = z^n + 1$ . Let us use Rouché to show that all the zeros are on  $\partial\mathbb{D}$ .

The classical hypothesis  $|f(z) - g(z)| < |f(z)|$  is often sufficient.

**Example:** Consider  $P(z) = z^n + 1$ . Let us use Rouché to show that all the zeros are on  $\partial\mathbb{D}$ .

On  $\partial\Delta_{1-\epsilon}(0)$ ,

$$|P(z) - 1| = |z|^n < 1 = |1|.$$

The classical hypothesis  $|f(z) - g(z)| < |f(z)|$  is often sufficient.

**Example:** Consider  $P(z) = z^n + 1$ . Let us use Rouché to show that all the zeros are on  $\partial\mathbb{D}$ .

On  $\partial\Delta_{1-\epsilon}(0)$ ,

$$|P(z) - 1| = |z|^n < 1 = |1|.$$

By Rouché  $P(z)$  and 1 have the same number of zeros in  $\Delta_{1-\epsilon}(0)$ .



The classical hypothesis  $|f(z) - g(z)| < |f(z)|$  is often sufficient.

**Example:** Consider  $P(z) = z^n + 1$ . Let us use Rouché to show that all the zeros are on  $\partial\mathbb{D}$ .

On  $\partial\Delta_{1-\epsilon}(0)$ ,

$$|P(z) - 1| = |z|^n < 1 = |1|.$$

By Rouché  $P(z)$  and 1 have the same number of zeros in  $\Delta_{1-\epsilon}(0)$ .

On  $\partial\Delta_{1+\epsilon}(0)$ ,

$$|P(z) - z^n| = 1 < |z^n|.$$

By Rouché,  $P(z)$  and  $z^n$  have the same number of zeros in  $\Delta_{1+\epsilon}(0)$ .

**Example:** Consider  $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$ .

**Example:** Consider  $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$ . On  $\partial\mathbb{D}$ ,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

**Example:** Consider  $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$ . On  $\partial\mathbb{D}$ ,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$  has zeros at  $\pm\sqrt{24}i$  (outside  $\mathbb{D}$ ) and two zeros at the origin (inside  $\mathbb{D}$ ). So  $P(z)$  also has two zeros in  $\mathbb{D}$ .

**Example:** Consider  $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$ . On  $\partial\mathbb{D}$ ,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$  has zeros at  $\pm\sqrt{24}i$  (outside  $\mathbb{D}$ ) and two zeros at the origin (inside  $\mathbb{D}$ ). So  $P(z)$  also has two zeros in  $\mathbb{D}$ .

If  $|z| = 46 + \epsilon$ , then

$$|P(z) - z^4| = |12z^3 + 24z^2 + 4z + 6| \leq 46|z|^3 < |z|^4 = |z^4|.$$

**Example:** Consider  $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$ . On  $\partial\mathbb{D}$ ,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$  has zeros at  $\pm\sqrt{24}i$  (outside  $\mathbb{D}$ ) and two zeros at the origin (inside  $\mathbb{D}$ ). So  $P(z)$  also has two zeros in  $\mathbb{D}$ .

If  $|z| = 46 + \epsilon$ , then

$$|P(z) - z^4| = |12z^3 + 24z^2 + 4z + 6| \leq 46|z|^3 < |z|^4 = |z^4|.$$

So all four zeros satisfy  $|z| < 46 + \epsilon$ , that is,  $|z| \leq 46$ .

**Example:** Consider  $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$ . On  $\partial\mathbb{D}$ ,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$  has zeros at  $\pm\sqrt{24}i$  (outside  $\mathbb{D}$ ) and two zeros at the origin (inside  $\mathbb{D}$ ). So  $P(z)$  also has two zeros in  $\mathbb{D}$ .

If  $|z| = 46 + \epsilon$ , then

$$|P(z) - z^4| = |12z^3 + 24z^2 + 4z + 6| \leq 46|z|^3 < |z|^4 = |z^4|.$$

So all four zeros satisfy  $|z| < 46 + \epsilon$ , that is,  $|z| \leq 46$ .

(Actually the largest zero of  $P$  has modulus less than 10).