

# Cultivating Complex Analysis: Simply connected domains (4.3 part 1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

A few remarks are in order:

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

A few remarks are in order:

**Remark:** This is the “wrong” definition. It happens to work in the setting of domains in  $\mathbb{C}$ . It is wrong for two reasons:

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

A few remarks are in order:

**Remark:** This is the “wrong” definition. It happens to work in the setting of domains in  $\mathbb{C}$ . It is wrong for two reasons:

- 1) It is in terms of homology and not homotopy (an optional section in the book).



## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

A few remarks are in order:

**Remark:** This is the “wrong” definition. It happens to work in the setting of domains in  $\mathbb{C}$ . It is wrong for two reasons:

- 1) It is in terms of homology and not homotopy (an optional section in the book).

One could say *simply connected in the sense of homology* to emphasize.

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

A few remarks are in order:

**Remark:** This is the “wrong” definition. It happens to work in the setting of domains in  $\mathbb{C}$ . It is wrong for two reasons:

- 1) It is in terms of homology and not homotopy (an optional section in the book).

One could say *simply connected in the sense of homology* to emphasize.

- 2) We defined cycles as “piecewise- $C^1$ ” instead of “continuous.”

## Definition

A domain  $U \subset \mathbb{C}$  is *simply connected* if every cycle in  $U$  is homologous to zero in  $U$ .

That is,  $U$  is simply connected if  $n(\Gamma; p) = 0$  for every cycle  $\Gamma$  in  $U$  and every  $p \in \mathbb{C} \setminus U$ .

### Exercise:

- a) star-like domains (e.g.,  $\mathbb{C}$ ,  $\mathbb{D}$ , and  $\mathbb{H}$ ) in  $\mathbb{C}$  are simply connected.
- b)  $\mathbb{C} \setminus \{0\}$  is not simply connected.

A few remarks are in order:

**Remark:** This is the “wrong” definition. It happens to work in the setting of domains in  $\mathbb{C}$ . It is wrong for two reasons:

- 1) It is in terms of homology and not homotopy (an optional section in the book).

One could say *simply connected in the sense of homology* to emphasize.

- 2) We defined cycles as “piecewise- $C^1$ ” instead of “continuous.”

**Remark:** Can a disconnected set be simply connected? We remain neutral on this.

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

*Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic.*

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

*Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then*

$$\int_{\Gamma} f(z) dz = 0.$$

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Since  $U$  is simply connected,  $n(\Gamma; p) = 0$  for every  $p \in \mathbb{C} \setminus U$ , so Cauchy applies.  $\square$

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Since  $U$  is simply connected,  $n(\Gamma; p) = 0$  for every  $p \in \mathbb{C} \setminus U$ , so Cauchy applies.  $\square$

If we have Cauchy's theorem we expect primitives:

### Theorem

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic.

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Since  $U$  is simply connected,  $n(\Gamma; p) = 0$  for every  $p \in \mathbb{C} \setminus U$ , so Cauchy applies.  $\square$

If we have Cauchy's theorem we expect primitives:

### Theorem

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. Then  $f$  has a primitive in  $U$ .



Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Since  $U$  is simply connected,  $n(\Gamma; p) = 0$  for every  $p \in \mathbb{C} \setminus U$ , so Cauchy applies.  $\square$

If we have Cauchy's theorem we expect primitives:

### Theorem

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. Then  $f$  has a primitive in  $U$ .

**Proof:** Fix  $p \in U$  and note that  $U$  is path connected.

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Since  $U$  is simply connected,  $n(\Gamma; p) = 0$  for every  $p \in \mathbb{C} \setminus U$ , so Cauchy applies.  $\square$

If we have Cauchy's theorem we expect primitives:

### Theorem

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. Then  $f$  has a primitive in  $U$ .

**Proof:** Fix  $p \in U$  and note that  $U$  is path connected.

For every  $z \in U$ , pick some piecewise- $C^1$  path  $\gamma$  from  $p$  to  $z$ , and

Simply connected domains satisfy Cauchy's theorem.

### Theorem (Cauchy's theorem (simply connected version))

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. If  $\Gamma$  is a cycle in  $U$ , then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Since  $U$  is simply connected,  $n(\Gamma; p) = 0$  for every  $p \in \mathbb{C} \setminus U$ , so Cauchy applies.  $\square$

If we have Cauchy's theorem we expect primitives:

### Theorem

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  holomorphic. Then  $f$  has a primitive in  $U$ .

**Proof:** Fix  $p \in U$  and note that  $U$  is path connected.

For every  $z \in U$ , pick some piecewise- $C^1$  path  $\gamma$  from  $p$  to  $z$ , and

Define 
$$F(z) = \int_{\gamma} f(\zeta) d\zeta.$$

If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0$$

If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0 \quad \Rightarrow \quad F(z) = \int_{\gamma} f(\zeta) d\zeta \text{ does not depend on } \gamma.$$

If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0 \quad \Rightarrow \quad F(z) = \int_{\gamma} f(\zeta) d\zeta \text{ does not depend on } \gamma.$$


Consider  $q \in U$ ,  $\gamma$  a path from  $p$  to  $q$ , and  $\Delta_r(q) \subset U$ .

If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0 \quad \Rightarrow \quad F(z) = \int_{\gamma} f(\zeta) d\zeta \text{ does not depend on } \gamma.$$

Consider  $q \in U$ ,  $\gamma$  a path from  $p$  to  $q$ , and  $\Delta_r(q) \subset U$ . For  $z \in \Delta_r(q)$

$$F(z) = \int_{\gamma+[q,z]} f(\zeta) d\zeta$$




If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0 \quad \Rightarrow \quad F(z) = \int_{\gamma} f(\zeta) d\zeta \text{ does not depend on } \gamma.$$

Consider  $q \in U$ ,  $\gamma$  a path from  $p$  to  $q$ , and  $\Delta_r(q) \subset U$ . For  $z \in \Delta_r(q)$

$$\begin{aligned} F(z) &= \int_{\gamma+[q,z]} f(\zeta) d\zeta \\ &= \int_{\gamma} f(\zeta) d\zeta + \int_{[q,z]} f(\zeta) d\zeta. \end{aligned}$$






If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0 \quad \Rightarrow \quad F(z) = \int_{\gamma} f(\zeta) d\zeta \text{ does not depend on } \gamma.$$

Consider  $q \in U$ ,  $\gamma$  a path from  $p$  to  $q$ , and  $\Delta_r(q) \subset U$ . For  $z \in \Delta_r(q)$

$$\begin{aligned} F(z) &= \int_{\gamma+[q,z]} f(\zeta) d\zeta \\ &= \int_{\gamma} f(\zeta) d\zeta + \int_{[q,z]} f(\zeta) d\zeta. \end{aligned}$$

The first term is a constant.



If  $\alpha$  is another path from  $p$  to  $z$ , then by Cauchy

$$\int_{\gamma} f(\zeta) d\zeta - \int_{\alpha} f(\zeta) d\zeta = \int_{\gamma-\alpha} f(\zeta) d\zeta = 0 \quad \Rightarrow \quad F(z) = \int_{\gamma} f(\zeta) d\zeta \text{ does not depend on } \gamma.$$


Consider  $q \in U$ ,  $\gamma$  a path from  $p$  to  $q$ , and  $\Delta_r(q) \subset U$ . For  $z \in \Delta_r(q)$

$$\begin{aligned} F(z) &= \int_{\gamma+[q,z]} f(\zeta) d\zeta \\ &= \int_{\gamma} f(\zeta) d\zeta + \int_{[q,z]} f(\zeta) d\zeta. \end{aligned}$$

The first term is a constant.

The second term is how we defined a primitive in a star-like domain ( $\Delta_r(q)$ ).

See Proposition 3.2.11.



## Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic.*

## Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$e^{g(z)} = f(z).$$

## Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ .

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ .

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ . Then

$$\frac{d}{dz} \left[ \frac{e^{g(z)}}{f(z)} \right]$$



## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ . Then

$$\frac{d}{dz} \left[ \frac{e^{g(z)}}{f(z)} \right] = \frac{e^{g(z)} g'(z) f(z) - e^{g(z)} f'(z)}{(f(z))^2}$$

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ . Then

$$\frac{d}{dz} \left[ \frac{e^{g(z)}}{f(z)} \right] = \frac{e^{g(z)}g'(z)f(z) - e^{g(z)}f'(z)}{(f(z))^2} = \frac{e^{g(z)}f'(z) - e^{g(z)}f'(z)}{(f(z))^2}$$

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ . Then

$$\frac{d}{dz} \left[ \frac{e^{g(z)}}{f(z)} \right] = \frac{e^{g(z)}g'(z)f(z) - e^{g(z)}f'(z)}{(f(z))^2} = \frac{e^{g(z)}f'(z) - e^{g(z)}f'(z)}{(f(z))^2} = 0.$$

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ . Then

$$\frac{d}{dz} \left[ \frac{e^{g(z)}}{f(z)} \right] = \frac{e^{g(z)}g'(z)f(z) - e^{g(z)}f'(z)}{(f(z))^2} = \frac{e^{g(z)}f'(z) - e^{g(z)}f'(z)}{(f(z))^2} = 0.$$

$\Rightarrow \frac{e^{g(z)}}{f(z)}$  is constant.

## Corollary

Let  $U \subset \mathbb{C}$  be a simply connected domain and  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic. Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that

$$e^{g(z)} = f(z).$$

**Example:** If  $U \subset \mathbb{C} \setminus \{0\}$  is a simply connected domain, then  $\exists$  a holomorphic  $L: U \rightarrow \mathbb{C}$  such that  $e^{L(z)} = z$  (a branch of the log).

**Proof:**  $\frac{f'(z)}{f(z)}$  is holomorphic on  $U$ . Find a primitive  $g(z)$ . Then

$$\frac{d}{dz} \left[ \frac{e^{g(z)}}{f(z)} \right] = \frac{e^{g(z)}g'(z)f(z) - e^{g(z)}f'(z)}{(f(z))^2} = \frac{e^{g(z)}f'(z) - e^{g(z)}f'(z)}{(f(z))^2} = 0.$$

$\Rightarrow \frac{e^{g(z)}}{f(z)}$  is constant.

$\Rightarrow \exists C \in \mathbb{C}$  such that  $e^{g(z)+C} = f(z)$ .



If we have the logarithm, we can take roots.

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ .*

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$



If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$

**Proof:** Find a  $\psi: U \rightarrow \mathbb{C}$  such that  $e^{\psi(z)} = f(z)$ .

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$

**Proof:** Find a  $\psi: U \rightarrow \mathbb{C}$  such that  $e^{\psi(z)} = f(z)$ .

Let  $g(z) = e^{\frac{1}{k}\psi(z)}$ .

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$

**Proof:** Find a  $\psi: U \rightarrow \mathbb{C}$  such that  $e^{\psi(z)} = f(z)$ .

Let  $g(z) = e^{\frac{1}{k}\psi(z)}$ .

Check:  $(g(z))^k$

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$

**Proof:** Find a  $\psi: U \rightarrow \mathbb{C}$  such that  $e^{\psi(z)} = f(z)$ .

Let  $g(z) = e^{\frac{1}{k}\psi(z)}$ .

Check:  $(g(z))^k = \left(e^{\frac{1}{k}\psi(z)}\right)^k$

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$

**Proof:** Find a  $\psi: U \rightarrow \mathbb{C}$  such that  $e^{\psi(z)} = f(z)$ .

Let  $g(z) = e^{\frac{1}{k}\psi(z)}$ .

Check:  $(g(z))^k = \left(e^{\frac{1}{k}\psi(z)}\right)^k = e^{\psi(z)}$

If we have the logarithm, we can take roots.

### Corollary

*Let  $U \subset \mathbb{C}$  be a simply connected domain,  $f: U \rightarrow \mathbb{C}$  nowhere zero and holomorphic, and  $k \in \mathbb{N}$ . Then there exists a holomorphic  $g: U \rightarrow \mathbb{C}$  such that*

$$(g(z))^k = f(z).$$

**Proof:** Find a  $\psi: U \rightarrow \mathbb{C}$  such that  $e^{\psi(z)} = f(z)$ .

Let  $g(z) = e^{\frac{1}{k}\psi(z)}$ .

Check:  $(g(z))^k = \left(e^{\frac{1}{k}\psi(z)}\right)^k = e^{\psi(z)} = f(z).$

