

Cultivating Complex Analysis:  
The geometry and topology of the plane (1.1.2)  
Complex-valued functions (1.1.3)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$0 \leq |z\bar{w} - \bar{z}w|^2$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$0 \leq |z\bar{w} - \bar{z}w|^2 = (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w})$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$0 \leq |z\bar{w} - \bar{z}w|^2 = (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2$$



We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$\begin{aligned} 0 \leq |z\bar{w} - \bar{z}w|^2 &= (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2 \\ &= 4z\bar{z}w\bar{w} - (z\bar{w} + \bar{z}w)^2 \end{aligned}$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$\begin{aligned} 0 \leq |z\bar{w} - \bar{z}w|^2 &= (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2 \\ &= 4z\bar{z}w\bar{w} - (z\bar{w} + \bar{z}w)^2 = (2|z||w|)^2 - (2\operatorname{Re} z\bar{w})^2. \end{aligned}$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$\begin{aligned} 0 \leq |z\bar{w} - \bar{z}w|^2 &= (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2 \\ &= 4z\bar{z}w\bar{w} - (z\bar{w} + \bar{z}w)^2 = (2|z||w|)^2 - (2\operatorname{Re} z\bar{w})^2. \end{aligned}$$

Triangle inequality:

$$|z + w|^2 = (z + w)(\bar{z} + \bar{w})$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$\begin{aligned} 0 \leq |z\bar{w} - \bar{z}w|^2 &= (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2 \\ &= 4z\bar{z}w\bar{w} - (z\bar{w} + \bar{z}w)^2 = (2|z||w|)^2 - (2\operatorname{Re} z\bar{w})^2. \end{aligned}$$

Triangle inequality:

$$|z + w|^2 = (z + w)(\bar{z} + \bar{w}) = z\bar{z} + w\bar{w} + z\bar{w} + \bar{z}w$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$\begin{aligned} 0 \leq |z\bar{w} - \bar{z}w|^2 &= (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2 \\ &= 4z\bar{z}w\bar{w} - (z\bar{w} + \bar{z}w)^2 = (2|z||w|)^2 - (2\operatorname{Re} z\bar{w})^2. \end{aligned}$$

Triangle inequality:

$$|z + w|^2 = (z + w)(\bar{z} + \bar{w}) = z\bar{z} + w\bar{w} + z\bar{w} + \bar{z}w \leq z\bar{z} + w\bar{w} + 2|z||w|$$

We measure distance with the *modulus* (euclidean distance from zero):

$$|z| \stackrel{\text{def}}{=} \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}.$$

The (euclidean) distance  $|z - w|$  makes  $\mathbb{C}$  into a complete metric space.

### Proposition (Cauchy–Schwarz and the triangle inequality)

If  $z, w \in \mathbb{C}$ , then

- (i)  $|\operatorname{Re} z\bar{w}| \leq |z||w|$  (Cauchy–Schwarz inequality, note:  $\operatorname{Re} z\bar{w}$  is the real dot product),
- (ii)  $|z + w| \leq |z| + |w|$  (Triangle inequality).

**Proof:** Cauchy–Schwarz:

$$\begin{aligned} 0 \leq |z\bar{w} - \bar{z}w|^2 &= (z\bar{w} - \bar{z}w)(\bar{z}w - z\bar{w}) = 2z\bar{z}w\bar{w} - z^2\bar{w}^2 - \bar{z}^2w^2 \\ &= 4z\bar{z}w\bar{w} - (z\bar{w} + \bar{z}w)^2 = (2|z||w|)^2 - (2\operatorname{Re} z\bar{w})^2. \end{aligned}$$

Triangle inequality:

$$|z + w|^2 = (z + w)(\bar{z} + \bar{w}) = z\bar{z} + w\bar{w} + z\bar{w} + \bar{z}w \leq z\bar{z} + w\bar{w} + 2|z||w| = (|z| + |w|)^2. \quad \square$$

## Proposition

*Complex addition, multiplication, division, and conjugation are continuous: Suppose  $\{a_n\}$  and  $\{b_n\}$  are two convergent sequences of complex numbers. Then,*

$$(i) \quad \lim_{n \rightarrow \infty} (a_n + b_n) = \left( \lim_{n \rightarrow \infty} a_n \right) + \left( \lim_{n \rightarrow \infty} b_n \right),$$

$$(ii) \quad \lim_{n \rightarrow \infty} a_n b_n = \left( \lim_{n \rightarrow \infty} a_n \right) \left( \lim_{n \rightarrow \infty} b_n \right),$$

$$(iii) \quad \lim_{n \rightarrow \infty} \frac{1}{a_n} = \frac{1}{\lim_{n \rightarrow \infty} a_n}, \text{ as long as } \lim_{n \rightarrow \infty} a_n \neq 0,$$

$$(iv) \quad \lim_{n \rightarrow \infty} \bar{a}_n = \overline{\lim_{n \rightarrow \infty} a_n}.$$

## Proposition

*Complex addition, multiplication, division, and conjugation are continuous: Suppose  $\{a_n\}$  and  $\{b_n\}$  are two convergent sequences of complex numbers. Then,*

- (i)  $\lim_{n \rightarrow \infty} (a_n + b_n) = \left( \lim_{n \rightarrow \infty} a_n \right) + \left( \lim_{n \rightarrow \infty} b_n \right),$
- (ii)  $\lim_{n \rightarrow \infty} a_n b_n = \left( \lim_{n \rightarrow \infty} a_n \right) \left( \lim_{n \rightarrow \infty} b_n \right),$
- (iii)  $\lim_{n \rightarrow \infty} \frac{1}{a_n} = \frac{1}{\lim_{n \rightarrow \infty} a_n},$  as long as  $\lim_{n \rightarrow \infty} a_n \neq 0,$
- (iv)  $\lim_{n \rightarrow \infty} \bar{a}_n = \overline{\lim_{n \rightarrow \infty} a_n}.$

All these operations are defined in terms of operations on the real and imaginary parts which are continuous. Details left as exercise.



If  $p \in \mathbb{C}$  and  $r > 0$ , define the *disc* of radius  $r$  around  $p$  as

$$\Delta_r(p) \stackrel{\text{def}}{=} \{z \in \mathbb{C} : |z - p| < r\}.$$

If  $p \in \mathbb{C}$  and  $r > 0$ , define the *disc* of radius  $r$  around  $p$  as

$$\Delta_r(p) \stackrel{\text{def}}{=} \{z \in \mathbb{C} : |z - p| < r\}.$$

$$\mathbb{D} \stackrel{\text{def}}{=} \Delta_1(0) = \{z \in \mathbb{C} : |z| < 1\} \quad \textit{unit disc}.$$

If  $p \in \mathbb{C}$  and  $r > 0$ , define the *disc* of radius  $r$  around  $p$  as

$$\Delta_r(p) \stackrel{\text{def}}{=} \{z \in \mathbb{C} : |z - p| < r\}.$$

$$\mathbb{D} \stackrel{\text{def}}{=} \Delta_1(0) = \{z \in \mathbb{C} : |z| < 1\} \quad \textit{unit disc}.$$

A useful “version” of  $\mathbb{D}$  is the *upper half-plane*:

$$\mathbb{H} \stackrel{\text{def}}{=} \{z \in \mathbb{C} : \text{Im } z > 0\}.$$

If  $p \in \mathbb{C}$  and  $r > 0$ , define the *disc* of radius  $r$  around  $p$  as

$$\Delta_r(p) \stackrel{\text{def}}{=} \{z \in \mathbb{C} : |z - p| < r\}.$$

$$\mathbb{D} \stackrel{\text{def}}{=} \Delta_1(0) = \{z \in \mathbb{C} : |z| < 1\} \quad \text{unit disc.}$$

A useful “version” of  $\mathbb{D}$  is the *upper half-plane*:

$$\mathbb{H} \stackrel{\text{def}}{=} \{z \in \mathbb{C} : \text{Im } z > 0\}.$$

### Definition

An open and connected set  $U \subset \mathbb{C}$  is called a *domain*.

If  $f: X \rightarrow \mathbb{C}$ , write  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ :  $f = u + iv$ .

If  $f: X \rightarrow \mathbb{C}$ , write  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ :  $f = u + iv$ .

If  $X \subset \mathbb{C}$  ( $z = x + iy$ ):  $\frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$  and  $\frac{\partial f}{\partial y} = \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y}$ .

If  $f: X \rightarrow \mathbb{C}$ , write  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ :  $f = u + iv$ .

If  $X \subset \mathbb{C}$  ( $z = x + iy$ ):  $\frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$  and  $\frac{\partial f}{\partial y} = \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y}$ .

If  $X \subset \mathbb{R}$ :  $f' = u' + iv'$ .

If  $f: X \rightarrow \mathbb{C}$ , write  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ :  $f = u + iv$ .

If  $X \subset \mathbb{C}$  ( $z = x + iy$ ):  $\frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$  and  $\frac{\partial f}{\partial y} = \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y}$ .

If  $X \subset \mathbb{R}$ :  $f' = u' + iv'$ .

If  $f: [a, b] \rightarrow \mathbb{C}$ ,  $f$  is (Riemann) integrable if  $u$  and  $v$  are, and

$$\int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$



If  $f: X \rightarrow \mathbb{C}$ , write  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ :  $f = u + iv$ .

If  $X \subset \mathbb{C}$  ( $z = x + iy$ ):  $\frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$  and  $\frac{\partial f}{\partial y} = \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y}$ .

If  $X \subset \mathbb{R}$ :  $f' = u' + iv'$ .

If  $f: [a, b] \rightarrow \mathbb{C}$ ,  $f$  is (Riemann) integrable if  $u$  and  $v$  are, and

$$\int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

## Proposition

Suppose  $f: [a, b] \rightarrow \mathbb{C}$  is (Riemann) integrable. Then  $|f|$  is (Riemann) integrable and

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt.$$