

Cultivating Complex Analysis: Inverse function theorem and automorphisms (2.2.3)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Let $U, V \subset \mathbb{C}$ be open.

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

If $U = V$, then f is called an *automorphism*.

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

If $U = V$, then f is called an *automorphism*.

(Remark that the word “automorphism” appears in many different contexts.)

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

If $U = V$, then f is called an *automorphism*.

(Remark that the word “automorphism” appears in many different contexts.)

Denote the set of automorphism by $\text{Aut}(U)$, which is a group under composition (exercise).

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

If $U = V$, then f is called an *automorphism*.

(Remark that the word “automorphism” appears in many different contexts.)

Denote the set of automorphism by $\text{Aut}(U)$, which is a group under composition (exercise).

Remark: Traditionally, a biholomorphism is called a *conformal mapping* and biholomorphic U and V are said to be *conformally equivalent*.

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

If $U = V$, then f is called an *automorphism*.

(Remark that the word “automorphism” appears in many different contexts.)

Denote the set of automorphism by $\text{Aut}(U)$, which is a group under composition (exercise).

Remark: Traditionally, a biholomorphism is called a *conformal mapping* and biholomorphic U and V are said to be *conformally equivalent*.

Example: The Cayley map $C(z) = \frac{z-i}{z+i}$ takes $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ to \mathbb{D} , and has an inverse. That is, $C|_{\mathbb{H}}: \mathbb{H} \rightarrow \mathbb{D}$ is a biholomorphism.

Let $U, V \subset \mathbb{C}$ be open.

If $f: U \rightarrow V$ is holomorphic and bijective and f^{-1} is holomorphic, then f is called a *biholomorphism*, and say U and V are *biholomorphic*.

(surprisingly, we will later see that f^{-1} is automatically holomorphic).

If $U = V$, then f is called an *automorphism*.

(Remark that the word “automorphism” appears in many different contexts.)

Denote the set of automorphism by $\text{Aut}(U)$, which is a group under composition (exercise).

Remark: Traditionally, a biholomorphism is called a *conformal mapping* and biholomorphic U and V are said to be *conformally equivalent*.

Example: The Cayley map $C(z) = \frac{z-i}{z+i}$ takes $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$ to \mathbb{D} , and has an inverse. That is, $C|_{\mathbb{H}}: \mathbb{H} \rightarrow \mathbb{D}$ is a biholomorphism.

Example: For any $a, b \in \mathbb{C}$, $a \neq 0$, the function $az + b$ is an automorphism of \mathbb{C} .

Suppose f is a biholomorphism.

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z ,

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$.

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Using the Cauchy–Riemann equations, we compute the Jacobian determinant,

$$\det Df$$

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Using the Cauchy–Riemann equations, we compute the Jacobian determinant,

$$\det Df = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x}$$

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Using the Cauchy–Riemann equations, we compute the Jacobian determinant,

$$\det Df = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x} = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2$$

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Using the Cauchy–Riemann equations, we compute the Jacobian determinant,

$$\det Df = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x} = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 = |f'(z)|^2.$$

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Using the Cauchy–Riemann equations, we compute the Jacobian determinant,

$$\det Df = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x} = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 = |f'(z)|^2.$$

$\det Df$ is nonzero (positive) and Df is invertible $\Leftrightarrow f'(z) \neq 0$.

Suppose f is a biholomorphism.

Differentiate $f^{-1}(f(z)) = z$ to find $(f^{-1})'(f(z))f'(z) = 1$.

So $f'(z) \neq 0$ for all z , and if $w = f(z)$, then

$$(f^{-1})'(w) = \frac{1}{f'(z)} \quad \text{or} \quad f'(z) = \frac{1}{(f^{-1})'(w)}.$$

Consider a holomorphic $f = u + iv$. Its real derivative is (as usual $z = x + iy$)

$$Df = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

Using the Cauchy–Riemann equations, we compute the Jacobian determinant,

$$\det Df = \frac{\partial u}{\partial x} \frac{\partial v}{\partial y} - \frac{\partial u}{\partial y} \frac{\partial v}{\partial x} = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 = |f'(z)|^2.$$

$\det Df$ is nonzero (positive) and Df is invertible $\Leftrightarrow f'(z) \neq 0$.

Note that we also found that a holomorphic f preserves orientation (positive $\det Df$).

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

If a 2×2 matrix represents a complex number, its inverse represents the reciprocal.

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

If a 2×2 matrix represents a complex number, its inverse represents the reciprocal.

So in the theorem, if f is holomorphic, then so is f^{-1} .

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

If a 2×2 matrix represents a complex number, its inverse represents the reciprocal.

So in the theorem, if f is holomorphic, then so is f^{-1} . We proved:

Theorem (Inverse function theorem for holomorphic functions)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $p \in U$, and $f'(p) \neq 0$. Suppose further that f is continuously differentiable.

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

If a 2×2 matrix represents a complex number, its inverse represents the reciprocal.

So in the theorem, if f is holomorphic, then so is f^{-1} . We proved:

Theorem (Inverse function theorem for holomorphic functions)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $p \in U$, and $f'(p) \neq 0$. Suppose further that f is continuously differentiable. Then there exist open sets $V, W \subset \mathbb{C}$ such that $p \in V \subset U$, $f(V) = W$, the restriction $f|_V$ is injective (one-to-one), and hence a $g: W \rightarrow V$ exists such that $g(w) = (f|_V)^{-1}(w)$ for all $w \in W$.

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

If a 2×2 matrix represents a complex number, its inverse represents the reciprocal.

So in the theorem, if f is holomorphic, then so is f^{-1} . We proved:

Theorem (Inverse function theorem for holomorphic functions)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $p \in U$, and $f'(p) \neq 0$. Suppose further that f is continuously differentiable. Then there exist open sets $V, W \subset \mathbb{C}$ such that $p \in V \subset U$, $f(V) = W$, the restriction $f|_V$ is injective (one-to-one), and hence a $g: W \rightarrow V$ exists such that $g(w) = (f|_V)^{-1}(w)$ for all $w \in W$. Furthermore, g is holomorphic and

$$g'(w) = \frac{1}{f'(g(w))} \quad \text{for all } w \in W.$$

The real inverse function theorem for continuously differentiable functions of \mathbb{R}^2 to \mathbb{R}^2 says that if Df is invertible at p , then f takes a neighborhood V of p bijectively to a neighborhood $f(V)$ of $f(p)$ and the inverse on that neighborhood is continuously differentiable with $D(f^{-1})|_{f(p)} = (Df|_p)^{-1}$.

If a 2×2 matrix represents a complex number, its inverse represents the reciprocal.

So in the theorem, if f is holomorphic, then so is f^{-1} . We proved:

Theorem (Inverse function theorem for holomorphic functions)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, $p \in U$, and $f'(p) \neq 0$. Suppose further that f is continuously differentiable. Then there exist open sets $V, W \subset \mathbb{C}$ such that $p \in V \subset U$, $f(V) = W$, the restriction $f|_V$ is injective (one-to-one), and hence a $g: W \rightarrow V$ exists such that $g(w) = (f|_V)^{-1}(w)$ for all $w \in W$. Furthermore, g is holomorphic and

$$g'(w) = \frac{1}{f'(g(w))} \quad \text{for all } w \in W.$$

Remark: We will prove later a holomorphic f is always continuously differentiable.

Just because f' is never zero doesn't mean that f is (globally) bijective.

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

However, for a fixed U and V , there are very few biholomorphisms of U and V .

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

However, for a fixed U and V , there are very few biholomorphisms of U and V .

E.g., if $f \in \text{Aut}(\mathbb{D})$, then $f(z) = e^{i\theta} \frac{z-a}{1-\bar{a}z}$,

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

However, for a fixed U and V , there are very few biholomorphisms of U and V .

E.g., if $f \in \text{Aut}(\mathbb{D})$, then $f(z) = e^{i\theta} \frac{z-a}{1-\bar{a}z}$, if $f \in \text{Aut}(\mathbb{C})$, then $f(z) = az + b$.

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

However, for a fixed U and V , there are very few biholomorphisms of U and V .

E.g., if $f \in \text{Aut}(\mathbb{D})$, then $f(z) = e^{i\theta} \frac{z-a}{1-\bar{a}z}$, if $f \in \text{Aut}(\mathbb{C})$, then $f(z) = az + b$.

And comparatively, \mathbb{D} and \mathbb{C} have more automorphisms than most.

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

However, for a fixed U and V , there are very few biholomorphisms of U and V .

E.g., if $f \in \text{Aut}(\mathbb{D})$, then $f(z) = e^{i\theta} \frac{z-a}{1-\bar{a}z}$, if $f \in \text{Aut}(\mathbb{C})$, then $f(z) = az + b$.

And comparatively, \mathbb{D} and \mathbb{C} have more automorphisms than most.

E.g., a disc minus three generically placed points only has the identity automorphism.

Just because f' is never zero doesn't mean that f is (globally) bijective.

e^z is never zero (so neither is its derivative), but $e^z = e^{z+2\pi i}$.

But locally we can invert e^z and we will obtain (locally) the complex logarithm.

Remark: Locally, there are lots of local biholomorphisms (there are lots of holomorphic functions whose derivative is nonzero at a point).

However, for a fixed U and V , there are very few biholomorphisms of U and V .

E.g., if $f \in \text{Aut}(\mathbb{D})$, then $f(z) = e^{i\theta} \frac{z-a}{1-\bar{a}z}$, if $f \in \text{Aut}(\mathbb{C})$, then $f(z) = az + b$.

And comparatively, \mathbb{D} and \mathbb{C} have more automorphisms than most.

E.g., a disc minus three generically placed points only has the identity automorphism.

(Although we don't yet have enough machinery to prove these statements.)