

# Cultivating Complex Analysis: Inverses of holomorphic functions (5.6)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Let us restate the inverse function theorem.

### Theorem (Inverse function theorem for holomorphic functions)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $p \in U$ , and  $f'(p) \neq 0$ . Then there exist open sets  $V, W \subset \mathbb{C}$  such that  $p \in V \subset U$ ,  $f(V) = W$ , the restriction  $f|_V$  is injective (one-to-one), and hence a  $g: W \rightarrow V$  exists such that  $g(w) = (f|_V)^{-1}(w)$  for all  $w \in W$ . Furthermore,  $g$  is holomorphic and*

$$g'(w) = \frac{1}{f'(g(w))} \quad \text{for all } w \in W.$$

Let us restate the inverse function theorem.

### Theorem (Inverse function theorem for holomorphic functions)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $p \in U$ , and  $f'(p) \neq 0$ . Then there exist open sets  $V, W \subset \mathbb{C}$  such that  $p \in V \subset U$ ,  $f(V) = W$ , the restriction  $f|_V$  is injective (one-to-one), and hence a  $g: W \rightarrow V$  exists such that  $g(w) = (f|_V)^{-1}(w)$  for all  $w \in W$ . Furthermore,  $g$  is holomorphic and*

$$g'(w) = \frac{1}{f'(g(w))} \quad \text{for all } w \in W.$$

In other words, if  $f'$  is nonzero somewhere,  $f$  is injective near that point.

Let us restate the inverse function theorem.

### Theorem (Inverse function theorem for holomorphic functions)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $p \in U$ , and  $f'(p) \neq 0$ . Then there exist open sets  $V, W \subset \mathbb{C}$  such that  $p \in V \subset U$ ,  $f(V) = W$ , the restriction  $f|_V$  is injective (one-to-one), and hence a  $g: W \rightarrow V$  exists such that  $g(w) = (f|_V)^{-1}(w)$  for all  $w \in W$ . Furthermore,  $g$  is holomorphic and*

$$g'(w) = \frac{1}{f'(g(w))} \quad \text{for all } w \in W.$$

In other words, if  $f'$  is nonzero somewhere,  $f$  is injective near that point.

Only local:  $f(z) = z^2$  maps  $\mathbb{C} \setminus \{0\}$  to itself,  $f'$  does not vanish, but  $f$  is 2-to-1 globally.

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .



Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Let  $w \in \Delta_\delta(f(p)) \setminus \{f(p)\}$

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Let  $w \in \Delta_\delta(f(p)) \setminus \{f(p)\} \Rightarrow z \mapsto f(z) - w$  has at least two zeros (Rouché).

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Let  $w \in \Delta_\delta(f(p)) \setminus \{f(p)\} \Rightarrow z \mapsto f(z) - w$  has at least two zeros (Rouché).

$f' \neq 0$  in  $\Delta_r(p) \setminus \{p\}$

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Let  $w \in \Delta_\delta(f(p)) \setminus \{f(p)\} \Rightarrow z \mapsto f(z) - w$  has at least two zeros (Rouché).

$f' \neq 0$  in  $\Delta_r(p) \setminus \{p\} \Rightarrow$  zeros of  $z \mapsto f(z) - w$  are simple

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Let  $w \in \Delta_\delta(f(p)) \setminus \{f(p)\} \Rightarrow z \mapsto f(z) - w$  has at least two zeros (Rouché).

$f' \neq 0$  in  $\Delta_r(p) \setminus \{p\} \Rightarrow$  zeros of  $z \mapsto f(z) - w$  are simple

$\Rightarrow f(z) - w$  has at least two distinct zeros

Real functions can be injective and the derivative can vanish:

$f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^3$ , is injective but  $f'(0) = 0$ .

Holomorphic functions locally all behave like  $z \mapsto z^k$ , and that is injective only if  $k = 1$ .

### Lemma

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f'$  is never zero.*

**Proof:** Suppose  $f$  nonconstant and  $f'(p) = 0$ .

Let  $\overline{\Delta_r(p)} \subset U$  be so that  $f' \neq 0$  on  $\Delta_r(p) \setminus \{p\}$ , and  $|f(z) - f(p)| > \delta > 0$  for  $z \in \partial\Delta_r(p)$ .

$z \mapsto f(z) - f(p)$  has a zero of multiplicity at least two.

Let  $w \in \Delta_\delta(f(p)) \setminus \{f(p)\} \Rightarrow z \mapsto f(z) - w$  has at least two zeros (Rouché).

$f' \neq 0$  in  $\Delta_r(p) \setminus \{p\} \Rightarrow$  zeros of  $z \mapsto f(z) - w$  are simple

$\Rightarrow f(z) - w$  has at least two distinct zeros  $\Rightarrow f$  is not injective.





We can actually compute the inverse:

### Lemma

If  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, and  $\overline{\Delta_r(p)} \subset U$ . Then for all  $w \in f(\Delta_r(p))$ ,

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz.$$

We can actually compute the inverse:

### Lemma

If  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, and  $\overline{\Delta_r(p)} \subset U$ . Then for all  $w \in f(\Delta_r(p))$ ,

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz.$$

**Proof:** Fix  $w \in f(\Delta_r(p))$  and  $\zeta \in \Delta_r(p)$  such that  $f(\zeta) = w$ .

We can actually compute the inverse:

### Lemma

If  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, and  $\overline{\Delta_r(p)} \subset U$ . Then for all  $w \in f(\Delta_r(p))$ ,

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz.$$

**Proof:** Fix  $w \in f(\Delta_r(p))$  and  $\zeta \in \Delta_r(p)$  such that  $f(\zeta) = w$ .

$f'$  is never zero, so  $z \mapsto f(z) - w$  has a simple zero at  $z = \zeta$ .

We can actually compute the inverse:

### Lemma

If  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, and  $\overline{\Delta_r(p)} \subset U$ . Then for all  $w \in f(\Delta_r(p))$ ,

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz.$$

**Proof:** Fix  $w \in f(\Delta_r(p))$  and  $\zeta \in \Delta_r(p)$  such that  $f(\zeta) = w$ .

$f'$  is never zero, so  $z \mapsto f(z) - w$  has a simple zero at  $z = \zeta$ .

By the residue theorem

$$\frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz = \text{Res} \left( \frac{f'(z)z}{f(z) - w}; \zeta \right)$$

We can actually compute the inverse:

### Lemma

If  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, and  $\overline{\Delta_r(p)} \subset U$ . Then for all  $w \in f(\Delta_r(p))$ ,

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz.$$

**Proof:** Fix  $w \in f(\Delta_r(p))$  and  $\zeta \in \Delta_r(p)$  such that  $f(\zeta) = w$ .

$f'$  is never zero, so  $z \mapsto f(z) - w$  has a simple zero at  $z = \zeta$ .

By the residue theorem

$$\frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz = \operatorname{Res} \left( \frac{f'(z)z}{f(z) - w}; \zeta \right) = \frac{f'(\zeta)\zeta}{f'(\zeta)} = \zeta = f^{-1}(w). \quad \square$$

We can actually compute the inverse:

### Lemma

If  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, and  $\overline{\Delta_r(p)} \subset U$ . Then for all  $w \in f(\Delta_r(p))$ ,

$$f^{-1}(w) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz.$$

**Proof:** Fix  $w \in f(\Delta_r(p))$  and  $\zeta \in \Delta_r(p)$  such that  $f(\zeta) = w$ .

$f'$  is never zero, so  $z \mapsto f(z) - w$  has a simple zero at  $z = \zeta$ .

By the residue theorem

$$\frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f'(z)z}{f(z) - w} dz = \operatorname{Res} \left( \frac{f'(z)z}{f(z) - w}; \zeta \right) = \frac{f'(\zeta)\zeta}{f'(\zeta)} = \zeta = f^{-1}(w). \quad \square$$

Consequently,  $f^{-1}$  is holomorphic without even using the inverse function theorem.

## Theorem

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f(U)$  is open,  $f'$  is never zero on  $U$ , and  $f^{-1}: f(U) \rightarrow U$  is holomorphic.*

## Theorem

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f(U)$  is open,  $f'$  is never zero on  $U$ , and  $f^{-1}: f(U) \rightarrow U$  is holomorphic.*

**Proof:**  $f(U)$  is open by the open mapping theorem.



## Theorem

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f(U)$  is open,  $f'$  is never zero on  $U$ , and  $f^{-1}: f(U) \rightarrow U$  is holomorphic.*

**Proof:**  $f(U)$  is open by the open mapping theorem.

By one of the lemmas,  $f'$  is never zero on  $U$ .

## Theorem

*If  $U \subset \mathbb{C}$  is open and  $f: U \rightarrow \mathbb{C}$  is holomorphic and injective, then  $f(U)$  is open,  $f'$  is never zero on  $U$ , and  $f^{-1}: f(U) \rightarrow U$  is holomorphic.*

**Proof:**  $f(U)$  is open by the open mapping theorem.

By one of the lemmas,  $f'$  is never zero on  $U$ .

By the other (or IFT),  $f^{-1}$  is holomorphic.

