

Cultivating Complex Analysis: Cauchy–Goursat, the “Cauchy for triangles” (3.2.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

Let $a, b, c \in \mathbb{C}$ be noncollinear.

A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

Let $a, b, c \in \mathbb{C}$ be noncollinear.

A *triangle* T with vertices a, b, c is the convex hull of $\{a, b, c\}$, that is, the smallest convex set containing the points.

A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

Let $a, b, c \in \mathbb{C}$ be noncollinear.

A *triangle* T with vertices a, b, c is the convex hull of $\{a, b, c\}$, that is, the smallest convex set containing the points.

In other words, T is the set of points

$$t_1a + t_2b + t_3c,$$

where $t_1, t_2, t_3 \in [0, 1]$ and $t_1 + t_2 + t_3 = 1$.

A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

Let $a, b, c \in \mathbb{C}$ be noncollinear.


A *triangle* T with vertices a, b, c is the convex hull of $\{a, b, c\}$, that is, the smallest convex set containing the points.

In other words, T is the set of points

$$t_1a + t_2b + t_3c,$$

where $t_1, t_2, t_3 \in [0, 1]$ and $t_1 + t_2 + t_3 = 1$.

The triangle is oriented positively if the vertices are ordered so that a, b, c goes counterclockwise.



A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

Let $a, b, c \in \mathbb{C}$ be noncollinear.

A *triangle* T with vertices a, b, c is the convex hull of $\{a, b, c\}$, that is, the smallest convex set containing the points.

In other words, T is the set of points


$$t_1a + t_2b + t_3c,$$

where $t_1, t_2, t_3 \in [0, 1]$ and $t_1 + t_2 + t_3 = 1$.

The triangle is oriented positively if the vertices are ordered so that a, b, c goes counterclockwise.

The boundary ∂T of T is defined as the cycle

$$\partial T = [a, b] + [b, c] + [c, a].$$



A set X is *convex* if $[a, b] \subset X$ for all $a, b \in X$.

Let $a, b, c \in \mathbb{C}$ be noncollinear.

A *triangle* T with vertices a, b, c is the convex hull of $\{a, b, c\}$, that is, the smallest convex set containing the points.

In other words, T is the set of points

$$t_1 a + t_2 b + t_3 c,$$


where $t_1, t_2, t_3 \in [0, 1]$ and $t_1 + t_2 + t_3 = 1$.

The triangle is oriented positively if the vertices are ordered so that a, b, c goes counterclockwise.

The boundary ∂T of T is defined as the cycle

$$\partial T = [a, b] + [b, c] + [c, a].$$

Note that our triangle T is the **solid** triangle (includes the interior).



Theorem (Cauchy–Goursat)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, and $T \subset U$ is a triangle. Then

$$\int_{\partial T} f(z) dz = 0.$$

Theorem (Cauchy–Goursat)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, and $T \subset U$ is a triangle. Then

$$\int_{\partial T} f(z) dz = 0.$$

It is important is that $T \subset U$ means the whole solid triangle is in U , not just the boundary.

Theorem (Cauchy–Goursat)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, and $T \subset U$ is a triangle. Then

$$\int_{\partial T} f(z) dz = 0.$$

It is important is that $T \subset U$ means the whole solid triangle is in U , not just the boundary.

Remark: It is a “Goursat” theorem not just “Cauchy” because of the proof: We do not assume that f' is continuous as we have not proved that yet.

Theorem (Cauchy–Goursat)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, and $T \subset U$ is a triangle. Then

$$\int_{\partial T} f(z) dz = 0.$$

It is important is that $T \subset U$ means the whole solid triangle is in U , not just the boundary.

Remark: It is a “Goursat” theorem not just “Cauchy” because of the proof: We do not assume that f' is continuous as we have not proved that yet.

Proof: We prove the contrapositive.

Theorem (Cauchy–Goursat)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, and $T \subset U$ is a triangle. Then

$$\int_{\partial T} f(z) dz = 0.$$

It is important is that $T \subset U$ means the whole solid triangle is in U , not just the boundary.

Remark: It is a “Goursat” theorem not just “Cauchy” because of the proof: We do not assume that f' is continuous as we have not proved that yet.

Proof: We prove the contrapositive.

Suppose f is continuous and suppose $\exists T \subset U$ such that

$$\left| \int_{\partial T} f(z) dz \right| = c \neq 0.$$

Theorem (Cauchy–Goursat)

Suppose $U \subset \mathbb{C}$ is open, $f: U \rightarrow \mathbb{C}$ is holomorphic, and $T \subset U$ is a triangle. Then

$$\int_{\partial T} f(z) dz = 0.$$

It is important is that $T \subset U$ means the whole solid triangle is in U , not just the boundary.

Remark: It is a “Goursat” theorem not just “Cauchy” because of the proof: We do not assume that f' is continuous as we have not proved that yet.


Proof: We prove the contrapositive.

Suppose f is continuous and suppose $\exists T \subset U$ such that

$$\left| \int_{\partial T} f(z) dz \right| = c \neq 0.$$

We will find a point where f is not complex differentiable.


Cut T into four subtriangles T_1, T_2, T_3, T_4
(cut each side in half)



Cut T into four subtriangles T_1, T_2, T_3, T_4
(cut each side in half)

Orient each T_j positively:

The inner sides cancel.




Cut T into four subtriangles T_1, T_2, T_3, T_4
(cut each side in half)

Orient each T_j positively:

The inner sides cancel.

So



$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

Cut T into four subtriangles T_1, T_2, T_3, T_4
(cut each side in half)


Orient each T_j positively:

The inner sides cancel.

So

$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

So for some triangle T_j , the integral is at least $\frac{c}{4}$.




Cut T into four subtriangles T_1, T_2, T_3, T_4
 (cut each side in half)

Orient each T_j positively:

The inner sides cancel.

So



$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

So for some triangle T_j , the integral is at least $\frac{c}{4}$.


Label that subtriangle $T^1 = T_j$ and $\left| \int_{\partial T^1} f(z) dz \right| \geq \frac{c}{4}.$

Cut T into four subtriangles T_1, T_2, T_3, T_4
 (cut each side in half)

Orient each T_j positively:

The inner sides cancel.

So



$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

So for some triangle T_j , the integral is at least $\frac{c}{4}$.

Label that subtriangle $T^1 = T_j$ and $\left| \int_{\partial T^1} f(z) dz \right| \geq \frac{c}{4}$.


Cut T^1 into subtriangles $T_1^1, T_2^1, T_3^1, T_4^1$.

Cut T into four subtriangles T_1, T_2, T_3, T_4
 (cut each side in half)

Orient each T_j positively:

The inner sides cancel.

So



$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

So for some triangle T_j , the integral is at least $\frac{c}{4}$.

Label that subtriangle $T^1 = T_j$ and $\left| \int_{\partial T^1} f(z) dz \right| \geq \frac{c}{4}$.


Cut T^1 into subtriangles $T_1^1, T_2^1, T_3^1, T_4^1$. Integral over some ∂T_j^1 is at least $\frac{c}{4^2}$, so label it T^2 .

Cut T into four subtriangles T_1, T_2, T_3, T_4
 (cut each side in half)

Orient each T_j positively:

The inner sides cancel.

So



$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

So for some triangle T_j , the integral is at least $\frac{c}{4}$.

Label that subtriangle $T^1 = T_j$ and $\left| \int_{\partial T^1} f(z) dz \right| \geq \frac{c}{4}.$

Cut T^1 into subtriangles $T_1^1, T_2^1, T_3^1, T_4^1$. Integral over some ∂T_j^1 is at least $\frac{c}{4^2}$, so label it T^2 .


Rinse and repeat.

Cut T into four subtriangles T_1, T_2, T_3, T_4
(cut each side in half)

Orient each T_j positively:

The inner sides cancel.

So



$$c = \left| \int_{\partial T} f(z) dz \right| = \left| \int_{\partial T_1} f(z) dz + \int_{\partial T_2} f(z) dz + \int_{\partial T_3} f(z) dz + \int_{\partial T_4} f(z) dz \right|.$$

So for some triangle T_j , the integral is at least $\frac{c}{4}$.

Label that subtriangle $T^1 = T_j$ and $\left| \int_{\partial T^1} f(z) dz \right| \geq \frac{c}{4}$.

Cut T^1 into subtriangles $T_1^1, T_2^1, T_3^1, T_4^1$. Integral over some ∂T_j^1 is at least $\frac{c}{4^2}$, so label it T^2 .

Rinse and repeat.

After k iterations for the k^{th} triangle T^k , $\left| \int_{\partial T^k} f(z) dz \right| \geq \frac{c}{4^k}$.

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

Were f complex differentiable at z_0 , then for some α , $\frac{g(z)}{z-z_0}$ would go to zero as $z \rightarrow z_0$.

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

Were f complex differentiable at z_0 , then for some α , $\frac{g(z)}{z-z_0}$ would go to zero as $z \rightarrow z_0$.

We will prove $\frac{g(z)}{z-z_0}$ never goes to zero (no matter what α is).

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

Were f complex differentiable at z_0 , then for some α , $\frac{g(z)}{z-z_0}$ would go to zero as $z \rightarrow z_0$.

We will prove $\frac{g(z)}{z-z_0}$ never goes to zero (no matter what α is).

Fix α .

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

Were f complex differentiable at z_0 , then for some α , $\frac{g(z)}{z-z_0}$ would go to zero as $z \rightarrow z_0$.

We will prove $\frac{g(z)}{z-z_0}$ never goes to zero (no matter what α is).

Fix α . If $g(z_0) \neq 0$, we are done.

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

Were f complex differentiable at z_0 , then for some α , $\frac{g(z)}{z-z_0}$ would go to zero as $z \rightarrow z_0$.

We will prove $\frac{g(z)}{z-z_0}$ never goes to zero (no matter what α is).

Fix α . If $g(z_0) \neq 0$, we are done.

So assume $g(z_0) = 0$.

$T^k \subset T^{k-1} \subset \dots \subset T$ and in each step the triangle is exactly half the size (similar triangles):

$$\text{diam}(T^k) = \frac{1}{2} \text{diam}(T^{k-1}) = \frac{1}{2^k} \text{diam}(T).$$

The triangles are compact \Rightarrow the intersection is nonempty.

The diameter goes to zero \Rightarrow the intersection is a single point:

$$\{z_0\} = \bigcap_{k=1}^{\infty} T^k.$$

Write $f(z) = f(z_0) + \alpha(z - z_0) + g(z)$ for some $\alpha \in \mathbb{C}$.

Were f complex differentiable at z_0 , then for some α , $\frac{g(z)}{z-z_0}$ would go to zero as $z \rightarrow z_0$.

We will prove $\frac{g(z)}{z-z_0}$ never goes to zero (no matter what α is).

Fix α . If $g(z_0) \neq 0$, we are done.

So assume $g(z_0) = 0$. Cauchy's theorem for polynomials says

$$\int_{\partial T^k} f(z) dz = \int_{\partial T^k} (f(z_0) + \alpha(z - z_0) + g(z)) dz = \int_{\partial T^k} g(z) dz.$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) \, dz \right|$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) \, dz \right| = \left| \int_{\partial T^k} g(z) \, dz \right|$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) \, dz \right| = \left| \int_{\partial T^k} g(z) \, dz \right| \leq \int_{\partial T^k} |g(z)| \, |dz|.$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$,

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right|$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right| \geq \frac{2^k |g(z_k)|}{d}$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right| \geq \frac{2^k |g(z_k)|}{d} = \frac{4^k}{d\ell} \int_{\partial T^k} |g(z)| |dz|$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right| \geq \frac{2^k |g(z_k)|}{d} = \frac{4^k}{d\ell} \int_{\partial T^k} |g(z)| |dz| \geq \frac{4^k}{d\ell} \frac{c}{4^k}$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right| \geq \frac{2^k |g(z_k)|}{d} = \frac{4^k}{d\ell} \int_{\partial T^k} |g(z)| |dz| \geq \frac{4^k}{d\ell} \frac{c}{4^k} = \frac{c}{d\ell}.$$

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right| \geq \frac{2^k |g(z_k)|}{d} = \frac{4^k}{d\ell} \int_{\partial T^k} |g(z)| |dz| \geq \frac{4^k}{d\ell} \frac{c}{4^k} = \frac{c}{d\ell}.$$

Since $z_k \rightarrow z_0$, we have that $\frac{g(z)}{z-z_0}$ does not go to zero as $z \rightarrow z_0$.

$$\frac{c}{4^k} \leq \left| \int_{\partial T^k} f(z) dz \right| = \left| \int_{\partial T^k} g(z) dz \right| \leq \int_{\partial T^k} |g(z)| |dz|.$$

Let ℓ be the length of ∂T .

The length of ∂T^k is $\frac{\ell}{2^k}$, by the mean value theorem for integrals, $\exists z_k \in \partial T^k$ such that

$$|g(z_k)| = \frac{2^k}{\ell} \int_{\partial T^k} |g(z)| |dz|.$$

$z_k \neq z_0$ as $g(z_0) = 0$.

Let $d = \text{diam}(T)$. Then $|z_k - z_0| \leq \frac{d}{2^k}$ and

$$\left| \frac{g(z_k)}{z_k - z_0} \right| \geq \frac{2^k |g(z_k)|}{d} = \frac{4^k}{d\ell} \int_{\partial T^k} |g(z)| |dz| \geq \frac{4^k}{d\ell} \frac{c}{4^k} = \frac{c}{d\ell}.$$

Since $z_k \rightarrow z_0$, we have that $\frac{g(z)}{z-z_0}$ does not go to zero as $z \rightarrow z_0$.

So f is not complex differentiable at z_0 .

□

A useful version of this result is the following exercise:

Exercise: Suppose $T \subset \mathbb{C}$ is a triangle and $f: T \rightarrow \mathbb{C}$ a continuous function whose restriction to the interior of T is holomorphic. Prove that $\int_{\partial T} f(z) dz = 0$.

A useful version of this result is the following exercise:

Exercise: Suppose $T \subset \mathbb{C}$ is a triangle and $f: T \rightarrow \mathbb{C}$ a continuous function whose restriction to the interior of T is holomorphic. Prove that $\int_{\partial T} f(z) dz = 0$.

Hint: Passing some sort of limit under the integral is required.