

# Cultivating Complex Analysis: Homology versions of Cauchy (4.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

Let  $U \subset \mathbb{C}$  be open and  $\Gamma$  a cycle in  $U$  such that  $n(\Gamma; p) = 0$  for all  $p \in \mathbb{C} \setminus U$ , then we say  $\Gamma$  is *homologous to zero in  $U$* .

## Definition

Let  $U \subset \mathbb{C}$  be open and  $\Gamma$  a cycle in  $U$  such that  $n(\Gamma; p) = 0$  for all  $p \in \mathbb{C} \setminus U$ , then we say  $\Gamma$  is *homologous to zero in  $U$* .

“homologous to zero in  $U$ ” = “does not wind around any point in  $\mathbb{C} \setminus U$ .”

## Definition

Let  $U \subset \mathbb{C}$  be open and  $\Gamma$  a cycle in  $U$  such that  $n(\Gamma; p) = 0$  for all  $p \in \mathbb{C} \setminus U$ , then we say  $\Gamma$  is *homologous to zero in  $U$* .

“homologous to zero in  $U$ ” = “does not wind around any point in  $\mathbb{C} \setminus U$ .”

**Note:** “homologous to zero” does **not** mean “equivalent to zero.”

## Definition

Let  $U \subset \mathbb{C}$  be open and  $\Gamma$  a cycle in  $U$  such that  $n(\Gamma; p) = 0$  for all  $p \in \mathbb{C} \setminus U$ , then we say  $\Gamma$  is *homologous to zero in  $U$* .

“homologous to zero in  $U$ ” = “does not wind around any point in  $\mathbb{C} \setminus U$ .”

**Note:** “homologous to zero” does **not** mean “equivalent to zero.”

**Example:** In  $U = \mathbb{C}$ , every  $\Gamma$  is homologous to zero.

## Definition

Let  $U \subset \mathbb{C}$  be open and  $\Gamma$  a cycle in  $U$  such that  $n(\Gamma; p) = 0$  for all  $p \in \mathbb{C} \setminus U$ , then we say  $\Gamma$  is *homologous to zero in  $U$* .

“homologous to zero in  $U$ ” = “does not wind around any point in  $\mathbb{C} \setminus U$ .”

**Note:** “homologous to zero” does **not** mean “equivalent to zero.”

**Example:** In  $U = \mathbb{C}$ , every  $\Gamma$  is homologous to zero.

**Example:** The unit circle is homologous to zero in  $\mathbb{C}$ , but not homologous to zero in  $\mathbb{C} \setminus \{0\}$ .

## Definition

Let  $U \subset \mathbb{C}$  be open and  $\Gamma$  a cycle in  $U$  such that  $n(\Gamma; p) = 0$  for all  $p \in \mathbb{C} \setminus U$ , then we say  $\Gamma$  is *homologous to zero in  $U$* .

“homologous to zero in  $U$ ” = “does not wind around any point in  $\mathbb{C} \setminus U$ .”

**Note:** “homologous to zero” does **not** mean “equivalent to zero.”

**Example:** In  $U = \mathbb{C}$ , every  $\Gamma$  is homologous to zero.

**Example:** The unit circle is homologous to zero in  $\mathbb{C}$ , but not homologous to zero in  $\mathbb{C} \setminus \{0\}$ .

## Theorem (Cauchy integral formula (homology version))

Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then for  $z \in U \setminus \Gamma$ ,

$$n(\Gamma; z)f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta) - f(z)}{\zeta - z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$



**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta) - f(z)}{\zeta - z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  
 $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  
 $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

Suppose  $z \in U \setminus \Gamma$  and  $n(\Gamma; z) = 0$ .

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

Suppose  $z \in U \setminus \Gamma$  and  $n(\Gamma; z) = 0$ .

$$\int_{\Gamma} g(\zeta, z) d\zeta$$

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

Suppose  $z \in U \setminus \Gamma$  and  $n(\Gamma; z) = 0$ .

$$\int_{\Gamma} g(\zeta, z) d\zeta = \int_{\Gamma} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta$$



**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

Suppose  $z \in U \setminus \Gamma$  and  $n(\Gamma; z) = 0$ .

$$\int_{\Gamma} g(\zeta, z) d\zeta = \int_{\Gamma} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - f(z)(2\pi i)n(\Gamma; z)$$

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

Suppose  $z \in U \setminus \Gamma$  and  $n(\Gamma; z) = 0$ .

$$\int_{\Gamma} g(\zeta, z) d\zeta = \int_{\Gamma} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - f(z)(2\pi i)n(\Gamma; z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

**Proof:** Define  $g: U \times U \rightarrow \mathbb{C}$  by

$$g(\zeta, z) = \begin{cases} \frac{f(\zeta)-f(z)}{\zeta-z} & \text{if } \zeta \neq z, \\ f'(\zeta) & \text{if } \zeta = z. \end{cases}$$

**Exercise:**  $g(\zeta, z)$  is continuous in  $U \times U$ , and  $z \mapsto g(\zeta, z)$  is holomorphic for every fixed  $\zeta \in U$ .

Let

$$h(z) = \begin{cases} \int_{\Gamma} g(\zeta, z) d\zeta & \text{if } z \in U, \\ \int_{\Gamma} \frac{f(\zeta)}{\zeta-z} d\zeta & \text{if } z \notin \Gamma \text{ and } n(\Gamma; z) = 0. \end{cases}$$

$h(z)$  is defined for all  $z \in \mathbb{C}$  (as  $n(\Gamma; z) = 0$  for all  $z \in \mathbb{C} \setminus U$ ).

But, at some points we have two definitions!

Suppose  $z \in U \setminus \Gamma$  and  $n(\Gamma; z) = 0$ .

$$\int_{\Gamma} g(\zeta, z) d\zeta = \int_{\Gamma} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - f(z)(2\pi i)n(\Gamma; z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

So  $h: \mathbb{C} \rightarrow \mathbb{C}$  is well-defined.

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ .



WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)|$$

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right|$$

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta|$$

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta| \leq \frac{M\ell}{d(z, \Gamma)} \quad (d(z, \Gamma) \text{ is the distance of } z \text{ and } \Gamma).$$

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta| \leq \frac{M\ell}{d(z, \Gamma)} \quad (d(z, \Gamma) \text{ is the distance of } z \text{ and } \Gamma).$$

$$z \rightarrow \infty \quad \Rightarrow \quad d(z, \Gamma) \rightarrow \infty \quad \Rightarrow \quad h(z) \rightarrow 0$$



WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta| \leq \frac{M\ell}{d(z, \Gamma)} \quad (d(z, \Gamma) \text{ is the distance of } z \text{ and } \Gamma).$$

$$z \rightarrow \infty \quad \Rightarrow \quad d(z, \Gamma) \rightarrow \infty \quad \Rightarrow \quad h(z) \rightarrow 0$$

So  $h$  is bounded, by Liouville  $h$  is constant, and the constant is zero.

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta| \leq \frac{M\ell}{d(z, \Gamma)} \quad (d(z, \Gamma) \text{ is the distance of } z \text{ and } \Gamma).$$

$$z \rightarrow \infty \quad \Rightarrow \quad d(z, \Gamma) \rightarrow \infty \quad \Rightarrow \quad h(z) \rightarrow 0$$

So  $h$  is bounded, by Liouville  $h$  is constant, and the constant is zero.

Suppose  $z \in U \setminus \Gamma$ . Then

$$0 = h(z)$$

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta| \leq \frac{M\ell}{d(z, \Gamma)} \quad (d(z, \Gamma) \text{ is the distance of } z \text{ and } \Gamma).$$

$$z \rightarrow \infty \quad \Rightarrow \quad d(z, \Gamma) \rightarrow \infty \quad \Rightarrow \quad h(z) \rightarrow 0$$

So  $h$  is bounded, by Liouville  $h$  is constant, and the constant is zero.

Suppose  $z \in U \setminus \Gamma$ . Then

$$0 = h(z) = \int_{\Gamma} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta$$

WTS  $h: \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic.

Holomorphicity is a local property, so only consider  $h$  on open sets covering  $\mathbb{C}$ .

The set of  $z \notin \Gamma$  where  $n(\Gamma; z) = 0$  is open, and  $z \mapsto \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$  is holomorphic there.

Similarly  $U$  is open and  $z \mapsto \int_{\Gamma} g(\zeta, z) d\zeta$  is holomorphic there, so  $h$  is holomorphic.

Consider  $z$  in the unbounded component of  $\mathbb{C} \setminus \Gamma$ . Then  $n(\Gamma; z) = 0$ , so  $h(z) = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$ .

Let  $M$  be such that  $|f(\zeta)| \leq M$  for  $\zeta \in \Gamma$  and  $\ell$  be the length of  $\Gamma$ .

$$|h(z)| = \left| \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta \right| \leq \int_{\Gamma} \left| \frac{f(\zeta)}{\zeta - z} \right| |d\zeta| \leq \frac{M\ell}{d(z, \Gamma)} \quad (d(z, \Gamma) \text{ is the distance of } z \text{ and } \Gamma).$$

$$z \rightarrow \infty \quad \Rightarrow \quad d(z, \Gamma) \rightarrow \infty \quad \Rightarrow \quad h(z) \rightarrow 0$$

So  $h$  is bounded, by Liouville  $h$  is constant, and the constant is zero.

Suppose  $z \in U \setminus \Gamma$ . Then

$$0 = h(z) = \int_{\Gamma} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta = \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - f(z) (2\pi i) n(\Gamma; z). \quad \square$$

### Theorem (Cauchy's theorem (homology version))

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then*

$$\int_{\Gamma} f(z) dz = 0.$$

### Theorem (Cauchy's theorem (homology version))

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then*

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Fix  $z \in U \setminus \Gamma$ .

### Theorem (Cauchy's theorem (homology version))

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then*

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Fix  $z \in U \setminus \Gamma$ .

Apply the Cauchy integral formula to  $\zeta \mapsto (\zeta - z)f(\zeta)$  at  $\zeta = z$ :

### Theorem (Cauchy's theorem (homology version))

Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Fix  $z \in U \setminus \Gamma$ .

Apply the Cauchy integral formula to  $\zeta \mapsto (\zeta - z)f(\zeta)$  at  $\zeta = z$ :

$$0 = n(\Gamma; z)(z - z)f(z)$$



### Theorem (Cauchy's theorem (homology version))

Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Fix  $z \in U \setminus \Gamma$ .

Apply the Cauchy integral formula to  $\zeta \mapsto (\zeta - z)f(\zeta)$  at  $\zeta = z$ :

$$0 = n(\Gamma; z)(z - z)f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{(\zeta - z)f(\zeta)}{\zeta - z} d\zeta$$

### Theorem (Cauchy's theorem (homology version))

Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Fix  $z \in U \setminus \Gamma$ .

Apply the Cauchy integral formula to  $\zeta \mapsto (\zeta - z)f(\zeta)$  at  $\zeta = z$ :

$$0 = n(\Gamma; z)(z - z)f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{(\zeta - z)f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\Gamma} f(\zeta) d\zeta. \quad \square$$

### Theorem (Cauchy's theorem (homology version))

Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic, and  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ . Then

$$\int_{\Gamma} f(z) dz = 0.$$

**Proof:** Fix  $z \in U \setminus \Gamma$ .

Apply the Cauchy integral formula to  $\zeta \mapsto (\zeta - z)f(\zeta)$  at  $\zeta = z$ :

$$0 = n(\Gamma; z)(z - z)f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{(\zeta - z)f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\Gamma} f(\zeta) d\zeta. \quad \square$$

**Remark:** Cauchy integral formula and Cauchy's theorem are equivalent logically (if you prove one the other follows).

## Definition

Cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U \subset \mathbb{C}$  are *homologous* in  $U$  if  $n(\Gamma_0; p) = n(\Gamma_1; p)$  for all  $p \in \mathbb{C} \setminus U$ .

Equivalently,  $\Gamma_0 - \Gamma_1$  is homologous to zero in  $U$ .

## Definition

Cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U \subset \mathbb{C}$  are *homologous* in  $U$  if  $n(\Gamma_0; p) = n(\Gamma_1; p)$  for all  $p \in \mathbb{C} \setminus U$ .

Equivalently,  $\Gamma_0 - \Gamma_1$  is homologous to zero in  $U$ .

## Corollary

Let  $U \subset \mathbb{C}$  be open and  $f: U \rightarrow \mathbb{C}$  holomorphic. If two cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U$  are homologous in  $U$ , then

$$\int_{\Gamma_0} f(z) dz = \int_{\Gamma_1} f(z) dz.$$

## Definition

Cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U \subset \mathbb{C}$  are *homologous* in  $U$  if  $n(\Gamma_0; p) = n(\Gamma_1; p)$  for all  $p \in \mathbb{C} \setminus U$ .

Equivalently,  $\Gamma_0 - \Gamma_1$  is homologous to zero in  $U$ .

## Corollary

Let  $U \subset \mathbb{C}$  be open and  $f: U \rightarrow \mathbb{C}$  holomorphic. If two cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U$  are homologous in  $U$ , then

$$\int_{\Gamma_0} f(z) dz = \int_{\Gamma_1} f(z) dz.$$

That is useful for computing integrals:

Given a complicated cycle, find a simple one that is homologous.

## Definition

Cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U \subset \mathbb{C}$  are *homologous* in  $U$  if  $n(\Gamma_0; p) = n(\Gamma_1; p)$  for all  $p \in \mathbb{C} \setminus U$ .

Equivalently,  $\Gamma_0 - \Gamma_1$  is homologous to zero in  $U$ .

## Corollary

Let  $U \subset \mathbb{C}$  be open and  $f: U \rightarrow \mathbb{C}$  holomorphic. If two cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U$  are homologous in  $U$ , then

$$\int_{\Gamma_0} f(z) dz = \int_{\Gamma_1} f(z) dz.$$

That is useful for computing integrals:

Given a complicated cycle, find a simple one that is homologous.

E.g. (exercise), any cycle in  $\mathbb{C} \setminus \{0\}$  is homologous to  $n\partial\mathbb{D}$  for some  $n \in \mathbb{Z}$ .

## Definition

Cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U \subset \mathbb{C}$  are *homologous* in  $U$  if  $n(\Gamma_0; p) = n(\Gamma_1; p)$  for all  $p \in \mathbb{C} \setminus U$ .

Equivalently,  $\Gamma_0 - \Gamma_1$  is homologous to zero in  $U$ .

## Corollary

Let  $U \subset \mathbb{C}$  be open and  $f: U \rightarrow \mathbb{C}$  holomorphic. If two cycles  $\Gamma_0$  and  $\Gamma_1$  in  $U$  are homologous in  $U$ , then

$$\int_{\Gamma_0} f(z) dz = \int_{\Gamma_1} f(z) dz.$$

That is useful for computing integrals:

Given a complicated cycle, find a simple one that is homologous.

E.g. (exercise), any cycle in  $\mathbb{C} \setminus \{0\}$  is homologous to  $n\partial\mathbb{D}$  for some  $n \in \mathbb{Z}$ .

**Remark:** Being “homologous” is an equivalence relation and the set of equivalence classes of cycles is an abelian group (under the cycle addition, exercise). This group is called the *first homology group* of  $U$ .