

Cultivating Complex Analysis: Residue theorem (5.3 part 1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Suppose f has an isolated singularity at p .

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n.$$

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n (z-p)^n.$$

$(z-p)^n$ has a primitive in $\Delta_r(p) \setminus \{p\}$ for all n except $n = -1$.

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n.$$

$(z-p)^n$ has a primitive in $\Delta_r(p) \setminus \{p\}$ for all n except $n = -1$.

So c_{-1} is the only thing left after integration.

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n.$$

$(z-p)^n$ has a primitive in $\Delta_r(p) \setminus \{p\}$ for all n except $n = -1$.

So c_{-1} is the only thing left after integration. It's the "residue."

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n.$$

$(z-p)^n$ has a primitive in $\Delta_r(p) \setminus \{p\}$ for all n except $n = -1$.

So c_{-1} is the only thing left after integration. It's the "residue."

Definition

Let the *residue* of f at p be

$$\operatorname{Res}(f; p) \stackrel{\text{def}}{=} c_{-1}.$$

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n.$$

$(z-p)^n$ has a primitive in $\Delta_r(p) \setminus \{p\}$ for all n except $n = -1$.

So c_{-1} is the only thing left after integration. It's the "residue."

Definition

Let the *residue* of f at p be

$$\operatorname{Res}(f; p) \stackrel{\text{def}}{=} c_{-1}.$$

For small enough $s > 0$,

$$\operatorname{Res}(f; p) = c_{-1} = \frac{1}{2\pi i} \int_{\partial \Delta_s(p)} f(z) dz.$$

Suppose f has an isolated singularity at p . Expand as a Laurent series on $\Delta_r(p) \setminus \{p\}$:

$$f(z) = \sum_{n=-\infty}^{\infty} c_n (z-p)^n.$$

$(z-p)^n$ has a primitive in $\Delta_r(p) \setminus \{p\}$ for all n except $n = -1$.

So c_{-1} is the only thing left after integration. It's the "residue."

Definition

Let the *residue* of f at p be

$$\operatorname{Res}(f; p) \stackrel{\text{def}}{=} c_{-1}.$$

For small enough $s > 0$,

$$\operatorname{Res}(f; p) = c_{-1} = \frac{1}{2\pi i} \int_{\partial \Delta_s(p)} f(z) dz.$$

By Cauchy, we can compute the integral over any cycle via the residues inside. That's the Residue theorem.

Theorem (Residue theorem)

Suppose $U \subset \mathbb{C}$ is open, $S \subset U$ is a finite subset, and Γ is a cycle in $U \setminus S$ homologous to zero in U . Suppose $f: U \setminus S \rightarrow \mathbb{C}$ is holomorphic (isolated singularities on S). Then

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = \sum_{p \in S} n(\Gamma; p) \operatorname{Res}(f; p).$$

(Recall, “homologous to zero in U ” means $n(\Gamma; z) = 0$ for all $z \in \mathbb{C} \setminus U$.)

Theorem (Residue theorem)

Suppose $U \subset \mathbb{C}$ is open, $S \subset U$ is a finite subset, and Γ is a cycle in $U \setminus S$ homologous to zero in U . Suppose $f: U \setminus S \rightarrow \mathbb{C}$ is holomorphic (isolated singularities on S). Then

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = \sum_{p \in S} n(\Gamma; p) \operatorname{Res}(f; p).$$

(Recall, “homologous to zero in U ” means $n(\Gamma; z) = 0$ for all $z \in \mathbb{C} \setminus U$.)

Proof: Write $S = \{w_1, \dots, w_{\ell}\}$.

Theorem (Residue theorem)

Suppose $U \subset \mathbb{C}$ is open, $S \subset U$ is a finite subset, and Γ is a cycle in $U \setminus S$ homologous to zero in U . Suppose $f: U \setminus S \rightarrow \mathbb{C}$ is holomorphic (isolated singularities on S). Then

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = \sum_{p \in S} n(\Gamma; p) \operatorname{Res}(f; p).$$

(Recall, “homologous to zero in U ” means $n(\Gamma; z) = 0$ for all $z \in \mathbb{C} \setminus U$.)

Proof: Write $S = \{w_1, \dots, w_{\ell}\}$.

Let r_1, \dots, r_{ℓ} be positive such that

$\overline{\Delta_{r_1}(w_1)}, \dots, \overline{\Delta_{r_{\ell}}(w_{\ell})}$ are mutually disjoint

Theorem (Residue theorem)

Suppose $U \subset \mathbb{C}$ is open, $S \subset U$ is a finite subset, and Γ is a cycle in $U \setminus S$ homologous to zero in U . Suppose $f: U \setminus S \rightarrow \mathbb{C}$ is holomorphic (isolated singularities on S). Then

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = \sum_{p \in S} n(\Gamma; p) \operatorname{Res}(f; p).$$

(Recall, “homologous to zero in U ” means $n(\Gamma; z) = 0$ for all $z \in \mathbb{C} \setminus U$.)

Proof: Write $S = \{w_1, \dots, w_\ell\}$.

Let r_1, \dots, r_ℓ be positive such that

$\overline{\Delta_{r_1}(w_1)}, \dots, \overline{\Delta_{r_\ell}(w_\ell)}$ are mutually disjoint

and $\Delta_{r_j}(w_j) \subset U$ for all j .

Theorem (Residue theorem)

Suppose $U \subset \mathbb{C}$ is open, $S \subset U$ is a finite subset, and Γ is a cycle in $U \setminus S$ homologous to zero in U . Suppose $f: U \setminus S \rightarrow \mathbb{C}$ is holomorphic (isolated singularities on S). Then

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = \sum_{p \in S} n(\Gamma; p) \operatorname{Res}(f; p).$$


(Recall, “homologous to zero in U ” means $n(\Gamma; z) = 0$ for all $z \in \mathbb{C} \setminus U$.)

Proof: Write $S = \{w_1, \dots, w_\ell\}$.

Let r_1, \dots, r_ℓ be positive such that $\overline{\Delta_{r_1}(w_1)}, \dots, \overline{\Delta_{r_\ell}(w_\ell)}$ are mutually disjoint and $\Delta_{r_j}(w_j) \subset U$ for all j .

Define the cycle

$$\Lambda = \Gamma - n(\Gamma; w_1) \partial \Delta_{r_1}(w_1) - \dots - n(\Gamma; w_\ell) \partial \Delta_{r_\ell}(w_\ell).$$



Theorem (Residue theorem)

Suppose $U \subset \mathbb{C}$ is open, $S \subset U$ is a finite subset, and Γ is a cycle in $U \setminus S$ homologous to zero in U . Suppose $f: U \setminus S \rightarrow \mathbb{C}$ is holomorphic (isolated singularities on S). Then

$$\frac{1}{2\pi i} \int_{\Gamma} f(z) dz = \sum_{p \in S} n(\Gamma; p) \operatorname{Res}(f; p).$$

(Recall, “homologous to zero in U ” means $n(\Gamma; z) = 0$ for all $z \in \mathbb{C} \setminus U$.)


Proof: Write $S = \{w_1, \dots, w_\ell\}$.

Let r_1, \dots, r_ℓ be positive such that $\overline{\Delta_{r_1}(w_1)}, \dots, \overline{\Delta_{r_\ell}(w_\ell)}$ are mutually disjoint and $\Delta_{r_j}(w_j) \subset U$ for all j .

Define the cycle

$$\Lambda = \Gamma - n(\Gamma; w_1) \partial \Delta_{r_1}(w_1) - \dots - n(\Gamma; w_\ell) \partial \Delta_{r_\ell}(w_\ell).$$

In the picture $n(\Gamma; w_1) = 1$, $n(\Gamma; w_2) = 0$, and $n(\Gamma; w_3) = 2$.



Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial \Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial \Delta_{r_\ell}(w_\ell).$

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

and for all j , $n(\partial\Delta_{r_j}(w_j); p) = 0$ (as $\overline{\Delta_{r_j}(w_j)} \subset U$).

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

and for all j , $n(\partial\Delta_{r_j}(w_j); p) = 0$ (as $\overline{\Delta_{r_j}(w_j)} \subset U$).

If $p = w_k \in S \Rightarrow n(\partial\Delta_{r_j}(w_j); p) = 0$ if $j \neq k$, and $n(\partial\Delta_{r_k}(w_k); p) = 1$.

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

and for all j , $n(\partial\Delta_{r_j}(w_j); p) = 0$ (as $\overline{\Delta_{r_j}(w_j)} \subset U$).

If $p = w_k \in S \Rightarrow n(\partial\Delta_{r_j}(w_j); p) = 0$ if $j \neq k$, and $n(\partial\Delta_{r_k}(w_k); p) = 1$.

$\Rightarrow n(\Lambda; p) = 0$ for all $p \notin U \setminus S$.

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

and for all j , $n(\partial\Delta_{r_j}(w_j); p) = 0$ (as $\overline{\Delta_{r_j}(w_j)} \subset U$).

If $p = w_k \in S \Rightarrow n(\partial\Delta_{r_j}(w_j); p) = 0$ if $j \neq k$, and $n(\partial\Delta_{r_k}(w_k); p) = 1$.

$\Rightarrow n(\Lambda; p) = 0$ for all $p \notin U \setminus S$.

We can apply the Cauchy theorem in $U \setminus S$:

$$0 = \frac{1}{2\pi i} \int_{\Lambda} f(z) dz$$

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

and for all j , $n(\partial\Delta_{r_j}(w_j); p) = 0$ (as $\overline{\Delta_{r_j}(w_j)} \subset U$).

If $p = w_k \in S \Rightarrow n(\partial\Delta_{r_j}(w_j); p) = 0$ if $j \neq k$, and $n(\partial\Delta_{r_k}(w_k); p) = 1$.

$\Rightarrow n(\Lambda; p) = 0$ for all $p \notin U \setminus S$.

We can apply the Cauchy theorem in $U \setminus S$:

$$0 = \frac{1}{2\pi i} \int_{\Lambda} f(z) dz = \frac{1}{2\pi i} \int_{\Gamma} f(z) dz - \sum_{k=1}^{\ell} n(\Gamma; w_k) \frac{1}{2\pi i} \int_{\partial\Delta_{r_k}(w_k)} f(z) dz.$$

Recall: $\Lambda = \Gamma - n(\Gamma; w_1) \partial\Delta_{r_1}(w_1) - \cdots - n(\Gamma; w_\ell) \partial\Delta_{r_\ell}(w_\ell)$.

Winding number is an integral so it is linear:

$$n(\Lambda; p) = n(\Gamma; p) - n(\Gamma; w_1) n(\partial\Delta_{r_1}(w_1); p) - \cdots - n(\Gamma; w_\ell) n(\partial\Delta_{r_\ell}(w_\ell); p).$$

If $p \notin U \Rightarrow n(\Gamma; p) = 0$ (as Γ is homologous to zero in U),

and for all j , $n(\partial\Delta_{r_j}(w_j); p) = 0$ (as $\overline{\Delta_{r_j}(w_j)} \subset U$).

If $p = w_k \in S \Rightarrow n(\partial\Delta_{r_j}(w_j); p) = 0$ if $j \neq k$, and $n(\partial\Delta_{r_k}(w_k); p) = 1$.

$\Rightarrow n(\Lambda; p) = 0$ for all $p \notin U \setminus S$.

We can apply the Cauchy theorem in $U \setminus S$:

$$0 = \frac{1}{2\pi i} \int_{\Lambda} f(z) dz = \frac{1}{2\pi i} \int_{\Gamma} f(z) dz - \sum_{k=1}^{\ell} n(\Gamma; w_k) \frac{1}{2\pi i} \int_{\partial\Delta_{r_k}(w_k)} f(z) dz.$$

Recognize the formula for c_{-1} at w_k :

$$\frac{1}{2\pi i} \int_{\partial\Delta_{r_k}(w_k)} f(z) dz = \text{Res}(f; w_k).$$