

# Cultivating Complex Analysis: Winding numbers (4.1.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Let  $\Gamma$  be a cycle, and  $p \notin \Gamma$ . Define

$$n(\Gamma; p) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz.$$

Let  $\Gamma$  be a cycle, and  $p \notin \Gamma$ . Define

$$n(\Gamma; p) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz.$$

$n(\Gamma; p)$  is called the *winding number* of  $\Gamma$  around  $p$ , or the *index* of  $\Gamma$  with respect to  $p$ .

Let  $\Gamma$  be a cycle, and  $p \notin \Gamma$ . Define

$$n(\Gamma; p) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz.$$

$n(\Gamma; p)$  is called the *winding number* of  $\Gamma$  around  $p$ , or the *index* of  $\Gamma$  with respect to  $p$ .

It is the number of times  $\Gamma$  winds around  $p$  in the counterclockwise direction.

Let  $\Gamma$  be a cycle, and  $p \notin \Gamma$ . Define

$$n(\Gamma; p) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz.$$

$n(\Gamma; p)$  is called the *winding number* of  $\Gamma$  around  $p$ , or the *index* of  $\Gamma$  with respect to  $p$ .

It is the number of times  $\Gamma$  winds around  $p$  in the counterclockwise direction.

**Example:**  $\gamma(t) = e^{it}$  for  $t \in [0, 2\pi]$  goes once around 0 in the counterclockwise direction,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz = 1.$$

Let  $\Gamma$  be a cycle, and  $p \notin \Gamma$ . Define

$$n(\Gamma; p) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz.$$

$n(\Gamma; p)$  is called the *winding number* of  $\Gamma$  around  $p$ , or the *index* of  $\Gamma$  with respect to  $p$ .

It is the number of times  $\Gamma$  winds around  $p$  in the counterclockwise direction.

**Example:**  $\gamma(t) = e^{it}$  for  $t \in [0, 2\pi]$  goes once around 0 in the counterclockwise direction,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz = 1.$$

**Example:**  $\gamma(t) = e^{i2t}$  for  $t \in [0, 2\pi]$  goes twice around 0,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz = 2.$$

Let  $\Gamma$  be a cycle, and  $p \notin \Gamma$ . Define

$$n(\Gamma; p) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz.$$

$n(\Gamma; p)$  is called the *winding number* of  $\Gamma$  around  $p$ , or the *index* of  $\Gamma$  with respect to  $p$ .

It is the number of times  $\Gamma$  winds around  $p$  in the counterclockwise direction.

**Example:**  $\gamma(t) = e^{it}$  for  $t \in [0, 2\pi]$  goes once around 0 in the counterclockwise direction,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz = 1.$$

**Example:**  $\gamma(t) = e^{i2t}$  for  $t \in [0, 2\pi]$  goes twice around 0,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz = 2.$$

**Example:**  $\gamma(t) = e^{-it}$  for  $t \in [0, 2\pi]$  goes once around in the clockwise direction,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{1}{z} dz = -1.$$

## Proposition

*Suppose  $\Gamma$  is a cycle and  $p \notin \Gamma$ . Then  $n(\Gamma; p)$  is an integer.*



## Proposition

*Suppose  $\Gamma$  is a cycle and  $p \notin \Gamma$ . Then  $n(\Gamma; p)$  is an integer.*

Idea: Follow a branch of  $\log$ , then the argument differs by an integer multiple of  $2\pi$ .

## Proposition

*Suppose  $\Gamma$  is a cycle and  $p \notin \Gamma$ . Then  $n(\Gamma; p)$  is an integer.*

Idea: Follow a branch of  $\log$ , then the argument differs by an integer multiple of  $2\pi$ .

**Proof:**  $\Gamma$  is a “sum” of closed paths, so WLOG consider a closed piecewise- $C^1$  path  $\gamma: [0, 1] \rightarrow \mathbb{C}$ .


## Proposition

Suppose  $\Gamma$  is a cycle and  $p \notin \Gamma$ . Then  $n(\Gamma; p)$  is an integer.

Idea: Follow a branch of log, then the argument differs by an integer multiple of  $2\pi$ .

**Proof:**  $\Gamma$  is a “sum” of closed paths, so WLOG consider a closed piecewise- $C^1$  path  $\gamma: [0, 1] \rightarrow \mathbb{C}$ .

$\gamma$  can be covered by finitely many discs  $D_1, \dots, D_n$  none of which contain  $p$ .




## Proposition

Suppose  $\Gamma$  is a cycle and  $p \notin \Gamma$ . Then  $n(\Gamma; p)$  is an integer.

Idea: Follow a branch of log, then the argument differs by an integer multiple of  $2\pi$ .

**Proof:**  $\Gamma$  is a “sum” of closed paths, so WLOG consider a closed piecewise- $C^1$  path  $\gamma: [0, 1] \rightarrow \mathbb{C}$ .

$\gamma$  can be covered by finitely many discs  $D_1, \dots, D_n$  none of which contain  $p$ .



(cover the whole closed curve, of course)

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ .



The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$n(\gamma; p) = \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz$$

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$n(\gamma; p) = \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt$$

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$n(\gamma; p) = \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt = \frac{1}{2\pi i} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \frac{\gamma'(t)}{\gamma(t) - p} dt$$

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$\begin{aligned} n(\gamma; p) &= \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt = \frac{1}{2\pi i} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \frac{\gamma'(t)}{\gamma(t) - p} dt \\ &= \frac{1}{2\pi i} \sum_{j=1}^n L_j(\gamma(t_j)) - L_j(\gamma(t_{j-1})) \end{aligned}$$

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$\begin{aligned} n(\gamma; p) &= \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt = \frac{1}{2\pi i} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \frac{\gamma'(t)}{\gamma(t) - p} dt \\ &= \frac{1}{2\pi i} \sum_{j=1}^n L_j(\gamma(t_j)) - L_j(\gamma(t_{j-1})) = \frac{1}{2\pi i} (L_n(z_0) - L_1(z_0)). \end{aligned}$$

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$\begin{aligned} n(\gamma; p) &= \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt = \frac{1}{2\pi i} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \frac{\gamma'(t)}{\gamma(t) - p} dt \\ &= \frac{1}{2\pi i} \sum_{j=1}^n L_j(\gamma(t_j)) - L_j(\gamma(t_{j-1})) = \frac{1}{2\pi i} (L_n(z_0) - L_1(z_0)). \end{aligned}$$

$L_n$  and  $L_1$  are branches of  $\log$ ,

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$\begin{aligned} n(\gamma; p) &= \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt = \frac{1}{2\pi i} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \frac{\gamma'(t)}{\gamma(t) - p} dt \\ &= \frac{1}{2\pi i} \sum_{j=1}^n L_j(\gamma(t_j)) - L_j(\gamma(t_{j-1})) = \frac{1}{2\pi i} (L_n(z_0) - L_1(z_0)). \end{aligned}$$

$L_n$  and  $L_1$  are branches of  $\log$ ,

each is  $\log|z_0| + i \arg z_0$  for some value of  $\arg$ ,

The discs  $D_1, \dots, D_n$  cover  $\gamma$ .

And the discs can be chosen (exercise) so that there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_n = 1$ , where  $\gamma([t_{j-1}, t_j]) \subset D_j$  for every  $j$ .

Each  $D_j$  is star-like and  $p \notin D_j \Rightarrow \exists$  a branch  $L_j$  of  $\log(z - p)$  on each  $D_j$ , such that

$$L_j(\gamma(t_j)) = L_{j+1}(\gamma(t_j)) \quad (L_1 \text{ is an arbitrary branch}).$$

Call  $z_0 = \gamma(0) = \gamma(1)$ . So

$$\begin{aligned} n(\gamma; p) &= \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - p} dz = \frac{1}{2\pi i} \int_0^1 \frac{\gamma'(t)}{\gamma(t) - p} dt = \frac{1}{2\pi i} \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \frac{\gamma'(t)}{\gamma(t) - p} dt \\ &= \frac{1}{2\pi i} \sum_{j=1}^n L_j(\gamma(t_j)) - L_j(\gamma(t_{j-1})) = \frac{1}{2\pi i} (L_n(z_0) - L_1(z_0)). \end{aligned}$$

$L_n$  and  $L_1$  are branches of  $\log$ ,  
each is  $\log|z_0| + i \arg z_0$  for some value of  $\arg$ ,  
their difference is  $2\pi ki$  for some  $k \in \mathbb{Z}$ .





## Proposition

*Given a cycle  $\Gamma$ , the function  $z \mapsto n(\Gamma; z)$  is constant on the topological components of  $\mathbb{C} \setminus \Gamma$ .*

## Proposition

*Given a cycle  $\Gamma$ , the function  $z \mapsto n(\Gamma; z)$  is constant on the topological components of  $\mathbb{C} \setminus \Gamma$ . Furthermore,  $n(\Gamma; z) = 0$  for  $z$  on the unbounded component of  $\mathbb{C} \setminus \Gamma$ .*

## Proposition

*Given a cycle  $\Gamma$ , the function  $z \mapsto n(\Gamma; z)$  is constant on the topological components of  $\mathbb{C} \setminus \Gamma$ . Furthermore,  $n(\Gamma; z) = 0$  for  $z$  on the unbounded component of  $\mathbb{C} \setminus \Gamma$ .*


As  $\Gamma$  is compact, is a unique unbounded component of  $\mathbb{C} \setminus \Gamma$ .

## Proposition

Given a cycle  $\Gamma$ , the function  $z \mapsto n(\Gamma; z)$  is constant on the topological components of  $\mathbb{C} \setminus \Gamma$ . Furthermore,  $n(\Gamma; z) = 0$  for  $z$  on the unbounded component of  $\mathbb{C} \setminus \Gamma$ .

As  $\Gamma$  is compact, there is a unique unbounded component of  $\mathbb{C} \setminus \Gamma$ .

**Example:**



**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z - p| \geq d/2$  for  $z \in \Gamma$ .

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z - p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ).



**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)|$$

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z - p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z - p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right|$$

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z - p_0||z - p|} |dz|$$

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z - p_0||z - p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z-p_0)(z-p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z-p_0||z-p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z - p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z - p_0||z - p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

Being continuous and integer-valued, it is constant on every component of  $\mathbb{C} \setminus \Gamma$ .

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z-p_0)(z-p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z-p_0||z-p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

Being continuous and integer-valued, it is constant on every component of  $\mathbb{C} \setminus \Gamma$ .

For  $p \in \mathbb{C} \setminus \Gamma$ ,

$$|n(\Gamma; p)|$$

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z-p_0)(z-p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z-p_0||z-p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

Being continuous and integer-valued, it is constant on every component of  $\mathbb{C} \setminus \Gamma$ .

For  $p \in \mathbb{C} \setminus \Gamma$ ,

$$|n(\Gamma; p)| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{1}{|z-p|} |dz|$$



**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z - p_0||z - p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

Being continuous and integer-valued, it is constant on every component of  $\mathbb{C} \setminus \Gamma$ .

For  $p \in \mathbb{C} \setminus \Gamma$ ,

$$|n(\Gamma; p)| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{1}{|z - p|} |dz| \leq \frac{1}{2\pi} \frac{\ell}{d(p, \Gamma)}.$$

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z-p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z - p_0||z - p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

Being continuous and integer-valued, it is constant on every component of  $\mathbb{C} \setminus \Gamma$ .

For  $p \in \mathbb{C} \setminus \Gamma$ ,

$$|n(\Gamma; p)| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{1}{|z-p|} |dz| \leq \frac{1}{2\pi} \frac{\ell}{d(p, \Gamma)}.$$

On the unbounded component, there are  $p$  with arbitrarily large  $d(p, \Gamma)$ .

**Proof:** We start by showing that  $p \mapsto n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$  is continuous on  $\mathbb{C} \setminus \Gamma$ .

Fix  $p_0 \in \mathbb{C} \setminus \Gamma$ , and let  $d = d(p_0, \Gamma)$  be the distance from  $p_0$  to  $\Gamma$  ( $d > 0$  as  $\Gamma$  is compact).

If  $p \in \Delta_{d/2}(p_0)$ , then  $|z - p| \geq d/2$  for  $z \in \Gamma$ .

Let  $\ell = \int_{\Gamma} |dz|$  (length of  $\Gamma$ ). Then

$$|n(\Gamma; p_0) - n(\Gamma; p)| = \left| \frac{1}{2\pi i} \int_{\Gamma} \frac{p_0 - p}{(z - p_0)(z - p)} dz \right| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{|p_0 - p|}{|z - p_0||z - p|} |dz| \leq \frac{\ell}{\pi d^2} |p_0 - p|.$$

So,  $p \mapsto n(\Gamma; p)$  is a continuous.

Being continuous and integer-valued, it is constant on every component of  $\mathbb{C} \setminus \Gamma$ .

For  $p \in \mathbb{C} \setminus \Gamma$ ,

$$|n(\Gamma; p)| \leq \frac{1}{2\pi} \int_{\Gamma} \frac{1}{|z - p|} |dz| \leq \frac{1}{2\pi} \frac{\ell}{d(p, \Gamma)}.$$

On the unbounded component, there are  $p$  with arbitrarily large  $d(p, \Gamma)$ .

So  $n(\Gamma; p) = 0$  on this component.



An exercise (direct computation) which we will often use is:

An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

$n(\partial\Delta_r(p); z) = 1$  if  $z \in \Delta_r(p)$ .

An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

$n(\partial\Delta_r(p); z) = 1$  if  $z \in \Delta_r(p)$ .

**Exercise:** Compute the winding numbers in an “annulus”:

An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

$n(\partial\Delta_r(p); z) = 1$  if  $z \in \Delta_r(p)$ .

**Exercise:** Compute the winding numbers in an “annulus”:

Suppose  $0 < r_1 < r_2 < \infty$  and  $\Gamma = \partial\Delta_{r_2}(p) - \partial\Delta_{r_1}(p)$ .



An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

$n(\partial\Delta_r(p); z) = 1$  if  $z \in \Delta_r(p)$ .

**Exercise:** Compute the winding numbers in an “annulus”:

Suppose  $0 < r_1 < r_2 < \infty$  and  $\Gamma = \partial\Delta_{r_2}(p) - \partial\Delta_{r_1}(p)$ .

Then

$n(\Gamma; z) = 0$  if  $|z - p| < r_1$ ,

An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

$n(\partial\Delta_r(p); z) = 1$  if  $z \in \Delta_r(p)$ .

**Exercise:** Compute the winding numbers in an “annulus”:

Suppose  $0 < r_1 < r_2 < \infty$  and  $\Gamma = \partial\Delta_{r_2}(p) - \partial\Delta_{r_1}(p)$ .

Then

$n(\Gamma; z) = 0$  if  $|z - p| < r_1$ ,

$n(\Gamma; z) = 1$  if  $r_1 < |z - p| < r_2$ ,

An exercise (direct computation) which we will often use is:

**Exercise:**

$n(\partial\Delta_r(p); z) = 0$  if  $z \notin \overline{\Delta_r(p)}$ , and

$n(\partial\Delta_r(p); z) = 1$  if  $z \in \Delta_r(p)$ .

**Exercise:** Compute the winding numbers in an “annulus”:

Suppose  $0 < r_1 < r_2 < \infty$  and  $\Gamma = \partial\Delta_{r_2}(p) - \partial\Delta_{r_1}(p)$ .

Then

$n(\Gamma; z) = 0$  if  $|z - p| < r_1$ ,

$n(\Gamma; z) = 1$  if  $r_1 < |z - p| < r_2$ ,

$n(\Gamma; z) = 0$  if  $r_2 < |z - p|$ .