

# Cultivating Complex Analysis: Holomorphic functions via integrals (3.4.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Lemma

*Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic.*

## Lemma

*Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then*

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

## Lemma

*Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then*

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

## Lemma

*Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then*

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual).

## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual). Then

$$\int_{\partial T} h(z) dz$$

## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual). Then

$$\int_{\partial T} h(z) dz = \int_{\partial T} \int_0^1 \psi(z, t) dt dz$$



## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual). Then

$$\int_{\partial T} h(z) dz = \int_{\partial T} \int_0^1 \psi(z, t) dt dz = \int_0^1 \int_{\partial T} \psi(z, t) dz dt$$

## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual). Then

$$\int_{\partial T} h(z) dz = \int_{\partial T} \int_0^1 \psi(z, t) dt dz = \int_0^1 \int_{\partial T} \psi(z, t) dz dt = \int_0^1 0 dt = 0.$$

## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual). Then

$$\int_{\partial T} h(z) dz = \int_{\partial T} \int_0^1 \psi(z, t) dt dz = \int_0^1 \int_{\partial T} \psi(z, t) dz dt = \int_0^1 0 dt = 0.$$

Fubini applies as the integrand is continuous if we think of each leg of  $\partial T$  separately.

## Lemma

Suppose  $U \subset \mathbb{C}$  is open, and  $\psi: U \times [0, 1] \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $t \in [0, 1]$ , the function  $z \mapsto \psi(z, t)$  is holomorphic. Then

$$h(z) = \int_0^1 \psi(z, t) dt \quad \text{is a holomorphic function on } U.$$

There are two common proofs of this kind of result.

**Proof A:** Morera together with Fubini and Cauchy–Goursat.

Let  $T \subset U$  be a triangle (solid triangle as usual). Then

$$\int_{\partial T} h(z) dz = \int_{\partial T} \int_0^1 \psi(z, t) dt dz = \int_0^1 \int_{\partial T} \psi(z, t) dz dt = \int_0^1 0 dt = 0.$$

Fubini applies as the integrand is continuous if we think of each leg of  $\partial T$  separately.

$h(z)$  is holomorphic by Morera.



**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)]$$

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt$$

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt$$



**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

We are really passing the partial derivatives in  $x$  and  $y$  (since  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$ ) under the integral via the Leibniz integral rule.

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

We are really passing the partial derivatives in  $x$  and  $y$  (since  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$ ) under the integral via the Leibniz integral rule.

There is a technicality! Could we apply Leibniz?

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

We are really passing the partial derivatives in  $x$  and  $y$  (since  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$ ) under the integral via the Leibniz integral rule.

There is a technicality! Could we apply Leibniz?

We only assumed  $z \mapsto \psi(z, t)$  is holomorphic for all  $t$ .

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

We are really passing the partial derivatives in  $x$  and  $y$  (since  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$ ) under the integral via the Leibniz integral rule.

There is a technicality! Could we apply Leibniz?

We only assumed  $z \mapsto \psi(z, t)$  is holomorphic for all  $t$ .

We did *not* assume  $\frac{\partial \psi}{\partial x}$  and  $\frac{\partial \psi}{\partial y}$  (components of  $\frac{\partial \psi}{\partial \bar{z}}$ ) were continuous functions on  $U \times [0, 1]$ .  
If we did, we would be done (that's what the standard Leibniz needs).

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

We are really passing the partial derivatives in  $x$  and  $y$  (since  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$ ) under the integral via the Leibniz integral rule.

There is a technicality! Could we apply Leibniz?

We only assumed  $z \mapsto \psi(z, t)$  is holomorphic for all  $t$ .

We did *not* assume  $\frac{\partial \psi}{\partial x}$  and  $\frac{\partial \psi}{\partial y}$  (components of  $\frac{\partial \psi}{\partial \bar{z}}$ ) were continuous functions on  $U \times [0, 1]$ . If we did, we would be done (that's what the standard Leibniz needs).

By an exercise we mentioned previously (using Cauchy's integral formula for derivatives): If  $z \mapsto \psi(z, t)$  is holomorphic for all  $t$  (and  $\psi$  continuous), then  $\frac{\partial \psi}{\partial x}$  and  $\frac{\partial \psi}{\partial y}$  are continuous.

**Proof B:** Apply Wirtinger derivatives and differentiate under the integral:

$$\frac{\partial}{\partial \bar{z}} [h(z)] = \frac{\partial}{\partial \bar{z}} \int_0^1 \psi(z, t) dt = \int_0^1 \frac{\partial}{\partial \bar{z}} [\psi(z, t)] dt = \int_0^1 0 dt = 0.$$

We are really passing the partial derivatives in  $x$  and  $y$  (since  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{i}{2} \frac{\partial}{\partial y}$ ) under the integral via the Leibniz integral rule.

There is a technicality! Could we apply Leibniz?

We only assumed  $z \mapsto \psi(z, t)$  is holomorphic for all  $t$ .

We did *not* assume  $\frac{\partial \psi}{\partial x}$  and  $\frac{\partial \psi}{\partial y}$  (components of  $\frac{\partial \psi}{\partial \bar{z}}$ ) were continuous functions on  $U \times [0, 1]$ . If we did, we would be done (that's what the standard Leibniz needs).

By an exercise we mentioned previously (using Cauchy's integral formula for derivatives): If  $z \mapsto \psi(z, t)$  is holomorphic for all  $t$  (and  $\psi$  continuous), then  $\frac{\partial \psi}{\partial x}$  and  $\frac{\partial \psi}{\partial y}$  are continuous.

OK, now we are done. □

## Corollary

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a chain, and  $\psi: U \times \Gamma \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $w \in \Gamma$ , the function  $z \mapsto \psi(z, w)$  is holomorphic. Then

$$h(z) = \int_{\Gamma} \psi(z, w) dw \quad \text{is a holomorphic function on } U.$$



## Corollary

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a chain, and  $\psi: U \times \Gamma \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $w \in \Gamma$ , the function  $z \mapsto \psi(z, w)$  is holomorphic. Then

$$h(z) = \int_{\Gamma} \psi(z, w) dw \quad \text{is a holomorphic function on } U.$$

For a continuous  $f: \partial\Delta_r(p) \rightarrow \mathbb{C}$ , define the *Cauchy transform*  $Cf: \Delta_r(p) \rightarrow \mathbb{C}$  by

$$Cf(z) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

## Corollary

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a chain, and  $\psi: U \times \Gamma \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $w \in \Gamma$ , the function  $z \mapsto \psi(z, w)$  is holomorphic. Then

$$h(z) = \int_{\Gamma} \psi(z, w) dw \quad \text{is a holomorphic function on } U.$$

For a continuous  $f: \partial\Delta_r(p) \rightarrow \mathbb{C}$ , define the Cauchy transform  $Cf: \Delta_r(p) \rightarrow \mathbb{C}$  by

$$Cf(z) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

## Corollary

For a continuous  $f: \partial\Delta_r(p) \rightarrow \mathbb{C}$ , the Cauchy transform  $Cf: \Delta_r(p) \rightarrow \mathbb{C}$  is holomorphic.

## Corollary

Suppose  $U \subset \mathbb{C}$  is open,  $\Gamma$  is a chain, and  $\psi: U \times \Gamma \rightarrow \mathbb{C}$  is a continuous function such that for each fixed  $w \in \Gamma$ , the function  $z \mapsto \psi(z, w)$  is holomorphic. Then

$$h(z) = \int_{\Gamma} \psi(z, w) dw \quad \text{is a holomorphic function on } U.$$

For a continuous  $f: \partial\Delta_r(p) \rightarrow \mathbb{C}$ , define the Cauchy transform  $Cf: \Delta_r(p) \rightarrow \mathbb{C}$  by

$$Cf(z) \stackrel{\text{def}}{=} \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

## Corollary

For a continuous  $f: \partial\Delta_r(p) \rightarrow \mathbb{C}$ , the Cauchy transform  $Cf: \Delta_r(p) \rightarrow \mathbb{C}$  is holomorphic.

For a random continuous  $f$ ,  $Cf$  may not tend to  $f$  as we approach  $\partial\Delta_r(p)$ .

The corollary gives a converse to Cauchy's formula:

The corollary gives a converse to Cauchy's formula:

If  $f: \overline{\Delta_r(p)} \rightarrow \mathbb{C}$  is continuous and

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in \Delta_r(p),$$

The corollary gives a converse to Cauchy's formula:

If  $f: \overline{\Delta_r(p)} \rightarrow \mathbb{C}$  is continuous and

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in \Delta_r(p),$$

then  $f|_{\Delta_r(p)}$  is holomorphic.