

Cultivating Complex Analysis: The complex numbers as the plane (1.1.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Modern mathematics takes a false statement such as

“All polynomials have a root.”

Modern mathematics takes a false statement such as

"All polynomials have a root."

and redefines what it means to have a *root* to make the statement true.

Modern mathematics takes a false statement such as

“All polynomials have a root.”

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with $\mathbb{N} = \{1, 2, 3, \dots\}$, and arrive at \mathbb{Z} to solve something like $n + 2 = 1$.

Modern mathematics takes a false statement such as

“All polynomials have a root.”

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with $\mathbb{N} = \{1, 2, 3, \dots\}$, and arrive at \mathbb{Z} to solve something like $n + 2 = 1$.

Start with \mathbb{Z} and arrive at \mathbb{Q} to solve $2x = 1$.

Modern mathematics takes a false statement such as

“All polynomials have a root.”

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with $\mathbb{N} = \{1, 2, 3, \dots\}$, and arrive at \mathbb{Z} to solve something like $n + 2 = 1$.

Start with \mathbb{Z} and arrive at \mathbb{Q} to solve $2x = 1$.

Start with \mathbb{Q} and arrive at \mathbb{R} to be able to take limits.

Modern mathematics takes a false statement such as

“All polynomials have a root.”

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with $\mathbb{N} = \{1, 2, 3, \dots\}$, and arrive at \mathbb{Z} to solve something like $n + 2 = 1$.

Start with \mathbb{Z} and arrive at \mathbb{Q} to solve $2x = 1$.

Start with \mathbb{Q} and arrive at \mathbb{R} to be able to take limits.

Start with \mathbb{R} and arrive at \mathbb{C} to be able to solve $z^2 + 1 = 0$.

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Mainly once we ask for the complex derivative we get lots of odd statements:

If you can differentiate once, you can differentiate twice.

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Mainly once we ask for the complex derivative we get lots of odd statements:

If you can differentiate once, you can differentiate twice.

Every function acts sort of like a linear function.

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Mainly once we ask for the complex derivative we get lots of odd statements:

If you can differentiate once, you can differentiate twice.

Every function acts sort of like a linear function.

If all derivatives are zero at a point, the function is constant.

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Mainly once we ask for the complex derivative we get lots of odd statements:

If you can differentiate once, you can differentiate twice.

Every function acts sort of like a linear function.

If all derivatives are zero at a point, the function is constant.

etc.

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Mainly once we ask for the complex derivative we get lots of odd statements:

If you can differentiate once, you can differentiate twice.

Every function acts sort of like a linear function.

If all derivatives are zero at a point, the function is constant.

etc.

A graduate real analysis course crushes the student's dreams.

Interestingly, analysis with \mathbb{C} is very different to analysis with \mathbb{R} .

Mainly once we ask for the complex derivative we get lots of odd statements:

If you can differentiate once, you can differentiate twice.

Every function acts sort of like a linear function.

If all derivatives are zero at a point, the function is constant.

etc.

A graduate real analysis course crushes the student's dreams.

A graduate complex analysis course fills the student with unrealistic optimism.

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + c, b + d),$$
$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$\begin{aligned}(a, b) + (c, d) &\stackrel{\text{def}}{=} (a + c, b + d), \\ (a, b)(c, d) &\stackrel{\text{def}}{=} (ac - bd, bc + da).\end{aligned}$$

This makes \mathbb{C} into a field (exercise).

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + c, b + d),$$
$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

This makes \mathbb{C} into a field (exercise).

$\mathbb{R} \subset \mathbb{C}$ by identifying $x \in \mathbb{R}$ with $(x, 0)$.

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$\begin{aligned}(a, b) + (c, d) &\stackrel{\text{def}}{=} (a + b, c + d), \\ (a, b)(c, d) &\stackrel{\text{def}}{=} (ac - bd, bc + da).\end{aligned}$$

This makes \mathbb{C} into a field (exercise).

$\mathbb{R} \subset \mathbb{C}$ by identifying $x \in \mathbb{R}$ with $(x, 0)$.

Imaginary unit:

$$i \stackrel{\text{def}}{=} (0, 1)$$

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + b, c + d),$$
$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

This makes \mathbb{C} into a field (exercise).

$\mathbb{R} \subset \mathbb{C}$ by identifying $x \in \mathbb{R}$ with $(x, 0)$.

Imaginary unit:

$$i \stackrel{\text{def}}{=} (0, 1)$$

$i^2 = -1$ so $z^2 + 1 = 0$ has the two solutions i and $-i$.

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

The evil twin of $z = x + iy$ is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

The evil twin of $z = x + iy$ is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the x and y :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

The evil twin of $z = x + iy$ is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the x and y :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

The evil twin of $z = x + iy$ is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the x and y :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

An expression in x, y can be written in terms of z, \bar{z} and vice versa:

$$x^3 + y^3 + 3ixy = \left(\frac{z + \bar{z}}{2}\right)^3 + \left(\frac{z - \bar{z}}{2i}\right)^3 + 3i\left(\frac{z + \bar{z}}{2}\right)\left(\frac{z - \bar{z}}{2i}\right),$$

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

The evil twin of $z = x + iy$ is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the x and y :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

An expression in x, y can be written in terms of z, \bar{z} and vice versa:

$$x^3 + y^3 + 3ixy = \left(\frac{z + \bar{z}}{2}\right)^3 + \left(\frac{z - \bar{z}}{2i}\right)^3 + 3i\left(\frac{z + \bar{z}}{2}\right)\left(\frac{z - \bar{z}}{2i}\right),$$

or

$$z^2 - i\bar{z}^2 + z\bar{z} = (x + iy)^2 - i(x - iy)^2 + (x + iy)(x - iy).$$

$(x, y) = x + iy$, so use $x + iy$ from now on (*cartesian form*).

The evil twin of $z = x + iy$ is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the x and y :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

An expression in x, y can be written in terms of z, \bar{z} and vice versa:

$$x^3 + y^3 + 3ixy = \left(\frac{z + \bar{z}}{2}\right)^3 + \left(\frac{z - \bar{z}}{2i}\right)^3 + 3i\left(\frac{z + \bar{z}}{2}\right)\left(\frac{z - \bar{z}}{2i}\right),$$

or

$$z^2 - i\bar{z}^2 + z\bar{z} = (x + iy)^2 - i(x - iy)^2 + (x + iy)(x - iy).$$

Almost looks as if z and \bar{z} were independent variables.