

Cultivating Complex Analysis: The open mapping theorem (5.5)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

A continuous function from \mathbb{R}^2 to \mathbb{R}^2 can do all sorts of things to the topology.

A continuous function from \mathbb{R}^2 to \mathbb{R}^2 can do all sorts of things to the topology.

$(x, y) \mapsto (x, xy)$, takes all of \mathbb{R}^2 , which is both open and closed, to the set $\{(x, y) : x \neq 0 \text{ or } y = 0\}$, which is neither open nor closed.

A continuous function from \mathbb{R}^2 to \mathbb{R}^2 can do all sorts of things to the topology.

$(x, y) \mapsto (x, xy)$, takes all of \mathbb{R}^2 , which is both open and closed, to the set $\{(x, y) : x \neq 0 \text{ or } y = 0\}$, which is neither open nor closed.

Holomorphic functions are always nice to your topology.

A continuous function from \mathbb{R}^2 to \mathbb{R}^2 can do all sorts of things to the topology.

$(x, y) \mapsto (x, xy)$, takes all of \mathbb{R}^2 , which is both open and closed, to the set $\{(x, y) : x \neq 0 \text{ or } y = 0\}$, which is neither open nor closed.

Holomorphic functions are always nice to your topology.

For a continuous map, $f^{-1}(V)$ is open whenever V is.

A continuous function from \mathbb{R}^2 to \mathbb{R}^2 can do all sorts of things to the topology.

$(x, y) \mapsto (x, xy)$, takes all of \mathbb{R}^2 , which is both open and closed, to the set $\{(x, y) : x \neq 0 \text{ or } y = 0\}$, which is neither open nor closed.

Holomorphic functions are always nice to your topology.

For a continuous map, $f^{-1}(V)$ is open whenever V is.

For a holomorphic map, $f(V)$ is open whenever V is, unless f is constant.

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Proof: Suppose f is not constant. As U is connected, f is not constant near every point.

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Proof: Suppose f is not constant. As U is connected, f is not constant near every point.

Given $p \in V$, $\exists \overline{\Delta_r(p)} \subset V$ and a $\delta > 0$ such that $|f(z) - f(p)| > \delta$ for all $z \in \partial\Delta_r(p)$.

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Proof: Suppose f is not constant. As U is connected, f is not constant near every point.

Given $p \in V$, $\exists \overline{\Delta_r(p)} \subset V$ and a $\delta > 0$ such that $|f(z) - f(p)| > \delta$ for all $z \in \partial\Delta_r(p)$.

$z \mapsto f(z) - f(p)$ has at least one zero in $\Delta_r(p)$.

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Proof: Suppose f is not constant. As U is connected, f is not constant near every point.

Given $p \in V$, $\exists \overline{\Delta_r(p)} \subset V$ and a $\delta > 0$ such that $|f(z) - f(p)| > \delta$ for all $z \in \partial\Delta_r(p)$.

$z \mapsto f(z) - f(p)$ has at least one zero in $\Delta_r(p)$.

Take $w \in \Delta_\delta(f(p))$. For all $z \in \partial\Delta_r(p)$,

$$|(f(z) - w) - (f(z) - f(p))| = |f(p) - w| < \delta < |f(z) - f(p)|.$$

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Proof: Suppose f is not constant. As U is connected, f is not constant near every point.

Given $p \in V$, $\exists \overline{\Delta_r(p)} \subset V$ and a $\delta > 0$ such that $|f(z) - f(p)| > \delta$ for all $z \in \partial\Delta_r(p)$.

$z \mapsto f(z) - f(p)$ has at least one zero in $\Delta_r(p)$.

Take $w \in \Delta_\delta(f(p))$. For all $z \in \partial\Delta_r(p)$,

$$|(f(z) - w) - (f(z) - f(p))| = |f(p) - w| < \delta < |f(z) - f(p)|.$$

By Rouché, $z \mapsto f(z) - w$ has at least one zero in $\Delta_r(p)$.

Theorem (Open mapping)

Let $U \subset \mathbb{C}$ be a domain and $f: U \rightarrow \mathbb{C}$ be holomorphic and nonconstant. Then $f(V)$ is an open set for every open set $V \subset U$.

Proof: Suppose f is not constant. As U is connected, f is not constant near every point.

Given $p \in V$, $\exists \overline{\Delta_r(p)} \subset V$ and a $\delta > 0$ such that $|f(z) - f(p)| > \delta$ for all $z \in \partial\Delta_r(p)$.

$z \mapsto f(z) - f(p)$ has at least one zero in $\Delta_r(p)$.

Take $w \in \Delta_\delta(f(p))$. For all $z \in \partial\Delta_r(p)$,

$$|(f(z) - w) - (f(z) - f(p))| = |f(p) - w| < \delta < |f(z) - f(p)|.$$

By Rouché, $z \mapsto f(z) - w$ has at least one zero in $\Delta_r(p)$. So

$$\Delta_\delta(f(p)) \subset f(\Delta_r(p)). \quad \square$$

The open mapping theorem is a stronger version of the maximum principle.

The open mapping theorem is a stronger version of the maximum principle.
 $f(p)$ is in the interior of $f(V)$ for any open neighborhood V of p .

The open mapping theorem is a stronger version of the maximum principle.

$f(p)$ is in the interior of $f(V)$ for any open neighborhood V of p .

So $|f(z)|$ cannot achieve a maximum at p .

The open mapping theorem is a stronger version of the maximum principle.

$f(p)$ is in the interior of $f(V)$ for any open neighborhood V of p .

So $|f(z)|$ cannot achieve a maximum at p .

The proof gives the more explicit:

$|f(z) - f(p)| > \delta$ for $z \in \partial\Delta_r(p)$, then $\Delta_\delta(f(p)) \subset f(\Delta_r(p))$.

Exercise: Suppose $U, V \subset \mathbb{C}$ are open and $f: U \rightarrow \mathbb{C}$ is holomorphic and bijective. Prove that $f^{-1}: V \rightarrow U$ is continuous.

Exercise: Suppose $U, V \subset \mathbb{C}$ are open and $f: U \rightarrow \mathbb{C}$ is holomorphic and bijective. Prove that $f^{-1}: V \rightarrow U$ is continuous.

We'll prove in the next lecture that f^{-1} is, in fact, holomorphic.

Exercise: Suppose $U, V \subset \mathbb{C}$ are open and $f: U \rightarrow \mathbb{C}$ is holomorphic and bijective. Prove that $f^{-1}: V \rightarrow U$ is continuous.

We'll prove in the next lecture that f^{-1} is, in fact, holomorphic.

Typical application/exercise of the open mapping theorem is something like:

Exercise: Let $U \subset \mathbb{C}$ be a domain and let $f: U \rightarrow \mathbb{C}$ be holomorphic. Prove that if $(\operatorname{Im} f(z))^2 - (\operatorname{Re} f(z))^2 = 1$ for all $z \in U$, then f is constant.