

# Cultivating Complex Analysis: The complex numbers as the plane (1.1.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

*Modern mathematics* takes a false statement such as

*“All polynomials have a root.”*

*Modern mathematics* takes a false statement such as

*“All polynomials have a root.”*

and redefines what it means to have a *root* to make the statement true.

*Modern mathematics* takes a false statement such as

*“All polynomials have a root.”*

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with  $\mathbb{N} = \{1, 2, 3, \dots\}$ , and arrive at  $\mathbb{Z}$  to solve something like  $n + 2 = 1$ .

*Modern mathematics* takes a false statement such as

*“All polynomials have a root.”*

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with  $\mathbb{N} = \{1, 2, 3, \dots\}$ , and arrive at  $\mathbb{Z}$  to solve something like  $n + 2 = 1$ .

Start with  $\mathbb{Z}$  and arrive at  $\mathbb{Q}$  to solve  $2x = 1$ .

*Modern mathematics* takes a false statement such as

*“All polynomials have a root.”*

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with  $\mathbb{N} = \{1, 2, 3, \dots\}$ , and arrive at  $\mathbb{Z}$  to solve something like  $n + 2 = 1$ .

Start with  $\mathbb{Z}$  and arrive at  $\mathbb{Q}$  to solve  $2x = 1$ .

Start with  $\mathbb{Q}$  and arrive at  $\mathbb{R}$  to be able to take limits.

*Modern mathematics* takes a false statement such as

*“All polynomials have a root.”*

and redefines what it means to have a *root* to make the statement true.

Examples:

Start with  $\mathbb{N} = \{1, 2, 3, \dots\}$ , and arrive at  $\mathbb{Z}$  to solve something like  $n + 2 = 1$ .

Start with  $\mathbb{Z}$  and arrive at  $\mathbb{Q}$  to solve  $2x = 1$ .

Start with  $\mathbb{Q}$  and arrive at  $\mathbb{R}$  to be able to take limits.

Start with  $\mathbb{R}$  and arrive at  $\mathbb{C}$  to be able to solve  $z^2 + 1 = 0$ .

Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .



Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .

Mainly once we ask for the complex derivative we get lots of odd statements:

*If you can differentiate once, you can differentiate twice.*

Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .

Mainly once we ask for the complex derivative we get lots of odd statements:

*If you can differentiate once, you can differentiate twice.*

*Every function acts sort of like a linear function.*

Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .

Mainly once we ask for the complex derivative we get lots of odd statements:

*If you can differentiate once, you can differentiate twice.*

*Every function acts sort of like a linear function.*

*If all derivatives are zero at a point, the function is constant.*

Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .

Mainly once we ask for the complex derivative we get lots of odd statements:

*If you can differentiate once, you can differentiate twice.*

*Every function acts sort of like a linear function.*

*If all derivatives are zero at a point, the function is constant.*

etc.

Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .

Mainly once we ask for the complex derivative we get lots of odd statements:

*If you can differentiate once, you can differentiate twice.*

*Every function acts sort of like a linear function.*

*If all derivatives are zero at a point, the function is constant.*

etc.

A graduate real analysis course crushes the student's dreams.

Interestingly, analysis with  $\mathbb{C}$  is very different to analysis with  $\mathbb{R}$ .

Mainly once we ask for the complex derivative we get lots of odd statements:

*If you can differentiate once, you can differentiate twice.*

*Every function acts sort of like a linear function.*

*If all derivatives are zero at a point, the function is constant.*

etc.

A graduate real analysis course crushes the student's dreams.

A graduate complex analysis course fills the student with unrealistic optimism.

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + c, b + d),$$

$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + c, b + d),$$
$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

This makes  $\mathbb{C}$  into a field (exercise).



Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$\begin{aligned}(a, b) + (c, d) &\stackrel{\text{def}}{=} (a + c, b + d), \\ (a, b)(c, d) &\stackrel{\text{def}}{=} (ac - bd, bc + da).\end{aligned}$$

This makes  $\mathbb{C}$  into a field (exercise).

$\mathbb{R} \subset \mathbb{C}$  by identifying  $x \in \mathbb{R}$  with  $(x, 0)$ .

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + c, b + d),$$

$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

This makes  $\mathbb{C}$  into a field (exercise).

$\mathbb{R} \subset \mathbb{C}$  by identifying  $x \in \mathbb{R}$  with  $(x, 0)$ .

*Imaginary unit:*

$$i \stackrel{\text{def}}{=} (0, 1)$$

Definition of the complex numbers (complex field, complex plane):

$$\mathbb{C} \stackrel{\text{def}}{=} \mathbb{R}^2$$

with multiplication:

$$(a, b) + (c, d) \stackrel{\text{def}}{=} (a + c, b + d),$$

$$(a, b)(c, d) \stackrel{\text{def}}{=} (ac - bd, bc + da).$$

This makes  $\mathbb{C}$  into a field (exercise).

$\mathbb{R} \subset \mathbb{C}$  by identifying  $x \in \mathbb{R}$  with  $(x, 0)$ .

*Imaginary unit:*

$$i \stackrel{\text{def}}{=} (0, 1)$$

$i^2 = -1$  so  $z^2 + 1 = 0$  has the two solutions  $i$  and  $-i$ .

$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

The evil twin of  $z = x + iy$  is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

The evil twin of  $z = x + iy$  is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the  $x$  and  $y$ :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

The evil twin of  $z = x + iy$  is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the  $x$  and  $y$ :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

The evil twin of  $z = x + iy$  is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the  $x$  and  $y$ :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

An expression in  $x, y$  can be written in terms of  $z, \bar{z}$  and vice versa:

$$x^3 + y^3 + 3ixy = \left(\frac{z + \bar{z}}{2}\right)^3 + \left(\frac{z - \bar{z}}{2i}\right)^3 + 3i\left(\frac{z + \bar{z}}{2}\right)\left(\frac{z - \bar{z}}{2i}\right),$$



$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

The evil twin of  $z = x + iy$  is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the  $x$  and  $y$ :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

An expression in  $x, y$  can be written in terms of  $z, \bar{z}$  and vice versa:

$$x^3 + y^3 + 3ixy = \left(\frac{z + \bar{z}}{2}\right)^3 + \left(\frac{z - \bar{z}}{2i}\right)^3 + 3i\left(\frac{z + \bar{z}}{2}\right)\left(\frac{z - \bar{z}}{2i}\right),$$

or

$$z^2 - i\bar{z}^2 + z\bar{z} = (x + iy)^2 - i(x - iy)^2 + (x + iy)(x - iy).$$

$(x, y) = x + iy$ , so use  $x + iy$  from now on (*cartesian form*).

The evil twin of  $z = x + iy$  is the *complex conjugate*

$$\bar{z} \stackrel{\text{def}}{=} x - iy.$$

Let's give names to the  $x$  and  $y$ :

$$\operatorname{Re} z = \operatorname{Re}(x + iy) \stackrel{\text{def}}{=} \frac{z + \bar{z}}{2} = x \quad \text{real part of } z.$$

$$\operatorname{Im} z = \operatorname{Im}(x + iy) \stackrel{\text{def}}{=} \frac{z - \bar{z}}{2i} = y \quad \text{imaginary part of } z.$$

An expression in  $x, y$  can be written in terms of  $z, \bar{z}$  and vice versa:

$$x^3 + y^3 + 3ixy = \left(\frac{z + \bar{z}}{2}\right)^3 + \left(\frac{z - \bar{z}}{2i}\right)^3 + 3i\left(\frac{z + \bar{z}}{2}\right)\left(\frac{z - \bar{z}}{2i}\right),$$

or

$$z^2 - i\bar{z}^2 + z\bar{z} = (x + iy)^2 - i(x - iy)^2 + (x + iy)(x - iy).$$

Almost looks as if  $z$  and  $\bar{z}$  were independent variables.