


22. Forced oscillations and resonance, part 2: Damped forced motion and practical resonance (Notes on Diffy Qs, 2.6)

Jiří Lebl


Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Recall our setup:



Recall our setup:



The equation is

$$mx'' + cx' + kx = F(t),$$


m is the mass,

c is friction,

k is the spring constant, and

$F(t)$ is an external force acting on the mass.

Recall our setup:



The equation is

$$mx'' + cx' + kx = F(t),$$

m is the mass,

c is friction,

k is the spring constant, and

$F(t)$ is an external force acting on the mass.

We are considering

$$F(t) = F_0 \cos(\omega t)$$

We consider damped ($c > 0$) motion:

$$mx'' + cx' + kx = F_0 \cos(\omega t).$$

We consider damped ($c > 0$) motion:

$$mx'' + cx' + kx = F_0 \cos(\omega t).$$

Write:

$$p = \frac{c}{2m}, \quad \omega_0 = \sqrt{\frac{k}{m}}.$$

Equation becomes:

$$x'' + 2px' + \omega_0^2 x = \frac{F_0}{m} \cos(\omega t).$$

We consider damped ($c > 0$) motion:

$$mx'' + cx' + kx = F_0 \cos(\omega t).$$

Write:

$$p = \frac{c}{2m}, \quad \omega_0 = \sqrt{\frac{k}{m}}.$$

Equation becomes:

$$x'' + 2px' + \omega_0^2 x = \frac{F_0}{m} \cos(\omega t).$$

Roots of the characteristic equation are $-p \pm \sqrt{p^2 - \omega_0^2}$, so the complementary solution is

$$x_c = \begin{cases} C_1 e^{r_1 t} + C_2 e^{r_2 t} & \text{if } c^2 > 4km, \\ C_1 e^{-pt} + C_2 t e^{-pt} & \text{if } c^2 = 4km, \\ e^{-pt} (C_1 \cos(\omega_1 t) + C_2 \sin(\omega_1 t)) & \text{if } c^2 < 4km, \end{cases}$$

where $\omega_1 = \sqrt{\omega_0^2 - p^2}$.

We consider damped ($c > 0$) motion:

$$mx'' + cx' + kx = F_0 \cos(\omega t).$$

Write:

$$p = \frac{c}{2m}, \quad \omega_0 = \sqrt{\frac{k}{m}}.$$

Equation becomes:

$$x'' + 2px' + \omega_0^2 x = \frac{F_0}{m} \cos(\omega t).$$

Roots of the characteristic equation are $-p \pm \sqrt{p^2 - \omega_0^2}$, so the complementary solution is

$$x_c = \begin{cases} C_1 e^{r_1 t} + C_2 e^{r_2 t} & \text{if } c^2 > 4km, \\ C_1 e^{-pt} + C_2 t e^{-pt} & \text{if } c^2 = 4km, \\ e^{-pt} (C_1 \cos(\omega_1 t) + C_2 \sin(\omega_1 t)) & \text{if } c^2 < 4km, \end{cases}$$

where $\omega_1 = \sqrt{\omega_0^2 - p^2}$.

Note that $x_c(t) \rightarrow 0$ as $t \rightarrow \infty$.

We need a particular solution. Try

$$x_p = A \cos(\omega t) + B \sin(\omega t).$$

We need a particular solution. Try

$$x_p = A \cos(\omega t) + B \sin(\omega t).$$

We plug into the equation $x'' + 2px' + \omega_0^2 x = \frac{F_0}{m} \cos(\omega t)$:

$$((\omega_0^2 - \omega^2)B - 2\omega pA) \sin(\omega t) + ((\omega_0^2 - \omega^2)A + 2\omega pB) \cos(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

We need a particular solution. Try

$$x_p = A \cos(\omega t) + B \sin(\omega t).$$

We plug into the equation $x'' + 2px' + \omega_0^2 x = \frac{F_0}{m} \cos(\omega t)$:

$$((\omega_0^2 - \omega^2)B - 2\omega pA) \sin(\omega t) + ((\omega_0^2 - \omega^2)A + 2\omega pB) \cos(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

Solve for A and B :

$$A = \frac{(\omega_0^2 - \omega^2)F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2}, \quad B = \frac{2\omega pF_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2}.$$

Compute the amplitude of x_p :

$$C = \sqrt{A^2 + B^2} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}}.$$

So our particular solution is

$$\begin{aligned}x_p &= A \cos(\omega t) + B \sin(\omega t) \\&= \frac{(\omega_0^2 - \omega^2)F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2} \cos(\omega t) + \frac{2\omega p F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2} \sin(\omega t).\end{aligned}$$

So our particular solution is

$$\begin{aligned}x_p &= A \cos(\omega t) + B \sin(\omega t) \\&= \frac{(\omega_0^2 - \omega^2)F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2} \cos(\omega t) + \frac{2\omega p F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2} \sin(\omega t).\end{aligned}$$

Or

$$x_p = C \cos(\omega t - \gamma) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$

So our particular solution is

$$\begin{aligned}x_p &= A \cos(\omega t) + B \sin(\omega t) \\&= \frac{(\omega_0^2 - \omega^2)F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2} \cos(\omega t) + \frac{2\omega p F_0}{m(2\omega p)^2 + m(\omega_0^2 - \omega^2)^2} \sin(\omega t).\end{aligned}$$

Or

$$x_p = C \cos(\omega t - \gamma) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$

Remark: γ can be computed as before, e.g. if $A \neq 0$ ($\omega \neq \omega_0$), then $\tan \gamma = \frac{B}{A} = \frac{2\omega p}{\omega_0^2 - \omega^2}$. If $A = 0$, then $B = C = \frac{F_0}{2m\omega p}$, and $\gamma = \pi/2$.

Call x_c the *transient solution* and denote it by x_{tr} .

Call x_c the *transient solution* and denote it by x_{tr} .

Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .

Call x_c the *transient solution* and denote it by x_{tr} .

Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .

So $x = x_c + x_p = x_{tr} + x_{sp}$.


Call x_c the *transient solution* and denote it by x_{tr} .

Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .

So $x = x_c + x_p = x_{tr} + x_{sp}$.

$x_{tr} \rightarrow 0$ as $t \rightarrow \infty$, so for large t , x_{tr} is negligible.

Let's ignore x_{tr} .



$k = 1, m = 1, F_0 = 1, c = 0.7$, and $\omega = 1.1$.

Call x_c the *transient solution* and denote it by x_{tr} .


Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .

So $x = x_c + x_p = x_{tr} + x_{sp}$.

$x_{tr} \rightarrow 0$ as $t \rightarrow \infty$, so for large t , x_{tr} is negligible.

Let's ignore x_{tr} .

The larger the p (that is, c), the faster $x_{tr} \rightarrow 0$,
 \Rightarrow smaller "transient region."



$k = 1, m = 1, F_0 = 1, c = 0.7$, and $\omega = 1.1$.

Call x_c the *transient solution* and denote it by x_{tr} .

Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .


So $x = x_c + x_p = x_{tr} + x_{sp}$.

$x_{tr} \rightarrow 0$ as $t \rightarrow \infty$, so for large t , x_{tr} is negligible.

Let's ignore x_{tr} .

The larger the p (that is, c), the faster $x_{tr} \rightarrow 0$,
 \Rightarrow smaller "transient region."

x_{sp} is not affected by initial conditions.



$k = 1, m = 1, F_0 = 1, c = 0.7$, and $\omega = 1.1$.

Call x_c the *transient solution* and denote it by x_{tr} .

Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .

So $x = x_c + x_p = x_{tr} + x_{sp}$.


$x_{tr} \rightarrow 0$ as $t \rightarrow \infty$, so for large t , x_{tr} is negligible.

Let's ignore x_{tr} .

The larger the p (that is, c), the faster $x_{tr} \rightarrow 0$,
 \Rightarrow smaller "transient region."

x_{sp} is not affected by initial conditions.

Remark: Undamped motion, $c = 0$, means
"infinite transient region" (i.e., not "transient")



$k = 1, m = 1, F_0 = 1, c = 0.7$, and $\omega = 1.1$.

Call x_c the *transient solution* and denote it by x_{tr} .

Call x_p (the one above) the *steady periodic solution* and denote it by x_{sp} .

So $x = x_c + x_p = x_{tr} + x_{sp}$.

$x_{tr} \rightarrow 0$ as $t \rightarrow \infty$, so for large t , x_{tr} is negligible.


Let's ignore x_{tr} .

The larger the p (that is, c), the faster $x_{tr} \rightarrow 0$,
 \Rightarrow smaller "transient region."

x_{sp} is not affected by initial conditions.

Remark: Undamped motion, $c = 0$, means
"infinite transient region" (i.e., not "transient")

We focus on x_{sp} and its amplitude C .



There is no (pure) resonance as there is no factor of t in

$$x_{sp} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$

There is no (pure) resonance as there is no factor of t in

$$x_{sp} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$

We focus on the amplitude as we vary the forcing frequency ω :

$$C = C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}}$$

There is no (pure) resonance as there is no factor of t in

$$x_{sp} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$


We focus on the amplitude as we vary the forcing frequency ω :

$$C = C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}}$$

Graph of $C(\omega)$ with $k = 1$, $m = 1$, $F_0 = 1$:

Top line: $c = 0.4$, middle line: $c = 0.8$,

bottom line: $c = 1.6$.



There is no (pure) resonance as there is no factor of t in

$$x_{sp} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$


We focus on the amplitude as we vary the forcing frequency ω :

$$C = C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}}$$

Graph of $C(\omega)$ with $k = 1$, $m = 1$, $F_0 = 1$:

Top line: $c = 0.4$, middle line: $c = 0.8$,
bottom line: $c = 1.6$.

Call the $\omega > 0$ that maximizes C the
practical resonance frequency.



There is no (pure) resonance as there is no factor of t in

$$x_{sp} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$

We focus on the amplitude as we vary the forcing frequency ω :


$$C = C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}}$$

Graph of $C(\omega)$ with $k = 1$, $m = 1$, $F_0 = 1$:

Top line: $c = 0.4$, middle line: $c = 0.8$,
bottom line: $c = 1.6$.

Call the $\omega > 0$ that maximizes C the
practical resonance frequency.

$C(\omega)$ is then the *practical resonance amplitude*.



There is no (pure) resonance as there is no factor of t in

$$x_{sp} = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}} \cos(\omega t - \gamma).$$

We focus on the amplitude as we vary the forcing frequency ω :

$$C = C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}}$$

Graph of $C(\omega)$ with $k = 1$, $m = 1$, $F_0 = 1$:


Top line: $c = 0.4$, middle line: $c = 0.8$,
bottom line: $c = 1.6$.

Call the $\omega > 0$ that maximizes C the
practical resonance frequency.

$C(\omega)$ is then the *practical resonance amplitude*.

We call this *practical resonance*.


Practical resonance gets bigger if c gets smaller, and disappears when damping is too large.



To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$

find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$

$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.




To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$

find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$

$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.

At most one critical point for $\omega > 0$.



To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$


find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$

$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.

At most one critical point for $\omega > 0$.

So there is practical resonance if $2p^2 + \omega^2 - \omega_0^2 = 0$ has a solution (if $\omega_0^2 - 2p^2 > 0$):

$$\text{practical resonance } \omega = \sqrt{\omega_0^2 - 2p^2}.$$



To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$

find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$


$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.

At most one critical point for $\omega > 0$.

So there is practical resonance if $2p^2 + \omega^2 - \omega_0^2 = 0$ has a solution (if $\omega_0^2 - 2p^2 > 0$):

$$\text{practical resonance } \omega = \sqrt{\omega_0^2 - 2p^2}.$$

If practical resonance occurs, it is smaller than ω_0 .



To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$

find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$

$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.


At most one critical point for $\omega > 0$.

So there is practical resonance if $2p^2 + \omega^2 - \omega_0^2 = 0$ has a solution (if $\omega_0^2 - 2p^2 > 0$):

$$\text{practical resonance } \omega = \sqrt{\omega_0^2 - 2p^2}.$$

If practical resonance occurs, it is smaller than ω_0 .

For very small damping c , practical resonance frequency $\omega \approx \omega_0$.
(Indeed, if $c = 0$, ω_0 is the (pure) resonance frequency.)



To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$

find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$

$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.

At most one critical point for $\omega > 0$.


So there is practical resonance if $2p^2 + \omega^2 - \omega_0^2 = 0$ has a solution (if $\omega_0^2 - 2p^2 > 0$):

$$\text{practical resonance } \omega = \sqrt{\omega_0^2 - 2p^2}.$$

If practical resonance occurs, it is smaller than ω_0 .

For very small damping c , practical resonance frequency $\omega \approx \omega_0$.
(Indeed, if $c = 0$, ω_0 is the (pure) resonance frequency.)

As c grows, practical resonance frequency ω goes to 0.



To find the max of $C(\omega) = \frac{F_0}{m\sqrt{(2\omega p)^2 + (\omega_0^2 - \omega^2)^2}},$

find where $0 = C'(\omega) = \frac{-2\omega(2p^2 + \omega^2 - \omega_0^2)F_0}{m((2\omega p)^2 + (\omega_0^2 - \omega^2)^2)^{3/2}}.$

$C'(\omega) = 0$ when $\omega = 0$ or $2p^2 + \omega^2 - \omega_0^2 = 0$.

At most one critical point for $\omega > 0$.

So there is practical resonance if $2p^2 + \omega^2 - \omega_0^2 = 0$ has a solution (if $\omega_0^2 - 2p^2 > 0$):

$$\text{practical resonance } \omega = \sqrt{\omega_0^2 - 2p^2}.$$

If practical resonance occurs, it is smaller than ω_0 .

For very small damping c , practical resonance frequency $\omega \approx \omega_0$.

(Indeed, if $c = 0$, ω_0 is the (pure) resonance frequency.)

As c grows, practical resonance frequency ω goes to 0.

Remark: $C(\omega) \rightarrow 0$ as $\omega \rightarrow \infty$. \Rightarrow Large forcing frequency means small amplitude.

