

## 26. Transforms of derivatives and ODEs, part 2 (Notes on Diffy Qs, 6.2)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider the initial conditions are zero.

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider the initial conditions are zero.

Taking the Laplace transform the equation becomes

$$A(s)X(s) = F(s).$$

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider the initial conditions are zero.

Taking the Laplace transform the equation becomes

$$A(s)X(s) = F(s).$$

The ratio  $\frac{X(s)}{F(s)} = \frac{1}{A(s)}$  is called the *transfer function* and we denote it by  $H(s)$ .

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider the initial conditions are zero.

Taking the Laplace transform the equation becomes

$$A(s)X(s) = F(s).$$

The ratio  $\frac{X(s)}{F(s)} = \frac{1}{A(s)}$  is called the *transfer function* and we denote it by  $H(s)$ . So

$$X(s) = H(s)F(s).$$

The output  $X(s)$  is just a multiplication away from the input  $F(s)$ .

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider the initial conditions are zero.

Taking the Laplace transform the equation becomes

$$A(s)X(s) = F(s).$$

The ratio  $\frac{X(s)}{F(s)} = \frac{1}{A(s)}$  is called the *transfer function* and we denote it by  $H(s)$ . So

$$X(s) = H(s)F(s).$$

The output  $X(s)$  is just a multiplication away from the input  $F(s)$ .

Moreover, as  $H(s) = \frac{X(s)}{F(s)}$ , we need to simply know the output  $X(s)$  for one input  $F(s)$  to find the output for all inputs.

Consider a constant coefficient linear differential equation  $Lx = f(t)$ .

Consider  $f(t)$  the input and  $x(t)$  the output.

We want a way to analyze outputs for all sorts of inputs (but same  $L$ ). E.g., we want to understand how a specific circuit responds to different inputs.

Consider the initial conditions are zero.

Taking the Laplace transform the equation becomes

$$A(s)X(s) = F(s).$$

The ratio  $\frac{X(s)}{F(s)} = \frac{1}{A(s)}$  is called the *transfer function* and we denote it by  $H(s)$ . So

$$X(s) = H(s)F(s).$$

The output  $X(s)$  is just a multiplication away from the input  $F(s)$ .

Moreover, as  $H(s) = \frac{X(s)}{F(s)}$ , we need to simply know the output  $X(s)$  for one input  $F(s)$  to find the output for all inputs. We don't even need to know  $L$  itself, just its output.

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

Laplace transform the equation:

$$s^2 X(s) + \omega_0^2 X(s) = F(s).$$

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

Laplace transform the equation:

$$s^2 X(s) + \omega_0^2 X(s) = F(s).$$

The transfer function is

$$H(s) = \frac{X(s)}{F(s)} = \frac{1}{s^2 + \omega_0^2}.$$

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

Laplace transform the equation:

$$s^2 X(s) + \omega_0^2 X(s) = F(s).$$

The transfer function is

$$H(s) = \frac{X(s)}{F(s)} = \frac{1}{s^2 + \omega_0^2}.$$

So if the input is the constant  $f(t) = 1$ , then  $F(s) = 1/s$ .

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

Laplace transform the equation:

$$s^2 X(s) + \omega_0^2 X(s) = F(s).$$

The transfer function is

$$H(s) = \frac{X(s)}{F(s)} = \frac{1}{s^2 + \omega_0^2}.$$

So if the input is the constant  $f(t) = 1$ , then  $F(s) = 1/s$ .

Then the output is

$$X(s) = H(s)F(s) = \frac{1}{s^2 + \omega_0^2} \frac{1}{s}.$$

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

Laplace transform the equation:

$$s^2 X(s) + \omega_0^2 X(s) = F(s).$$

The transfer function is

$$H(s) = \frac{X(s)}{F(s)} = \frac{1}{s^2 + \omega_0^2}.$$

So if the input is the constant  $f(t) = 1$ , then  $F(s) = 1/s$ .

Then the output is

$$X(s) = H(s)F(s) = \frac{1}{s^2 + \omega_0^2} \frac{1}{s}.$$

The inverse Laplace transform gives

$$x(t) = \frac{1 - \cos(\omega_0 t)}{\omega_0^2}.$$

**Example:** Consider  $x'' + \omega_0^2 x = f(t)$  (zero initial conditions).

Laplace transform the equation:

$$s^2 X(s) + \omega_0^2 X(s) = F(s).$$

The transfer function is

$$H(s) = \frac{X(s)}{F(s)} = \frac{1}{s^2 + \omega_0^2}.$$

So if the input is the constant  $f(t) = 1$ , then  $F(s) = 1/s$ .

Then the output is

$$X(s) = H(s)F(s) = \frac{1}{s^2 + \omega_0^2} \frac{1}{s}.$$

The inverse Laplace transform gives

$$x(t) = \frac{1 - \cos(\omega_0 t)}{\omega_0^2}.$$

For any other input  $f(t)$ , the output (in  $s$ -space) is again simply

$$X(s) = H(s)F(s) = \frac{1}{s^2 + \omega_0^2} F(s).$$

Laplace also deals with integrals easily.

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s)$$

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s) \quad \text{or perhaps} \quad \int_0^t f(\tau) d\tau = \mathcal{L}^{-1} \left\{ \frac{1}{s} F(s) \right\}.$$

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s) \quad \text{or perhaps} \quad \int_0^t f(\tau) d\tau = \mathcal{L}^{-1} \left\{ \frac{1}{s} F(s) \right\}.$$

**Example:**

$$\mathcal{L}^{-1} \left\{ \frac{1}{s(s^2 + 1)} \right\}$$

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s) \quad \text{or perhaps} \quad \int_0^t f(\tau) d\tau = \mathcal{L}^{-1} \left\{ \frac{1}{s} F(s) \right\}.$$

**Example:**

$$\mathcal{L}^{-1} \left\{ \frac{1}{s(s^2 + 1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{s} \frac{1}{s^2 + 1} \right\}$$

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s) \quad \text{or perhaps} \quad \int_0^t f(\tau) d\tau = \mathcal{L}^{-1} \left\{ \frac{1}{s} F(s) \right\}.$$

**Example:**

$$\mathcal{L}^{-1} \left\{ \frac{1}{s(s^2 + 1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{s} \frac{1}{s^2 + 1} \right\} = \int_0^t \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + 1} \right\} d\tau$$

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s) \quad \text{or perhaps} \quad \int_0^t f(\tau) d\tau = \mathcal{L}^{-1} \left\{ \frac{1}{s} F(s) \right\}.$$

**Example:**

$$\mathcal{L}^{-1} \left\{ \frac{1}{s(s^2 + 1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{s} \frac{1}{s^2 + 1} \right\} = \int_0^t \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + 1} \right\} d\tau = \int_0^t \sin \tau d\tau$$

Laplace also deals with integrals easily.

Applying the definition and a little bit of calculus gives the property

$$\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s) \quad \text{or perhaps} \quad \int_0^t f(\tau) d\tau = \mathcal{L}^{-1} \left\{ \frac{1}{s} F(s) \right\}.$$

**Example:**

$$\mathcal{L}^{-1} \left\{ \frac{1}{s(s^2 + 1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{s} \frac{1}{s^2 + 1} \right\} = \int_0^t \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + 1} \right\} d\tau = \int_0^t \sin \tau d\tau = 1 - \cos t.$$

An equation containing integrals is called an *integral equation*.

An equation containing integrals is called an *integral equation*.

**Example:** Solve for  $x(t)$  in

$$x(t) - t = \int_0^t x(\tau) d\tau.$$

An equation containing integrals is called an *integral equation*.

**Example:** Solve for  $x(t)$  in

$$x(t) - t = \int_0^t x(\tau) d\tau.$$

Apply the Laplace transform. Remember  $\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s)$ .

$$X(s) - \frac{1}{s^2} = \frac{1}{s} X(s).$$

An equation containing integrals is called an *integral equation*.

**Example:** Solve for  $x(t)$  in

$$x(t) - t = \int_0^t x(\tau) d\tau.$$

Apply the Laplace transform. Remember  $\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s)$ .

$$X(s) - \frac{1}{s^2} = \frac{1}{s} X(s).$$

So

$$X(s) = \frac{1}{s(s-1)}$$

An equation containing integrals is called an *integral equation*.

**Example:** Solve for  $x(t)$  in

$$x(t) - t = \int_0^t x(\tau) d\tau.$$

Apply the Laplace transform. Remember  $\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s)$ .

$$X(s) - \frac{1}{s^2} = \frac{1}{s} X(s).$$

So

$$X(s) = \frac{1}{s(s-1)} = \frac{1}{s-1} - \frac{1}{s}.$$

An equation containing integrals is called an *integral equation*.

**Example:** Solve for  $x(t)$  in

$$x(t) - t = \int_0^t x(\tau) d\tau.$$

Apply the Laplace transform. Remember  $\mathcal{L} \left\{ \int_0^t f(\tau) d\tau \right\} = \frac{1}{s} F(s)$ .

$$X(s) - \frac{1}{s^2} = \frac{1}{s} X(s).$$

So

$$X(s) = \frac{1}{s(s-1)} = \frac{1}{s-1} - \frac{1}{s}.$$

The inverse Laplace transform gives

$$x(t) = e^t - 1.$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt \end{aligned}$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \end{aligned}$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

Solve for  $F(s)$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

$$\text{Solve for } F(s) = \frac{1}{1 - e^{-Ps}} \int_0^P e^{-st} f(t) dt.$$

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

$$\text{Solve for } F(s) = \frac{1}{1 - e^{-Ps}} \int_0^P e^{-st} f(t) dt.$$

**Example:** Suppose  $f(t)$  is a *sawtooth*, that is,  $f(t) = t$  for  $0 \leq t < 1$  and  $f(t) = f(t + 1)$ .

How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).


$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

$$\text{Solve for } F(s) = \frac{1}{1 - e^{-Ps}} \int_0^P e^{-st} f(t) dt.$$

**Example:** Suppose  $f(t)$  is a *sawtooth*, that is,  $f(t) = t$  for  $0 \leq t < 1$  and  $f(t) = f(t + 1)$ .

So  $f(t) = t - 1$  for  $1 \leq t < 2$ ,

$f(t) = t - 2$  for  $2 \leq t < 3$ , etc.



How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

$$\text{Solve for } F(s) = \frac{1}{1 - e^{-Ps}} \int_0^P e^{-st} f(t) dt.$$


**Example:** Suppose  $f(t)$  is a *sawtooth*, that is,  $f(t) = t$  for  $0 \leq t < 1$  and  $f(t) = f(t + 1)$ .

So  $f(t) = t - 1$  for  $1 \leq t < 2$ ,

$f(t) = t - 2$  for  $2 \leq t < 3$ , etc.

Then  $P = 1$  and

$$F(s) = \frac{1}{1 - e^{-s}} \int_0^1 e^{-st} t dt$$



How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

$$\text{Solve for } F(s) = \frac{1}{1 - e^{-Ps}} \int_0^P e^{-st} f(t) dt.$$


**Example:** Suppose  $f(t)$  is a *sawtooth*, that is,  $f(t) = t$  for  $0 \leq t < 1$  and  $f(t) = f(t + 1)$ .

So  $f(t) = t - 1$  for  $1 \leq t < 2$ ,

$f(t) = t - 2$  for  $2 \leq t < 3$ , etc.

Then  $P = 1$  and

$$F(s) = \frac{1}{1 - e^{-s}} \int_0^1 e^{-st} t dt = \frac{1}{1 - e^{-s}} \left( \frac{-e^{-s}}{s} - \frac{e^{-s}}{s^2} + \frac{1}{s^2} \right)$$



How about periodic functions? That is, an  $f(t)$  where  $f(t) = f(t + P)$  for some  $P$  (the period).

$$\begin{aligned} F(s) &= \int_0^{\infty} e^{-st} f(t) dt = \int_0^P e^{-st} f(t) dt + \int_P^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + \int_0^{\infty} e^{-s(t+P)} f(t+P) dt = \int_0^P e^{-st} f(t) dt + e^{-Ps} \int_0^{\infty} e^{-st} f(t) dt \\ &= \int_0^P e^{-st} f(t) dt + e^{-Ps} F(s). \end{aligned}$$

Solve for  $F(s) = \frac{1}{1 - e^{-Ps}} \int_0^P e^{-st} f(t) dt$ .

**Example:** Suppose  $f(t)$  is a *sawtooth*, that is,  $f(t) = t$  for  $0 \leq t < 1$  and  $f(t) = f(t + 1)$ .

So  $f(t) = t - 1$  for  $1 \leq t < 2$ ,

$f(t) = t - 2$  for  $2 \leq t < 3$ , etc.

Then  $P = 1$  and

$$F(s) = \frac{1}{1 - e^{-s}} \int_0^1 e^{-st} t dt = \frac{1}{1 - e^{-s}} \left( \frac{-e^{-s}}{s} - \frac{e^{-s}}{s^2} + \frac{1}{s^2} \right) = \frac{-e^{-s}}{(1 - e^{-s})s} + \frac{1}{s^2}.$$

