

20. Nonhomogeneous equations, part 2: variation of parameters (Notes on Diffy Qs, 2.5)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

We're trying to solve $Ly = f(x)$.

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

$$f'(x) = \sec^2 x,$$

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

$$f'(x) = \sec^2 x,$$

$$f''(x) = 2 \sec^2 x \tan x,$$

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

$$f'(x) = \sec^2 x,$$

$$f''(x) = 2 \sec^2 x \tan x,$$

$$f'''(x) = 4 \sec^2 x \tan^2 x + 2 \sec^4 x,$$

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

$$f'(x) = \sec^2 x,$$

$$f''(x) = 2 \sec^2 x \tan x,$$

$$f'''(x) = 4 \sec^2 x \tan^2 x + 2 \sec^4 x,$$

$$f^{(4)}(x) = 8 \sec^2 x \tan^3 x + 16 \sec^4 x \tan x,$$

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

$$f'(x) = \sec^2 x,$$

$$f''(x) = 2 \sec^2 x \tan x,$$

$$f'''(x) = 4 \sec^2 x \tan^2 x + 2 \sec^4 x,$$

$$f^{(4)}(x) = 8 \sec^2 x \tan^3 x + 16 \sec^4 x \tan x,$$

$$f^{(5)}(x) = 16 \sec^2 x \tan^4 x + 88 \sec^4 x \tan^2 x + 16 \sec^6 x,$$

...

We're trying to solve $Ly = f(x)$.

Undetermined coefficients needs $f(x)$ and its derivatives to be of the same general form.

What about $Ly = \tan x$, that is, $f(x) = \tan x$?

$$f'(x) = \sec^2 x,$$

$$f''(x) = 2 \sec^2 x \tan x,$$

$$f'''(x) = 4 \sec^2 x \tan^2 x + 2 \sec^4 x,$$

$$f^{(4)}(x) = 8 \sec^2 x \tan^3 x + 16 \sec^4 x \tan x,$$

$$f^{(5)}(x) = 16 \sec^2 x \tan^4 x + 88 \sec^4 x \tan^2 x + 16 \sec^6 x,$$

...

No luck ...

Variation of parameters can handle $\tan x$ or any other $f(x)$ (but computations can get tedious).

Variation of parameters can handle $\tan x$ or any other $f(x)$ (but computations can get tedious).

The technique works for the general linear equations of any order, but we stick to constant coefficient second order equations.

Variation of parameters can handle $\tan x$ or any other $f(x)$ (but computations can get tedious).

The technique works for the general linear equations of any order, but we stick to constant coefficient second order equations.

Suppose $Ly = 0$ (associated homogeneous equation) has the general solution

$$C_1y_1 + C_2y_2,$$

where C_1, C_2 are constants.

Variation of parameters can handle $\tan x$ or any other $f(x)$ (but computations can get tedious).

The technique works for the general linear equations of any order, but we stick to constant coefficient second order equations.

Suppose $Ly = 0$ (associated homogeneous equation) has the general solution

$$C_1y_1 + C_2y_2,$$

where C_1, C_2 are constants.

We look for a solution of $Ly = f(x)$ of the form

$$y = u_1y_1 + u_2y_2,$$

where u_1, u_2 are functions.

Example: Solve $y'' + y = \tan x$.

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

So let's look for a solution of the form

$$y = u_1 \cos x + u_2 \sin x = u_1 y_1 + u_2 y_2.$$

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

So let's look for a solution of the form

$$y = u_1 \cos x + u_2 \sin x = u_1 y_1 + u_2 y_2.$$

Compute

$$y' = (u_1' y_1 + u_2' y_2) + (u_1 y_1' + u_2 y_2').$$

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

So let's look for a solution of the form

$$y = u_1 \cos x + u_2 \sin x = u_1 y_1 + u_2 y_2.$$

Compute

$$y' = (u_1' y_1 + u_2' y_2) + (u_1 y_1' + u_2 y_2').$$

We have two unknown functions (u_1, u_2) and one condition $(Ly = f(x))$, so we can still impose another condition.

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

So let's look for a solution of the form

$$y = u_1 \cos x + u_2 \sin x = u_1 y_1 + u_2 y_2.$$

Compute

$$y' = (u_1' y_1 + u_2' y_2) + (u_1 y_1' + u_2 y_2').$$

We have two unknown functions (u_1, u_2) and one condition $(Ly = f(x))$, so we can still impose another condition. Wouldn't it be convenient if

$$u_1' y_1 + u_2' y_2 = 0?$$

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

So let's look for a solution of the form

$$y = u_1 \cos x + u_2 \sin x = u_1 y_1 + u_2 y_2.$$

Compute

$$y' = (u_1' y_1 + u_2' y_2) + (u_1 y_1' + u_2 y_2').$$

We have two unknown functions (u_1, u_2) and one condition $(Ly = f(x))$, so we can still impose another condition. Wouldn't it be convenient if

$$u_1' y_1 + u_2' y_2 = 0?$$

Yes? So let's assume that!

Example: Solve $y'' + y = \tan x$.

Complementary solution:

$$y_c = C_1 \cos x + C_2 \sin x.$$

So let's look for a solution of the form

$$y = u_1 \cos x + u_2 \sin x = u_1 y_1 + u_2 y_2.$$

Compute

$$y' = (u'_1 y_1 + u'_2 y_2) + (u_1 y'_1 + u_2 y'_2).$$

We have two unknown functions (u_1, u_2) and one condition $(Ly = f(x))$, so we can still impose another condition. Wouldn't it be convenient if

$$u'_1 y_1 + u'_2 y_2 = 0?$$

Yes? So let's assume that! Then

$$y' = u_1 y'_1 + u_2 y'_2.$$

Start with

$$y' = u_1 y'_1 + u_2 y'_2.$$

Start with

$$y' = u_1 y'_1 + u_2 y'_2.$$

Differentiate

$$y'' = (u'_1 y'_1 + u'_2 y'_2) + (u_1 y''_1 + u_2 y''_2).$$

Start with

$$y' = u_1 y_1' + u_2 y_2'.$$

Differentiate

$$y'' = (u_1' y_1' + u_2' y_2') + (u_1 y_1'' + u_2 y_2'').$$

y_1 and y_2 solve $y'' + y = 0$, so $y_1'' = -y_1$ and $y_2'' = -y_2$.

Start with

$$y' = u_1 y_1' + u_2 y_2'.$$

Differentiate

$$y'' = (u_1' y_1' + u_2' y_2') + (u_1 y_1'' + u_2 y_2'').$$

y_1 and y_2 solve $y'' + y = 0$, so $y_1'' = -y_1$ and $y_2'' = -y_2$.

So

$$y'' = (u_1' y_1' + u_2' y_2') - (u_1 y_1 + u_2 y_2)$$

Start with

$$y' = u_1 y'_1 + u_2 y'_2.$$

Differentiate

$$y'' = (u'_1 y'_1 + u'_2 y'_2) + (u_1 y''_1 + u_2 y''_2).$$

y_1 and y_2 solve $y'' + y = 0$, so $y''_1 = -y_1$ and $y''_2 = -y_2$.

So

$$y'' = (u'_1 y'_1 + u'_2 y'_2) - (u_1 y_1 + u_2 y_2) = (u'_1 y'_1 + u'_2 y'_2) - y$$

Start with

$$y' = u_1 y'_1 + u_2 y'_2.$$

Differentiate

$$y'' = (u'_1 y'_1 + u'_2 y'_2) + (u_1 y''_1 + u_2 y''_2).$$

y_1 and y_2 solve $y'' + y = 0$, so $y''_1 = -y_1$ and $y''_2 = -y_2$.

So

$$y'' = (u'_1 y'_1 + u'_2 y'_2) - (u_1 y_1 + u_2 y_2) = (u'_1 y'_1 + u'_2 y'_2) - y$$

In other words

$$y'' + y = Ly = u'_1 y'_1 + u'_2 y'_2.$$

Start with

$$y' = u_1 y_1' + u_2 y_2'.$$

Differentiate

$$y'' = (u_1' y_1' + u_2' y_2') + (u_1 y_1'' + u_2 y_2'').$$

y_1 and y_2 solve $y'' + y = 0$, so $y_1'' = -y_1$ and $y_2'' = -y_2$.

So

$$y'' = (u_1' y_1' + u_2' y_2') - (u_1 y_1 + u_2 y_2) = (u_1' y_1' + u_2' y_2') - y$$

In other words

$$y'' + y = Ly = u_1' y_1' + u_2' y_2'.$$

Soooo ... for y to satisfy $Ly = f(x)$, we must have $f(x) = u_1' y_1' + u_2' y_2'$.

Start with

$$y' = u_1 y_1' + u_2 y_2'.$$

Differentiate

$$y'' = (u_1' y_1' + u_2' y_2') + (u_1 y_1'' + u_2 y_2'').$$

y_1 and y_2 solve $y'' + y = 0$, so $y_1'' = -y_1$ and $y_2'' = -y_2$.

So

$$y'' = (u_1' y_1' + u_2' y_2') - (u_1 y_1 + u_2 y_2) = (u_1' y_1' + u_2' y_2') - y$$

In other words

$$y'' + y = Ly = u_1' y_1' + u_2' y_2'.$$

Soooo ... for y to satisfy $Ly = f(x)$, we must have $f(x) = u_1' y_1' + u_2' y_2'$.

To summarize we need to solve the two imposed conditions:

$$\begin{aligned} u_1' y_1 + u_2' y_2 &= 0, \\ u_1' y_1' + u_2' y_2' &= f(x). \end{aligned}$$

In our case

$$\begin{aligned}u_1' \cos x + u_2' \sin x &= 0, \\ -u_1' \sin x + u_2' \cos x &= \tan x.\end{aligned}$$

In our case

$$\begin{aligned}u_1' \cos x + u_2' \sin x &= 0, \\ -u_1' \sin x + u_2' \cos x &= \tan x.\end{aligned}$$

Hence

$$\begin{aligned}u_1' \cos x \sin x + u_2' \sin^2 x &= 0, \\ -u_1' \sin x \cos x + u_2' \cos^2 x &= \tan x \cos x = \sin x.\end{aligned}$$

In our case

$$\begin{aligned}u_1' \cos x + u_2' \sin x &= 0, \\ -u_1' \sin x + u_2' \cos x &= \tan x.\end{aligned}$$

Hence

$$\begin{aligned}u_1' \cos x \sin x + u_2' \sin^2 x &= 0, \\ -u_1' \sin x \cos x + u_2' \cos^2 x &= \tan x \cos x = \sin x.\end{aligned}$$

And thus

$$\begin{aligned}u_2' (\sin^2 x + \cos^2 x) &= \sin x, \\ u_2' &= \sin x, \\ u_1' &= \frac{-\sin^2 x}{\cos x} = \cos x - \sec x.\end{aligned}$$

In our case

$$\begin{aligned}u_1' \cos x + u_2' \sin x &= 0, \\ -u_1' \sin x + u_2' \cos x &= \tan x.\end{aligned}$$

Hence

$$\begin{aligned}u_1' \cos x \sin x + u_2' \sin^2 x &= 0, \\ -u_1' \sin x \cos x + u_2' \cos^2 x &= \tan x \cos x = \sin x.\end{aligned}$$

And thus

$$\begin{aligned}u_2' (\sin^2 x + \cos^2 x) &= \sin x, \\ u_2' &= \sin x, \\ u_1' &= \frac{-\sin^2 x}{\cos x} = \cos x - \sec x.\end{aligned}$$

Integrate

$$\begin{aligned}u_1 &= \int u_1' dx = \int (\cos x - \sec x) dx = \sin x - \ln |\sec x + \tan x|, \\ u_2 &= \int u_2' dx = \int \sin x dx = -\cos x.\end{aligned}$$

In our case

$$\begin{aligned}u_1' \cos x + u_2' \sin x &= 0, \\ -u_1' \sin x + u_2' \cos x &= \tan x.\end{aligned}$$

Hence

$$\begin{aligned}u_1' \cos x \sin x + u_2' \sin^2 x &= 0, \\ -u_1' \sin x \cos x + u_2' \cos^2 x &= \tan x \cos x = \sin x.\end{aligned}$$

And thus

$$\begin{aligned}u_2' (\sin^2 x + \cos^2 x) &= \sin x, \\ u_2' &= \sin x, \\ u_1' &= \frac{-\sin^2 x}{\cos x} = \cos x - \sec x.\end{aligned}$$

Integrate

$$\begin{aligned}u_1 &= \int u_1' dx = \int (\cos x - \sec x) dx = \sin x - \ln |\sec x + \tan x|, \\ u_2 &= \int u_2' dx = \int \sin x dx = -\cos x.\end{aligned}$$

(Forget about constants of integration, we want a particular solution)

So

$$y_p = u_1 y_1 + u_2 y_2$$

So

$$y_p = u_1 y_1 + u_2 y_2$$

$$= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x$$

So

$$\begin{aligned}y_p &= u_1 y_1 + u_2 y_2 \\&= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x \\&= -\cos x \ln|\sec x + \tan x|.\end{aligned}$$

So

$$\begin{aligned}y_p &= u_1 y_1 + u_2 y_2 \\&= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x \\&= -\cos x \ln|\sec x + \tan x|.\end{aligned}$$

The general solution to $y'' + y = \tan x$ is

$$y = C_1 \cos x + C_2 \sin x - \cos x \ln|\sec x + \tan x|.$$

So

$$\begin{aligned}y_p &= u_1 y_1 + u_2 y_2 \\&= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x \\&= -\cos x \ln|\sec x + \tan x|.\end{aligned}$$

The general solution to $y'' + y = \tan x$ is

$$y = C_1 \cos x + C_2 \sin x - \cos x \ln|\sec x + \tan x|.$$

In general, the procedure for finding y_p using variation parameters (2nd order) is

1) Solve the associated homogeneous equation.

So

$$\begin{aligned}y_p &= u_1 y_1 + u_2 y_2 \\&= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x \\&= -\cos x \ln|\sec x + \tan x|.\end{aligned}$$

The general solution to $y'' + y = \tan x$ is

$$y = C_1 \cos x + C_2 \sin x - \cos x \ln|\sec x + \tan x|.$$

In general, the procedure for finding y_p using variation parameters (2nd order) is

- 1) Solve the associated homogeneous equation.
- 2) Solve for u'_1 and u'_2 in

$$\begin{aligned}u'_1 y_1 + u'_2 y_2 &= 0, \\u'_1 y'_1 + u'_2 y'_2 &= f(x).\end{aligned}$$

So

$$\begin{aligned}y_p &= u_1 y_1 + u_2 y_2 \\&= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x \\&= -\cos x \ln|\sec x + \tan x|.\end{aligned}$$

The general solution to $y'' + y = \tan x$ is

$$y = C_1 \cos x + C_2 \sin x - \cos x \ln|\sec x + \tan x|.$$

In general, the procedure for finding y_p using variation parameters (2nd order) is

1) Solve the associated homogeneous equation.

2) Solve for u'_1 and u'_2 in

$$\begin{aligned}u'_1 y_1 + u'_2 y_2 &= 0, \\u'_1 y'_1 + u'_2 y'_2 &= f(x).\end{aligned}$$

3) Integrate to get u_1 and u_2 to find $y_p = u_1 y_1 + u_2 y_2$.

So

$$\begin{aligned}y_p &= u_1 y_1 + u_2 y_2 \\&= \cos x \sin x - \cos x \ln|\sec x + \tan x| - \cos x \sin x \\&= -\cos x \ln|\sec x + \tan x|.\end{aligned}$$

The general solution to $y'' + y = \tan x$ is

$$y = C_1 \cos x + C_2 \sin x - \cos x \ln|\sec x + \tan x|.$$

In general, the procedure for finding y_p using variation parameters (2nd order) is

1) Solve the associated homogeneous equation.

2) Solve for u'_1 and u'_2 in

$$u'_1 y_1 + u'_2 y_2 = 0,$$

$$u'_1 y'_1 + u'_2 y'_2 = f(x).$$

3) Integrate to get u_1 and u_2 to find $y_p = u_1 y_1 + u_2 y_2$.

Note: Undetermined coefficients are usually less tedious if applicable.