

16. Higher order linear ODEs (Notes on Diffy Qs, 2.3)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Let us consider linear equations of order higher than 2 — Mostly the same idea.

Let us consider linear equations of order higher than 2 — Mostly the same idea.

Theorem (Superposition)

If y_1, y_2, \dots, y_n are solutions of the homogeneous equation

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + \dots + p_1(x)y' + p_0(x)y = 0,$$

then the linear combination

$$y(x) = C_1y_1(x) + C_2y_2(x) + \dots + C_ny_n(x)$$

is also a solution for arbitrary constants C_1, C_2, \dots, C_n .

Theorem (Existence and uniqueness)

Suppose p_0 through p_{n-1} , and f are continuous functions on some interval I , a is a number in I , and b_0, b_1, \dots, b_{n-1} are constants. Then the equation

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + \dots + p_1(x)y' + p_0(x)y = f(x)$$

has exactly one solution $y(x)$ defined on the same interval I satisfying the initial conditions

$$y(a) = b_0, \quad y'(a) = b_1, \quad \dots, \quad y^{(n-1)}(a) = b_{n-1}.$$

To use superposition, we need enough *linearly independent* solutions.

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1y_1 + c_2y_2 + \cdots + c_ny_n = 0$$

has only the trivial solution $c_1 = c_2 = \cdots = c_n = 0$.

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others.

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x$

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x \Rightarrow z^2 = e^{2x}$ and $z^3 = e^{3x}$

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x \Rightarrow z^2 = e^{2x}$ and $z^3 = e^{3x} \Rightarrow$

$$c_1 z + c_2 z^2 + c_3 z^3 = 0.$$

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x \Rightarrow z^2 = e^{2x}$ and $z^3 = e^{3x} \Rightarrow$

$$c_1 z + c_2 z^2 + c_3 z^3 = 0.$$

This polynomial must be zero for any positive number $z = e^x$

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x \Rightarrow z^2 = e^{2x}$ and $z^3 = e^{3x} \Rightarrow$

$$c_1 z + c_2 z^2 + c_3 z^3 = 0.$$

This polynomial must be zero for any positive number $z = e^x \Rightarrow$ it is identically zero

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x \Rightarrow z^2 = e^{2x}$ and $z^3 = e^{3x} \Rightarrow$

$$c_1 z + c_2 z^2 + c_3 z^3 = 0.$$

This polynomial must be zero for any positive number $z = e^x \Rightarrow$ it is identically zero
 $\Rightarrow c_1 = c_2 = c_3 = 0$

To use superposition, we need enough *linearly independent* solutions.

The functions y_1, y_2, \dots, y_n are *linearly independent* if the equation

$$c_1 y_1 + c_2 y_2 + \dots + c_n y_n = 0$$

has only the trivial solution $c_1 = c_2 = \dots = c_n = 0$.

If there is a solution where, e.g., $c_1 \neq 0$, then we can solve for y_1 as a linear combination of the others. The functions are then *linearly dependent*.

Example: Show that e^x, e^{2x}, e^{3x} are linearly independent.

Consider

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Write $z = e^x \Rightarrow z^2 = e^{2x}$ and $z^3 = e^{3x} \Rightarrow$

$$c_1 z + c_2 z^2 + c_3 z^3 = 0.$$

This polynomial must be zero for any positive number $z = e^x \Rightarrow$ it is identically zero
 $\Rightarrow c_1 = c_2 = c_3 = 0 \Rightarrow$ the functions are linearly independent.

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty$

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$.

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$. Rinse, repeat!

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$. Rinse, repeat!

A third way: Suppose

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

This has to be true for all x , so plug in $x = 0$, $x = 1$, and $x = 2$.

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$. Rinse, repeat!

A third way: Suppose

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

This has to be true for all x , so plug in $x = 0$, $x = 1$, and $x = 2$.

We must have

$$c_1 + c_2 + c_3 = 0 \quad \text{and} \quad c_1 e + c_2 e^2 + c_3 e^3 = 0 \quad \text{and} \quad c_1 e^2 + c_2 e^4 + c_3 e^6 = 0.$$

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$. Rinse, repeat!

A third way: Suppose

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

This has to be true for all x , so plug in $x = 0$, $x = 1$, and $x = 2$.

We must have

$$c_1 + c_2 + c_3 = 0 \quad \text{and} \quad c_1 e + c_2 e^2 + c_3 e^3 = 0 \quad \text{and} \quad c_1 e^2 + c_2 e^4 + c_3 e^6 = 0.$$

Solving the three equations yields $c_1 = c_2 = c_3 = 0$.

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$. Rinse, repeat!

A third way: Suppose

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

This has to be true for all x , so plug in $x = 0$, $x = 1$, and $x = 2$.

We must have

$$c_1 + c_2 + c_3 = 0 \quad \text{and} \quad c_1 e + c_2 e^2 + c_3 e^3 = 0 \quad \text{and} \quad c_1 e^2 + c_2 e^4 + c_3 e^6 = 0.$$

Solving the three equations yields $c_1 = c_2 = c_3 = 0$.

A so-called *Wronskian* can also be used, but let's skip it.

Let's show that e^x, e^{2x}, e^{3x} are linearly independent in another way:

Start with

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

Divide by e^{3x} :

$$c_1 e^{-2x} + c_2 e^{-x} + c_3 = 0.$$

True for all x , so let $x \rightarrow \infty \Rightarrow c_3 = 0$.

So $c_1 e^x + c_2 e^{2x} = 0$. Rinse, repeat!

A third way: Suppose

$$c_1 e^x + c_2 e^{2x} + c_3 e^{3x} = 0.$$

This has to be true for all x , so plug in $x = 0$, $x = 1$, and $x = 2$.

We must have

$$c_1 + c_2 + c_3 = 0 \quad \text{and} \quad c_1 e + c_2 e^2 + c_3 e^3 = 0 \quad \text{and} \quad c_1 e^2 + c_2 e^4 + c_3 e^6 = 0.$$

Solving the three equations yields $c_1 = c_2 = c_3 = 0$.

A so-called *Wronskian* can also be used, but let's skip it.

The main thing to understand is the meaning of linear independence!

Example: The functions e^x , e^{-x} , and $\cosh x$ are linearly dependent:

Example: The functions e^x , e^{-x} , and $\cosh x$ are linearly dependent:

$$\cosh x = \frac{e^x + e^{-x}}{2} \quad \text{or} \quad 2 \cosh x - e^x - e^{-x} = 0.$$

Example: The functions e^x , e^{-x} , and $\cosh x$ are linearly dependent:

$$\cosh x = \frac{e^x + e^{-x}}{2} \quad \text{or} \quad 2 \cosh x - e^x - e^{-x} = 0.$$

Note: You must consider all the functions at once. The functions e^x and e^{-x} are linearly independent.

Example: The functions e^x , e^{-x} , and $\cosh x$ are linearly dependent:

$$\cosh x = \frac{e^x + e^{-x}}{2} \quad \text{or} \quad 2 \cosh x - e^x - e^{-x} = 0.$$

Note: You must consider all the functions at once. The functions e^x and e^{-x} are linearly independent.

Theorem

If y_1, y_2, \dots, y_n are linearly independent solutions of the homogeneous equation

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + \dots + p_1(x)y' + p_0(x)y = 0,$$

then the general solution can be written as

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + \dots + C_n y_n(x).$$

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots!

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

The general solution is

$$y = C_1 e^{-x} + C_2 e^x + C_3 e^{3x}.$$

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

The general solution is

$$y = C_1 e^{-x} + C_2 e^x + C_3 e^{3x}.$$

Initial conditions say

$$1 = y(0) = C_1 + C_2 + C_3,$$

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

The general solution is

$$y = C_1 e^{-x} + C_2 e^x + C_3 e^{3x}.$$

Initial conditions say

$$1 = y(0) = C_1 + C_2 + C_3, \quad 2 = y'(0) = -C_1 + C_2 + 3C_3,$$

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

The general solution is

$$y = C_1 e^{-x} + C_2 e^x + C_3 e^{3x}.$$

Initial conditions say

$$1 = y(0) = C_1 + C_2 + C_3, \quad 2 = y'(0) = -C_1 + C_2 + 3C_3, \quad 3 = y''(0) = C_1 + C_2 + 9C_3.$$

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

The general solution is

$$y = C_1 e^{-x} + C_2 e^x + C_3 e^{3x}.$$

Initial conditions say

$$1 = y(0) = C_1 + C_2 + C_3, \quad 2 = y'(0) = -C_1 + C_2 + 3C_3, \quad 3 = y''(0) = C_1 + C_2 + 9C_3.$$

We solve these to find $C_1 = -1/4$, $C_2 = 1$, and $C_3 = 1/4$.

Example: Solve $y''' - 3y'' - y' + 3y = 0$ subject to $y(0) = 1$, $y'(0) = 2$, and $y''(0) = 3$.

Try: $y = e^{rx}$:

$$\underbrace{r^3 e^{rx}}_{y'''} - 3 \underbrace{r^2 e^{rx}}_{y''} - \underbrace{r e^{rx}}_{y'} + 3 \underbrace{e^{rx}}_y = 0.$$

We divide by e^{rx} : $r^3 - 3r^2 - r + 3 = 0$.

Find the roots! By trial and error $r = -1, 1, 3$.

So solutions are $y_1 = e^{-x}$, $y_2 = e^x$, $y_3 = e^{3x}$.

The general solution is

$$y = C_1 e^{-x} + C_2 e^x + C_3 e^{3x}.$$

Initial conditions say

$$1 = y(0) = C_1 + C_2 + C_3, \quad 2 = y'(0) = -C_1 + C_2 + 3C_3, \quad 3 = y''(0) = C_1 + C_2 + 9C_3.$$

We solve these to find $C_1 = -1/4$, $C_2 = 1$, and $C_3 = 1/4$.

So the particular solution to our problem is:

$$y = \frac{-1}{4} e^{-x} + e^x + \frac{1}{4} e^{3x}.$$

The real trick is finding the roots.

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

One often uses a computer, but a good strategy to do it by hand is to plug in some easy numbers to start with: Start with 0, then try 1 and -1 , then try other integers.

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

One often uses a computer, but a good strategy to do it by hand is to plug in some easy numbers to start with: Start with 0, then try 1 and -1 , then try other integers.

E.g., for $r^3 - 3r^2 - r + 3 = 0$, we find $r_1 = 1$ and $r_2 = -1$ are roots by trying.

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

One often uses a computer, but a good strategy to do it by hand is to plug in some easy numbers to start with: Start with 0, then try 1 and -1 , then try other integers.

E.g., for $r^3 - 3r^2 - r + 3 = 0$, we find $r_1 = 1$ and $r_2 = -1$ are roots by trying.

The third root r_3 is easy to find:

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

One often uses a computer, but a good strategy to do it by hand is to plug in some easy numbers to start with: Start with 0, then try 1 and -1 , then try other integers.

E.g., for $r^3 - 3r^2 - r + 3 = 0$, we find $r_1 = 1$ and $r_2 = -1$ are roots by trying.

The third root r_3 is easy to find:

$$r^3 - 3r^2 - r + 3 = (r - r_1)(r - r_2)(r - r_3)$$

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

One often uses a computer, but a good strategy to do it by hand is to plug in some easy numbers to start with: Start with 0, then try 1 and -1 , then try other integers.

E.g., for $r^3 - 3r^2 - r + 3 = 0$, we find $r_1 = 1$ and $r_2 = -1$ are roots by trying.

The third root r_3 is easy to find:

$$r^3 - 3r^2 - r + 3 = (r - r_1)(r - r_2)(r - r_3)$$

So the constant term is

$$3 = (-r_1)(-r_2)(-r_3).$$

The real trick is finding the roots.

There are complicated formulas for degree 3 and 4 polynomials.

There is **no** formula for degree 5 or higher.

But there are always n roots for an n^{th} degree polynomial, though they can be repeated, and they may be complex.

One often uses a computer, but a good strategy to do it by hand is to plug in some easy numbers to start with: Start with 0, then try 1 and -1 , then try other integers.

E.g., for $r^3 - 3r^2 - r + 3 = 0$, we find $r_1 = 1$ and $r_2 = -1$ are roots by trying.

The third root r_3 is easy to find:

$$r^3 - 3r^2 - r + 3 = (r - r_1)(r - r_2)(r - r_3)$$

So the constant term is

$$3 = (-r_1)(-r_2)(-r_3).$$

So $3 = (-1)(1)(-r_3)$ or $r_3 = 3$.

If a real root r is repeated k times, then we have solutions

$$e^{rx}, \quad xe^{rx}, \quad x^2e^{rx}, \quad \dots, \quad x^{k-1}e^{rx}.$$

If a real root r is repeated k times, then we have solutions

$$e^{rx}, \quad xe^{rx}, \quad x^2e^{rx}, \quad \dots, \quad x^{k-1}e^{rx}.$$

Example: Solve $y^{(4)} - 3y''' + 3y'' - y' = 0$.

If a real root r is repeated k times, then we have solutions

$$e^{rx}, \quad xe^{rx}, \quad x^2e^{rx}, \quad \dots, \quad x^{k-1}e^{rx}.$$

Example: Solve $y^{(4)} - 3y''' + 3y'' - y' = 0$.

The characteristic equation is $r^4 - 3r^3 + 3r^2 - r = 0$.

If a real root r is repeated k times, then we have solutions

$$e^{rx}, \quad xe^{rx}, \quad x^2e^{rx}, \quad \dots, \quad x^{k-1}e^{rx}.$$

Example: Solve $y^{(4)} - 3y''' + 3y'' - y' = 0$.

The characteristic equation is $r^4 - 3r^3 + 3r^2 - r = 0$.

By inspection $r^4 - 3r^3 + 3r^2 - r = r(r - 1)^3$.

If a real root r is repeated k times, then we have solutions

$$e^{rx}, \quad xe^{rx}, \quad x^2e^{rx}, \quad \dots, \quad x^{k-1}e^{rx}.$$

Example: Solve $y^{(4)} - 3y''' + 3y'' - y' = 0$.

The characteristic equation is $r^4 - 3r^3 + 3r^2 - r = 0$.

By inspection $r^4 - 3r^3 + 3r^2 - r = r(r-1)^3$.

The roots are $r = 0, 1, 1, 1$. ($r = 1$ has multiplicity 3)

If a real root r is repeated k times, then we have solutions

$$e^{rx}, \quad xe^{rx}, \quad x^2e^{rx}, \quad \dots, \quad x^{k-1}e^{rx}.$$

Example: Solve $y^{(4)} - 3y''' + 3y'' - y' = 0$.

The characteristic equation is $r^4 - 3r^3 + 3r^2 - r = 0$.

By inspection $r^4 - 3r^3 + 3r^2 - r = r(r-1)^3$.

The roots are $r = 0, 1, 1, 1$. ($r = 1$ has multiplicity 3)

The general solution is

$$y = \underbrace{(C_1 + C_2x + C_3x^2)e^x}_{\text{terms coming from } r=1} + \underbrace{C_4}_{\text{from } r=0}.$$

Complex roots come in pairs $r = \alpha \pm i\beta$.

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Example: Solve $y^{(4)} - 4y''' + 8y'' - 8y' + 4y = 0$.

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Example: Solve $y^{(4)} - 4y''' + 8y'' - 8y' + 4y = 0$.

The characteristic equation is $r^4 - 4r^3 + 8r^2 - 8r + 4 = 0$

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Example: Solve $y^{(4)} - 4y''' + 8y'' - 8y' + 4y = 0$.

The characteristic equation is $r^4 - 4r^3 + 8r^2 - 8r + 4 = 0$

$$\Rightarrow (r^2 - 2r + 2)^2 = 0$$

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Example: Solve $y^{(4)} - 4y''' + 8y'' - 8y' + 4y = 0$.

The characteristic equation is $r^4 - 4r^3 + 8r^2 - 8r + 4 = 0$

$$\Rightarrow (r^2 - 2r + 2)^2 = 0 \quad \Rightarrow \quad ((r - 1)^2 + 1)^2 = 0.$$

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Example: Solve $y^{(4)} - 4y''' + 8y'' - 8y' + 4y = 0$.

The characteristic equation is $r^4 - 4r^3 + 8r^2 - 8r + 4 = 0$

$$\Rightarrow (r^2 - 2r + 2)^2 = 0 \quad \Rightarrow \quad ((r - 1)^2 + 1)^2 = 0.$$

Roots are $1 \pm i$, both with multiplicity 2.

Complex roots come in pairs $r = \alpha \pm i\beta$.

If we have such a pair each repeated k times, the corresponding solution is

$$(C_0 + C_1x + \cdots + C_{k-1}x^{k-1})e^{\alpha x} \cos(\beta x) + (D_0 + D_1x + \cdots + D_{k-1}x^{k-1})e^{\alpha x} \sin(\beta x).$$

Example: Solve $y^{(4)} - 4y''' + 8y'' - 8y' + 4y = 0$.

The characteristic equation is $r^4 - 4r^3 + 8r^2 - 8r + 4 = 0$

$$\Rightarrow (r^2 - 2r + 2)^2 = 0 \quad \Rightarrow \quad ((r - 1)^2 + 1)^2 = 0.$$

Roots are $1 \pm i$, both with multiplicity 2.

The general solution is

$$y = (C_1 + C_2x)e^x \cos x + (C_3 + C_4x)e^x \sin x.$$