

19. Nonhomogeneous equations, part 1: undetermined coefficients (Notes on Diffy Qs, 2.5)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p)$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p = 0 + (2x + 1).$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p = 0 + (2x + 1).$$

Why is it the *general* solution? Suppose y_p and \tilde{y}_p be two particular solutions: Then

$$L(y_p - \tilde{y}_p)$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p = 0 + (2x + 1).$$

Why is it the *general* solution? Suppose y_p and \tilde{y}_p be two particular solutions: Then

$$L(y_p - \tilde{y}_p) = Ly_p - L\tilde{y}_p$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p = 0 + (2x + 1).$$

Why is it the *general* solution? Suppose y_p and \tilde{y}_p be two particular solutions: Then

$$L(y_p - \tilde{y}_p) = Ly_p - L\tilde{y}_p = (2x + 1) - (2x + 1) = 0.$$

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p = 0 + (2x + 1).$$

Why is it the *general* solution? Suppose y_p and \tilde{y}_p be two particular solutions: Then

$$L(y_p - \tilde{y}_p) = Ly_p - L\tilde{y}_p = (2x + 1) - (2x + 1) = 0.$$

The difference is a solution to the associated homogeneous equation.

How do we solve nonhomogeneous equations? E.g.,

$$y'' + 5y' + 6y = 2x + 1 \quad (\text{Write } Ly = 2x + 1 \text{ for short}).$$

Consider the *associated homogeneous equation*:

$$y'' + 5y' + 6y = 0 \quad (Ly = 0).$$

Its general solution y_c (i.e. $Ly_c = 0$) is called the *complementary solution*.

Find any one *particular solution* y_p of $Ly_p = 2x + 1$.

$y = y_c + y_p$ is the general solution to $Ly = 2x + 1$ as L is a linear operator:

$$Ly = L(y_c + y_p) = Ly_c + Ly_p = 0 + (2x + 1).$$

Why is it the *general* solution? Suppose y_p and \tilde{y}_p be two particular solutions: Then

$$L(y_p - \tilde{y}_p) = Ly_p - L\tilde{y}_p = (2x + 1) - (2x + 1) = 0.$$

The difference is a solution to the associated homogeneous equation.

Theorem: If y_c is the general solution to $Ly_c = 0$ and y_p is any particular solution to $Ly_p = f(x)$, then

$$y = y_c + y_p \quad \text{is the general solution to } Ly = f(x).$$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p$$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B)$$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B$$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B = 6Ax + (5A + 6B).$$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B = 6Ax + (5A + 6B).$$

We want $6Ax + (5A + 6B) = 2x + 1$ or $6A = 2$ and $5A + 6B = 1$.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B = 6Ax + (5A + 6B).$$

We want $6Ax + (5A + 6B) = 2x + 1$ or $6A = 2$ and $5A + 6B = 1$.

$\Rightarrow A = 1/3$ and $B = -1/9$.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B = 6Ax + (5A + 6B).$$

We want $6Ax + (5A + 6B) = 2x + 1$ or $6A = 2$ and $5A + 6B = 1$.

$\Rightarrow A = 1/3$ and $B = -1/9$.

Let $y_p = \frac{1}{3}x - \frac{1}{9} = \frac{3x - 1}{9}.$

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B = 6Ax + (5A + 6B).$$

We want $6Ax + (5A + 6B) = 2x + 1$ or $6A = 2$ and $5A + 6B = 1$.

$\Rightarrow A = 1/3$ and $B = -1/9$.

Let
$$y_p = \frac{1}{3}x - \frac{1}{9} = \frac{3x - 1}{9}.$$

Solving the complementary problem: $y_c = C_1e^{-2x} + C_2e^{-3x}$.

OK, so how do we find some particular y_p solving $y'' + 5y' + 6y = 2x + 1$.

Undetermined coefficients

We guess in a smart way. We want y_p such that $Ly_p = 2x + 1$. Derivatives of polynomials are polynomials. So are their linear combinations.

Try: $y_p = Ax + B$

$$y_p'' + 5y_p' + 6y_p = (Ax + B)'' + 5(Ax + B)' + 6(Ax + B) = 0 + 5A + 6Ax + 6B = 6Ax + (5A + 6B).$$

We want $6Ax + (5A + 6B) = 2x + 1$ or $6A = 2$ and $5A + 6B = 1$.

$\Rightarrow A = 1/3$ and $B = -1/9$.

Let
$$y_p = \frac{1}{3}x - \frac{1}{9} = \frac{3x - 1}{9}.$$

Solving the complementary problem: $y_c = C_1e^{-2x} + C_2e^{-3x}$.

The general solution is

$$y = C_1e^{-2x} + C_2e^{-3x} + \frac{3x - 1}{9}.$$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}.$$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

Then $0 = y(0)$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

Then $0 = y(0) = C_1 + C_2 - 1/9$,

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

$$\text{Then } 0 = y(0) = C_1 + C_2 - 1/9, \quad 1/3 = y'(0)$$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

$$\text{Then } 0 = y(0) = C_1 + C_2 - 1/9, \quad 1/3 = y'(0) = -2C_1 - 3C_2 + 1/3.$$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

$$\text{Then } 0 = y(0) = C_1 + C_2 - 1/9, \quad 1/3 = y'(0) = -2C_1 - 3C_2 + 1/3.$$

Solve to get $C_1 = 1/3$ and $C_2 = -2/9$.

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

$$\text{Then } 0 = y(0) = C_1 + C_2 - 1/9, \quad 1/3 = y'(0) = -2C_1 - 3C_2 + 1/3.$$

Solve to get $C_1 = 1/3$ and $C_2 = -2/9$.

The solution to the initial value problem is

$$y = \frac{1}{3}e^{-2x} - \frac{2}{9}e^{-3x} + \frac{3x-1}{9} = \frac{3e^{-2x} - 2e^{-3x} + 3x - 1}{9}.$$

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

$$\text{Then } 0 = y(0) = C_1 + C_2 - 1/9, \quad 1/3 = y'(0) = -2C_1 - 3C_2 + 1/3.$$

Solve to get $C_1 = 1/3$ and $C_2 = -2/9$.

The solution to the initial value problem is

$$y = \frac{1}{3}e^{-2x} - \frac{2}{9}e^{-3x} + \frac{3x-1}{9} = \frac{3e^{-2x} - 2e^{-3x} + 3x - 1}{9}.$$

Warning: Do not solve for constants in y_c before adding y_p ! Write $y = y_c + y_p$ and *then* solve for constants.

Suppose we had an initial value problem: $y'' + 5y' + 6y = 2x + 1$, $y(0) = 0$ and $y'(0) = 1/3$.

$$y = C_1 e^{-2x} + C_2 e^{-3x} + \frac{3x-1}{9}. \quad y' = -2C_1 e^{-2x} - 3C_2 e^{-3x} + 1/3.$$

$$\text{Then } 0 = y(0) = C_1 + C_2 - 1/9, \quad 1/3 = y'(0) = -2C_1 - 3C_2 + 1/3.$$

Solve to get $C_1 = 1/3$ and $C_2 = -2/9$.

The solution to the initial value problem is

$$y = \frac{1}{3}e^{-2x} - \frac{2}{9}e^{-3x} + \frac{3x-1}{9} = \frac{3e^{-2x} - 2e^{-3x} + 3x - 1}{9}.$$

Warning: Do not solve for constants in y_c before adding y_p ! Write $y = y_c + y_p$ and *then* solve for constants.

Note: Do not forget lower degree terms even if they don't appear on the right hand side:

E.g., for $Ly = x^3 + 1$, try $y_p = Ax^3 + Bx^2 + Cx + D$.

Sines, cosines and exponentials are similar.

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

$$\underbrace{-4A \cos(2x) - 4B \sin(2x)}_{y_p''} + \underbrace{2(-2A \sin(2x) + 2B \cos(2x))}_{y_p'} + 2 \underbrace{(A \cos(2x) + B \sin(2x))}_{y_p} = \cos(2x),$$

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

$$\underbrace{-4A \cos(2x) - 4B \sin(2x)}_{y_p''} + \underbrace{2(-2A \sin(2x) + 2B \cos(2x))}_{y_p'} + 2 \underbrace{(A \cos(2x) + B \sin(2x))}_{y_p} = \cos(2x),$$

$$\Rightarrow (-4A + 4B + 2A) \cos(2x) + (-4B - 4A + 2B) \sin(2x) = \cos(2x).$$

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

$$\underbrace{-4A \cos(2x) - 4B \sin(2x)}_{y_p''} + \underbrace{2(-2A \sin(2x) + 2B \cos(2x))}_{y_p'} + 2 \underbrace{(A \cos(2x) + B \sin(2x))}_{y_p} = \cos(2x),$$

$$\Rightarrow (-4A + 4B + 2A) \cos(2x) + (-4B - 4A + 2B) \sin(2x) = \cos(2x).$$

$$\Rightarrow -4A + 4B + 2A = 1 \text{ and } -4B - 4A + 2B = 0$$

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

$$\underbrace{-4A \cos(2x) - 4B \sin(2x)}_{y_p''} + \underbrace{2(-2A \sin(2x) + 2B \cos(2x))}_{y_p'} + 2 \underbrace{(A \cos(2x) + B \sin(2x))}_{y_p} = \cos(2x),$$

$$\Rightarrow (-4A + 4B + 2A) \cos(2x) + (-4B - 4A + 2B) \sin(2x) = \cos(2x).$$

$$\Rightarrow -4A + 4B + 2A = 1 \text{ and } -4B - 4A + 2B = 0 \quad \Rightarrow \quad -2A + 4B = 1 \text{ and } 2A + B = 0$$

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

$$\underbrace{-4A \cos(2x) - 4B \sin(2x)}_{y_p''} + 2 \underbrace{(-2A \sin(2x) + 2B \cos(2x))}_{y_p'} + 2 \underbrace{(A \cos(2x) + B \sin(2x))}_{y_p} = \cos(2x),$$

$$\Rightarrow (-4A + 4B + 2A) \cos(2x) + (-4B - 4A + 2B) \sin(2x) = \cos(2x).$$

$$\Rightarrow -4A + 4B + 2A = 1 \text{ and } -4B - 4A + 2B = 0 \quad \Rightarrow \quad -2A + 4B = 1 \text{ and } 2A + B = 0$$

$$\Rightarrow A = -1/10 \text{ and } B = 1/5.$$

Sines, cosines and exponentials are similar.

Example: Find a particular solution to $y'' + 2y' + 2y = \cos(2x)$.

Linear combinations of derivatives of $\cos(2x)$ will be combinations of $\cos(2x)$ and $\sin(2x)$.

Try: $y_p = A \cos(2x) + B \sin(2x)$.

$$\underbrace{-4A \cos(2x) - 4B \sin(2x)}_{y_p''} + 2 \underbrace{(-2A \sin(2x) + 2B \cos(2x))}_{y_p'} + 2 \underbrace{(A \cos(2x) + B \sin(2x))}_{y_p} = \cos(2x),$$

$$\Rightarrow (-4A + 4B + 2A) \cos(2x) + (-4B - 4A + 2B) \sin(2x) = \cos(2x).$$

$$\Rightarrow -4A + 4B + 2A = 1 \text{ and } -4B - 4A + 2B = 0 \Rightarrow -2A + 4B = 1 \text{ and } 2A + B = 0$$

$$\Rightarrow A = -1/10 \text{ and } B = 1/5.$$

$$\text{So } y_p = A \cos(2x) + B \sin(2x) = \frac{-\cos(2x) + 2 \sin(2x)}{10}.$$

Similarly for exponentials:

For $Ly = e^{3x}$, try $y_p = Ae^{3x}$.

Similarly for exponentials:

For $Ly = e^{3x}$, try $y_p = Ae^{3x}$.

And we can combine exponentials, sines, cosines, and polynomials:

Similarly for exponentials:

For $Ly = e^{3x}$, try $y_p = Ae^{3x}$.

And we can combine exponentials, sines, cosines, and polynomials:

For $Ly = (1 + 3x^2)e^{-x} \cos(\pi x)$,

try $y_p = (A + Bx + Cx^2)e^{-x} \cos(\pi x) + (D + Ex + Fx^2)e^{-x} \sin(\pi x)$.

Similarly for exponentials:

For $Ly = e^{3x}$, try $y_p = Ae^{3x}$.

And we can combine exponentials, sines, cosines, and polynomials:

For $Ly = (1 + 3x^2)e^{-x} \cos(\pi x)$,

try $y_p = (A + Bx + Cx^2)e^{-x} \cos(\pi x) + (D + Ex + Fx^2)e^{-x} \sin(\pi x)$.

Plug in, then solve for A, B, C, D, E , and F .

Those guesses may not *always* work.

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x})$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

$$y_p' = Ae^{3x} + 3Axe^{3x} \quad \text{and} \quad y_p'' = 6Ae^{3x} + 9Axe^{3x}$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

$$y_p' = Ae^{3x} + 3Axe^{3x} \quad \text{and} \quad y_p'' = 6Ae^{3x} + 9Axe^{3x}$$

$$y_p'' - 9y_p$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

$$y_p' = Ae^{3x} + 3Axe^{3x} \quad \text{and} \quad y_p'' = 6Ae^{3x} + 9Axe^{3x}$$

$$y_p'' - 9y_p = 6Ae^{3x} + 9Axe^{3x} - 9Axe^{3x}$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

$$y_p' = Ae^{3x} + 3Axe^{3x} \quad \text{and} \quad y_p'' = 6Ae^{3x} + 9Axe^{3x}$$

$$y_p'' - 9y_p = 6Ae^{3x} + 9Axe^{3x} - 9Axe^{3x} = 6Ae^{3x}.$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

$$y_p' = Ae^{3x} + 3Axe^{3x} \quad \text{and} \quad y_p'' = 6Ae^{3x} + 9Axe^{3x}$$

$$y_p'' - 9y_p = 6Ae^{3x} + 9Axe^{3x} - 9Axe^{3x} = 6Ae^{3x}.$$

$$\Rightarrow 6Ae^{3x} = e^{3x} \quad \Rightarrow \quad 6A = 1 \quad \Rightarrow \quad A = 1/6.$$

Those guesses may not *always* work. The guess could get eaten by the left hand side.

Example: Consider $y'' - 9y = e^{3x}$.

Try guessing $y_p = Ae^{3x}$:

$$y_p'' - 9y_p = (9Ae^{3x}) - 9(Ae^{3x}) = 0 \neq e^{3x}.$$

No way to solve for A .

Hint: To save work, compute the complimentary solution first. Here, $y_c = C_1e^{-3x} + C_2e^{3x}$.

Trick: Multiply guess by x : $y_p = Axe^{3x}$.

$$y_p' = Ae^{3x} + 3Axe^{3x} \quad \text{and} \quad y_p'' = 6Ae^{3x} + 9Axe^{3x}$$

$$y_p'' - 9y_p = 6Ae^{3x} + 9Axe^{3x} - 9Axe^{3x} = 6Ae^{3x}.$$

$$\Rightarrow 6Ae^{3x} = e^{3x} \Rightarrow 6A = 1 \Rightarrow A = 1/6.$$

The general solution is

$$y = y_c + y_p = C_1e^{-3x} + C_2e^{3x} + \frac{1}{6}xe^{3x}.$$

Multiplying by x once may not be enough:

Example: Consider $y'' - 6y' + 9y = e^{3x}$.

Multiplying by x once may not be enough:

Example: Consider $y'' - 6y' + 9y = e^{3x}$.

$$y_c = C_1 e^{3x} + C_2 x e^{3x}.$$

Multiplying by x once may not be enough:

Example: Consider $y'' - 6y' + 9y = e^{3x}$.

$$y_c = C_1 e^{3x} + C_2 x e^{3x}.$$

So $y_p = A x e^{3x}$ does not get rid of the duplication with y_c .

Multiplying by x once may not be enough:

Example: Consider $y'' - 6y' + 9y = e^{3x}$.

$$y_c = C_1 e^{3x} + C_2 x e^{3x}.$$

So $y_p = A x e^{3x}$ does not get rid of the duplication with y_c .

Solution?

Multiplying by x once may not be enough:

Example: Consider $y'' - 6y' + 9y = e^{3x}$.

$$y_c = C_1 e^{3x} + C_2 x e^{3x}.$$

So $y_p = A x e^{3x}$ does not get rid of the duplication with y_c .

Solution? Guess $y_p = A x^2 e^{3x}$.

Multiplying by x once may not be enough:

Example: Consider $y'' - 6y' + 9y = e^{3x}$.

$$y_c = C_1 e^{3x} + C_2 x e^{3x}.$$

So $y_p = A x e^{3x}$ does not get rid of the duplication with y_c .

Solution? Guess $y_p = A x^2 e^{3x}$.

So keep multiplying by x until duplication is gone but no more.

What if the right-hand side has several terms, e.g.,

$$Ly = e^{2x} + \cos x.$$

What if the right-hand side has several terms, e.g.,

$$Ly = e^{2x} + \cos x.$$

Find a particular u that solves

$$Lu = e^{2x}.$$

What if the right-hand side has several terms, e.g.,

$$Ly = e^{2x} + \cos x.$$

Find a particular u that solves

$$Lu = e^{2x}.$$

Find particular v that solves

$$Lv = \cos x.$$

What if the right-hand side has several terms, e.g.,

$$Ly = e^{2x} + \cos x.$$

Find a particular u that solves

$$Lu = e^{2x}.$$

Find particular v that solves

$$Lv = \cos x.$$

$y = u + v$ solves $Ly = e^{2x} + \cos x$:

$$Ly = L(u + v) = Lu + Lv = e^{2x} + \cos x.$$