


21. Forced oscillations and resonance,
part 1: Undamped forced motion and resonance
(Notes on Diffy Qs, 2.6)

Jiří Lebl


Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

We return to mass on a spring, but now with nonzero forcing function $F(t)$:



We return to mass on a spring, but now with nonzero forcing function $F(t)$:



The equation is

$$mx'' + cx' + kx = F(t),$$


m is the mass,

c is friction,

k is the spring constant, and

$F(t)$ is an external force acting on the mass.

We return to mass on a spring, but now with nonzero forcing function $F(t)$:



The equation is

$$mx'' + cx' + kx = F(t),$$

m is the mass,

c is friction,


k is the spring constant, and

$F(t)$ is an external force acting on the mass.

We consider periodic forces $F(t)$, and the simplest periodic force is

$$F(t) = F_0 \cos(\omega t)$$

We return to mass on a spring, but now with nonzero forcing function $F(t)$:



The equation is

$$mx'' + cx' + kx = F(t),$$

m is the mass,

c is friction,

k is the spring constant, and

$F(t)$ is an external force acting on the mass.

We consider periodic forces $F(t)$, and the simplest periodic force is

$$F(t) = F_0 \cos(\omega t)$$

Note: Using Fourier series, all periodic functions can be understood via this simple case.

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where } \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency}.$$

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where} \quad \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where } \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

First suppose $\omega_0 \neq \omega$.

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where } \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

First suppose $\omega_0 \neq \omega$.

Using the method of undetermined coefficients we try $x_p = A \cos(\omega t)$ and solve for A .

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where } \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

First suppose $\omega_0 \neq \omega$.

Using the method of undetermined coefficients we try $x_p = A \cos(\omega t)$ and solve for A .
(Including $B \sin(\omega t)$ doesn't hurt, you'll see $B = 0$).

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where } \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

First suppose $\omega_0 \neq \omega$.

Using the method of undetermined coefficients we try $x_p = A \cos(\omega t)$ and solve for A .

(Including $B \sin(\omega t)$ doesn't hurt, you'll see $B = 0$). We find (exercise)

$$x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t).$$

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where} \quad \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

First suppose $\omega_0 \neq \omega$.

Using the method of undetermined coefficients we try $x_p = A \cos(\omega t)$ and solve for A .

(Including $B \sin(\omega t)$ doesn't hurt, you'll see $B = 0$). We find (exercise)

$$x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t).$$

The general solution is

$$x = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t) + \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$$

We start with undamped ($c = 0$) motion:

$$mx'' + kx = F_0 \cos(\omega t).$$

The complementary solution is

$$x_c = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t), \quad \text{where} \quad \omega_0 = \sqrt{k/m} = \text{natural (angular) frequency.}$$

ω_0 is the frequency at which the system “wants to oscillate” without any force F .

First suppose $\omega_0 \neq \omega$.

Using the method of undetermined coefficients we try $x_p = A \cos(\omega t)$ and solve for A .
(Including $B \sin(\omega t)$ doesn't hurt, you'll see $B = 0$). We find (exercise)

$$x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t).$$

The general solution is

$$x = C_1 \cos(\omega_0 t) + C_2 \sin(\omega_0 t) + \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t) \quad \text{or} \\ x = C \cos(\omega_0 t - \gamma) + \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t).$$

A superposition of two phase shifted cosine waves at different frequencies.

Example: Solve $0.5x'' + 8x = 10 \cos(\pi t), \quad x(0) = 0, \quad x'(0) = 0.$

Example: Solve $0.5x'' + 8x = 10 \cos(\pi t), \quad x(0) = 0, \quad x'(0) = 0.$

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$

Example: Solve $0.5x'' + 8x = 10 \cos(\pi t), \quad x(0) = 0, \quad x'(0) = 0.$

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$

The general solution is

$$x = C_1 \cos(4t) + C_2 \sin(4t) + \frac{20}{16 - \pi^2} \cos(\pi t).$$

Example: Solve $0.5x'' + 8x = 10 \cos(\pi t)$, $x(0) = 0$, $x'(0) = 0$.

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$

The general solution is

$$x = C_1 \cos(4t) + C_2 \sin(4t) + \frac{20}{16 - \pi^2} \cos(\pi t).$$

Solve for C_1 and C_2 to find

$$C_1 = \frac{-20}{16 - \pi^2}, \quad C_2 = 0, \quad \text{so}$$

Example: Solve $0.5x'' + 8x = 10 \cos(\pi t)$, $x(0) = 0$, $x'(0) = 0$.

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$


The general solution is

$$x = C_1 \cos(4t) + C_2 \sin(4t) + \frac{20}{16 - \pi^2} \cos(\pi t).$$

Solve for C_1 and C_2 to find

$$C_1 = \frac{-20}{16 - \pi^2}, \quad C_2 = 0, \quad \text{so}$$

$$x = \frac{20}{16 - \pi^2} (\cos(\pi t) - \cos(4t)).$$



Example: Solve $0.5x'' + 8x = 10 \cos(\pi t)$, $x(0) = 0$, $x'(0) = 0$.

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$

The general solution is


$$x = C_1 \cos(4t) + C_2 \sin(4t) + \frac{20}{16 - \pi^2} \cos(\pi t).$$

Solve for C_1 and C_2 to find

$$C_1 = \frac{-20}{16 - \pi^2}, \quad C_2 = 0, \quad \text{so}$$

$$x = \frac{20}{16 - \pi^2} (\cos(\pi t) - \cos(4t)).$$

Notice the “beating” in the figure.



Example: Solve $0.5x'' + 8x = 10 \cos(\pi t)$, $x(0) = 0$, $x'(0) = 0$.

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$

The general solution is

$$x = C_1 \cos(4t) + C_2 \sin(4t) + \frac{20}{16 - \pi^2} \cos(\pi t).$$

Solve for C_1 and C_2 to find


$$C_1 = \frac{-20}{16 - \pi^2}, \quad C_2 = 0, \quad \text{so}$$

$$x = \frac{20}{16 - \pi^2} (\cos(\pi t) - \cos(4t)).$$

Notice the “beating” in the figure.

Using the identity $2 \sin\left(\frac{A-B}{2}\right) \sin\left(\frac{A+B}{2}\right) = \cos B - \cos A$, we find

$$x = \frac{20}{16 - \pi^2} \left(2 \sin\left(\frac{4 - \pi}{2}t\right) \sin\left(\frac{4 + \pi}{2}t\right) \right).$$



Example: Solve $0.5x'' + 8x = 10 \cos(\pi t)$, $x(0) = 0$, $x'(0) = 0$.

$$\omega = \pi, \quad \omega_0 = \sqrt{8/0.5} = 4, \quad F_0 = 10, \quad m = 0.5.$$

The general solution is

$$x = C_1 \cos(4t) + C_2 \sin(4t) + \frac{20}{16 - \pi^2} \cos(\pi t).$$

Solve for C_1 and C_2 to find

$$C_1 = \frac{-20}{16 - \pi^2}, \quad C_2 = 0, \quad \text{so}$$


$$x = \frac{20}{16 - \pi^2} (\cos(\pi t) - \cos(4t)).$$

Notice the “beating” in the figure.

Using the identity $2 \sin\left(\frac{A-B}{2}\right) \sin\left(\frac{A+B}{2}\right) = \cos B - \cos A$, we find

$$x = \frac{20}{16 - \pi^2} \left(2 \sin\left(\frac{4 - \pi}{2}t\right) \sin\left(\frac{4 + \pi}{2}t\right) \right).$$

The solution is a high frequency wave modulated by a low frequency wave.



What about the case $\omega_0 = \omega$?

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

Undetermined coefficients says we should try

$$x_p = At \cos(\omega t) + Bt \sin(\omega t) \quad (\text{the sine term is needed this time}).$$

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

Undetermined coefficients says we should try

$$x_p = At \cos(\omega t) + Bt \sin(\omega t) \quad (\text{the sine term is needed this time}).$$

Write the equation as $x'' + \omega^2 x = \frac{F_0}{m} \cos(\omega t)$ and plug in x_p to find

$$2B\omega \cos(\omega t) - 2A\omega \sin(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

Undetermined coefficients says we should try

$$x_p = At \cos(\omega t) + Bt \sin(\omega t) \quad (\text{the sine term is needed this time}).$$

Write the equation as $x'' + \omega^2 x = \frac{F_0}{m} \cos(\omega t)$ and plug in x_p to find

$$2B\omega \cos(\omega t) - 2A\omega \sin(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

$$A = 0 \text{ and } B = \frac{F_0}{2m\omega}.$$

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

Undetermined coefficients says we should try

$$x_p = At \cos(\omega t) + Bt \sin(\omega t) \quad (\text{the sine term is needed this time}).$$

Write the equation as $x'' + \omega^2 x = \frac{F_0}{m} \cos(\omega t)$ and plug in x_p to find

$$2B\omega \cos(\omega t) - 2A\omega \sin(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

$A = 0$ and $B = \frac{F_0}{2m\omega}$. The particular solution is $\frac{F_0}{2m\omega} t \sin(\omega t)$ and the general

$$x = x_c + x_p = C_1 \cos(\omega t) + C_2 \sin(\omega t) + \frac{F_0}{2m\omega} t \sin(\omega t).$$

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

Undetermined coefficients says we should try

$$x_p = At \cos(\omega t) + Bt \sin(\omega t) \quad (\text{the sine term is needed this time}).$$

Write the equation as $x'' + \omega^2 x = \frac{F_0}{m} \cos(\omega t)$ and plug in x_p to find

$$2B\omega \cos(\omega t) - 2A\omega \sin(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

$A = 0$ and $B = \frac{F_0}{2m\omega}$. The particular solution is $\frac{F_0}{2m\omega} t \sin(\omega t)$ and the general

$$x = x_c + x_p = C_1 \cos(\omega t) + C_2 \sin(\omega t) + \frac{F_0}{2m\omega} t \sin(\omega t).$$

Note the t in the x_p , so x_p grows without bound as $t \rightarrow \infty$.

What about the case $\omega_0 = \omega$?

From $x_p = \frac{F_0}{m(\omega_0^2 - \omega^2)} \cos(\omega t)$, we suspect that something “blows up” when ω approaches ω_0 .

Undetermined coefficients says we should try

$$x_p = At \cos(\omega t) + Bt \sin(\omega t) \quad (\text{the sine term is needed this time}).$$

Write the equation as $x'' + \omega^2 x = \frac{F_0}{m} \cos(\omega t)$ and plug in x_p to find

$$2B\omega \cos(\omega t) - 2A\omega \sin(\omega t) = \frac{F_0}{m} \cos(\omega t).$$

$A = 0$ and $B = \frac{F_0}{2m\omega}$. The particular solution is $\frac{F_0}{2m\omega} t \sin(\omega t)$ and the general

$$x = x_c + x_p = C_1 \cos(\omega t) + C_2 \sin(\omega t) + \frac{F_0}{2m\omega} t \sin(\omega t).$$


Note the t in the x_p , so x_p grows without bound as $t \rightarrow \infty$.

x_p oscillates between $\frac{F_0 t}{2m\omega}$ and $-\frac{F_0 t}{2m\omega}$.

x_c only oscillates between $\pm \sqrt{C_1^2 + C_2^2}$.

For $C_1 = C_2 = 0$, $F_0 = 2$, $m = 1$, $\omega = \pi$,


$$x = \frac{1}{\pi}t \sin(\pi t).$$



For $C_1 = C_2 = 0$, $F_0 = 2$, $m = 1$, $\omega = \pi$,

$$x = \frac{1}{\pi}t \sin(\pi t).$$

So when $\omega = \omega_0$, we produce wild ever growing oscillations. This behavior is called *resonance* or *pure resonance*.




For $C_1 = C_2 = 0$, $F_0 = 2$, $m = 1$, $\omega = \pi$,

$$x = \frac{1}{\pi}t \sin(\pi t).$$

So when $\omega = \omega_0$, we produce wild ever growing oscillations. This behavior is called *resonance* or *pure resonance*.

Resonance can be good:
We can create large oscillations
with small force.



For $C_1 = C_2 = 0$, $F_0 = 2$, $m = 1$, $\omega = \pi$,


$$x = \frac{1}{\pi}t \sin(\pi t).$$

So when $\omega = \omega_0$, we produce wild ever growing oscillations. This behavior is called *resonance* or *pure resonance*.

Resonance can be good:

We can create large oscillations with small force.

Examples: Swinging a child, quartz watches, NMR/MRI, ...



For $C_1 = C_2 = 0, F_0 = 2, m = 1, \omega = \pi$,

$$x = \frac{1}{\pi} t \sin(\pi t).$$

So when $\omega = \omega_0$, we produce wild ever growing oscillations. This behavior is called *resonance* or *pure resonance*.

Resonance can be good:

We can create large oscillations with small force.

Examples: Swinging a child, quartz watches, NMR/MRI, ...

Resonance can be bad:

Examples: Earthquakes, vibrations in engines, soldiers marching on a bridge, ...

