

14. Constant coefficient second order linear ODEs (part 1)

(Notes on Diffy Qs, 2.2)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution.

Consider the problem $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz")

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx}$ $y'' = r^2e^{rx}$

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx} \quad y'' = r^2e^{rx}$

$$\begin{array}{c} y'' - 5y' + 6y = 0, \\ \underbrace{r^2e^{rx}}_{y''} - 5 \underbrace{re^{rx}}_{y'} + 6 \underbrace{e^{rx}}_y = 0, \end{array}$$

Consider the problem $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx}$ $y'' = r^2e^{rx}$

$$\begin{aligned} y'' - 5y' + 6y &= 0, \\ \underbrace{r^2e^{rx}}_{y''} - 5 \underbrace{re^{rx}}_{y'} + 6 \underbrace{e^{rx}}_y &= 0, \\ r^2 - 5r + 6 &= 0 \quad (\text{divide through by } e^{rx}), \end{aligned}$$

Consider the problem $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx}$ $y'' = r^2e^{rx}$

$$\begin{array}{c} y'' - 5y' + 6y = 0, \\ \underbrace{r^2e^{rx}}_{y''} - 5 \underbrace{re^{rx}}_{y'} + 6 \underbrace{e^{rx}}_y = 0, \end{array}$$

$$r^2 - 5r + 6 = 0 \quad (\text{divide through by } e^{rx}),$$

$$(r - 2)(r - 3) = 0.$$

Consider the problem $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx}$ $y'' = r^2e^{rx}$

$$\begin{aligned} y'' - 5y' + 6y &= 0, \\ \underbrace{r^2e^{rx}}_{y''} - 5 \underbrace{re^{rx}}_{y'} + 6 \underbrace{e^{rx}}_y &= 0, \\ r^2 - 5r + 6 &= 0 \quad (\text{divide through by } e^{rx}), \\ (r - 2)(r - 3) &= 0. \end{aligned}$$

So $r = 2$ or $r = 3$.

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx} \quad y'' = r^2e^{rx}$

$$\begin{aligned} y'' - 5y' + 6y &= 0, \\ \underbrace{r^2e^{rx}}_{y''} - 5 \underbrace{re^{rx}}_{y'} + 6 \underbrace{e^{rx}}_y &= 0, \\ r^2 - 5r + 6 &= 0 \quad (\text{divide through by } e^{rx}), \\ (r - 2)(r - 3) &= 0. \end{aligned}$$

So $r = 2$ or $r = 3$. Let $y_1 = e^{2x}$ and $y_2 = e^{3x}$.

Consider the problem $y'' - 5y' + 6y = 0, \quad y(0) = 1, \quad y'(0) = 7.$

It's a second order linear homogeneous equation with constant coefficients.

(*Constant coefficients* means that in $y'' + p(x)y' + q(x)y = f(x)$, the p and q are constants.)

Let's guess a solution. What function stays more or less the same when differentiated?
The exponential!

We try $y = e^{rx}$ (Germans call this "ansatz") $y' = re^{rx} \quad y'' = r^2e^{rx}$

$$\begin{aligned} y'' - 5y' + 6y &= 0, \\ \underbrace{r^2e^{rx}}_{y''} - 5 \underbrace{re^{rx}}_{y'} + 6 \underbrace{e^{rx}}_y &= 0, \\ r^2 - 5r + 6 &= 0 \quad (\text{divide through by } e^{rx}), \\ (r - 2)(r - 3) &= 0. \end{aligned}$$

So $r = 2$ or $r = 3$. Let $y_1 = e^{2x}$ and $y_2 = e^{3x}$.

Exercise: Check that y_1 and y_2 are solutions.

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent:

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C .

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$.

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0)$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0)$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Solve!

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Solve!

$$\text{e.g., } 2 = 2C_1 + 2C_2$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Solve!

$$\text{e.g., } 2 = 2C_1 + 2C_2 \Rightarrow (7 - 2) = (2 - 2)C_1 + (3 - 2)C_2$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Solve!

$$\text{e.g., } 2 = 2C_1 + 2C_2 \Rightarrow (7 - 2) = (2 - 2)C_1 + (3 - 2)C_2 \Rightarrow C_2 = 5$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Solve!

$$\text{e.g., } 2 = 2C_1 + 2C_2 \Rightarrow (7 - 2) = (2 - 2)C_1 + (3 - 2)C_2 \Rightarrow C_2 = 5 \Rightarrow C_1 = -4.$$

Still considering $y'' - 5y' + 6y = 0$, $y(0) = 1$, $y'(0) = 7$.

e^{2x} and e^{3x} are linearly independent: If not, $e^{3x} = Ce^{2x}$ for a constant C . So $e^x = C$. Nonsense!

The general solution is $y = C_1e^{2x} + C_2e^{3x}$ $y' = 2C_1e^{2x} + 3C_2e^{3x}$

$$1 = y(0) = C_1 + C_2,$$

$$7 = y'(0) = 2C_1 + 3C_2.$$

Solve!

$$\text{e.g., } 2 = 2C_1 + 2C_2 \Rightarrow (7 - 2) = (2 - 2)C_1 + (3 - 2)C_2 \Rightarrow C_2 = 5 \Rightarrow C_1 = -4.$$

So the solution is: $y = -4e^{2x} + 5e^{3x}$

Given $ay'' + by' + cy = 0$

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Divide by e^{rx} to get the *characteristic equation* of the ODE:

$$ar^2 + br + c = 0.$$

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Divide by e^{rx} to get the *characteristic equation* of the ODE:

$$ar^2 + br + c = 0.$$

Solve for the r : $r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Divide by e^{rx} to get the *characteristic equation* of the ODE:

$$ar^2 + br + c = 0.$$

Solve for the r : $r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

$\Rightarrow e^{r_1x}$ and e^{r_2x} are solutions (a bit more complicated if $r_1 = r_2$)

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Divide by e^{rx} to get the *characteristic equation* of the ODE:

$$ar^2 + br + c = 0.$$

Solve for the r : $r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$

$\Rightarrow e^{r_1x}$ and e^{r_2x} are solutions (a bit more complicated if $r_1 = r_2$)

Theorem

Suppose that r_1 and r_2 are the roots of the characteristic equation.

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Divide by e^{rx} to get the *characteristic equation* of the ODE:

$$ar^2 + br + c = 0.$$

Solve for the r : $r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

$\Rightarrow e^{r_1x}$ and e^{r_2x} are solutions (a bit more complicated if $r_1 = r_2$)

Theorem

Suppose that r_1 and r_2 are the roots of the characteristic equation.

(i) If r_1 and r_2 are distinct and real (when $b^2 - 4ac > 0$), the general solution is

$$y = C_1e^{r_1x} + C_2e^{r_2x}.$$

Given $ay'' + by' + cy = 0$ try $y = e^{rx}$: $ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$

Divide by e^{rx} to get the *characteristic equation* of the ODE:

$$ar^2 + br + c = 0.$$

Solve for the r : $r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$

$\Rightarrow e^{r_1x}$ and e^{r_2x} are solutions (a bit more complicated if $r_1 = r_2$)

Theorem

Suppose that r_1 and r_2 are the roots of the characteristic equation.

(i) If r_1 and r_2 are distinct and real (when $b^2 - 4ac > 0$), the general solution is

$$y = C_1e^{r_1x} + C_2e^{r_2x}.$$

(ii) If $r_1 = r_2$ (when $b^2 - 4ac = 0$), the general solution is

$$y = (C_1 + C_2x)e^{r_1x}.$$

Example: Solve $y'' - k^2y = 0$.

Example: Solve $y'' - k^2y = 0$.

Characteristic equation is $r^2 - k^2 = 0$ or $(r - k)(r + k) = 0$.

Example: Solve $y'' - k^2y = 0$.

Characteristic equation is $r^2 - k^2 = 0$ or $(r - k)(r + k) = 0$.

$\Rightarrow e^{-kx}$ and e^{kx} are the two linearly independent solutions,

Example: Solve $y'' - k^2y = 0$.

Characteristic equation is $r^2 - k^2 = 0$ or $(r - k)(r + k) = 0$.

$\Rightarrow e^{-kx}$ and e^{kx} are the two linearly independent solutions, and the general solution is

$$y = C_1e^{kx} + C_2e^{-kx}.$$

Example: Solve $y'' - k^2y = 0$.

Characteristic equation is $r^2 - k^2 = 0$ or $(r - k)(r + k) = 0$.

$\Rightarrow e^{-kx}$ and e^{kx} are the two linearly independent solutions, and the general solution is

$$y = C_1e^{kx} + C_2e^{-kx}.$$

Example: Solve $y'' - 8y' + 16y = 0$.

Example: Solve $y'' - k^2y = 0$.

Characteristic equation is $r^2 - k^2 = 0$ or $(r - k)(r + k) = 0$.

$\Rightarrow e^{-kx}$ and e^{kx} are the two linearly independent solutions, and the general solution is

$$y = C_1e^{kx} + C_2e^{-kx}.$$

Example: Solve $y'' - 8y' + 16y = 0$.

Characteristic equation is $r^2 - 8r + 16 = (r - 4)^2 = 0$.

Example: Solve $y'' - k^2y = 0$.

Characteristic equation is $r^2 - k^2 = 0$ or $(r - k)(r + k) = 0$.

$\Rightarrow e^{-kx}$ and e^{kx} are the two linearly independent solutions, and the general solution is

$$y = C_1e^{kx} + C_2e^{-kx}.$$

Example: Solve $y'' - 8y' + 16y = 0$.

Characteristic equation is $r^2 - 8r + 16 = (r - 4)^2 = 0$.

We have a double root $r_1 = r_2 = 4$. The general solution is

$$y = (C_1 + C_2x)e^{4x} = C_1e^{4x} + C_2xe^{4x}.$$

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

Why does xe^{rx} work if we the root is doubled?

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

Why does xe^{rx} work if we the root is doubled?

Think of two distinct roots r_1 and r_2 very close.

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

Why does xe^{rx} work if we the root is doubled?

Think of two distinct roots r_1 and r_2 very close.

$$\frac{e^{r_2x} - e^{r_1x}}{r_2 - r_1} \quad \text{is a solution.}$$

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

Why does xe^{rx} work if we the root is doubled?

Think of two distinct roots r_1 and r_2 very close.

$$\frac{e^{r_2x} - e^{r_1x}}{r_2 - r_1} \quad \text{is a solution.}$$

Doubled root is like taking the limit $r_1 \rightarrow r_2$.

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

Why does xe^{rx} work if the root is doubled?

Think of two distinct roots r_1 and r_2 very close.

$$\frac{e^{r_2x} - e^{r_1x}}{r_2 - r_1} \quad \text{is a solution.}$$

Doubled root is like taking the limit $r_1 \rightarrow r_2$.

So we are taking the derivative of e^{rx} as if r is the variable:

$$\frac{d}{dr} [e^{rx}] = xe^{rx}$$

In a sense, a doubled root rarely happens for a randomly chosen equation, but it does occur in real-world problems (e.g., critically damped mass-spring system).

Why does xe^{rx} work if the root is doubled?

Think of two distinct roots r_1 and r_2 very close.

$$\frac{e^{r_2x} - e^{r_1x}}{r_2 - r_1} \quad \text{is a solution.}$$

Doubled root is like taking the limit $r_1 \rightarrow r_2$.

So we are taking the derivative of e^{rx} as if r is the variable:

$$\frac{d}{dr} [e^{rx}] = xe^{rx}$$

Voila!