

11. Exact equations (part 1)

(Notes on Diffy Qs, 1.8)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

An *exact equation* is an equation that encodes

$$F(x, y) = \text{constant}$$

for a *potential function* $F(x, y)$.

An *exact equation* is an equation that encodes

$$F(x, y) = \text{constant}$$

for a *potential function* $F(x, y)$.

Naming suggests electric potential or potential energy.

An *exact equation* is an equation that encodes


$$F(x, y) = \text{constant}$$

for a *potential function* $F(x, y)$.

Naming suggests electric potential or potential energy.

Such equations come up when there is some conservation law at play.
(e.g., conservation of energy)

Example: Let $F(x, y) = x^2 + y^2$




Solutions to $F(x, y) = x^2 + y^2 = C$.

Example: Let $F(x, y) = x^2 + y^2$

Take the *total derivative* of F :

$$dF = \frac{\partial F}{\partial x}dx + \frac{\partial F}{\partial y}dy = F_x dx + F_y dy.$$




Solutions to $F(x, y) = x^2 + y^2 = C$.

Example: Let $F(x, y) = x^2 + y^2$

Take the *total derivative* of F :

$$dF = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy = F_x dx + F_y dy.$$

So $dF = 2x dx + 2y dy$.



Solutions to $F(x, y) = x^2 + y^2 = C$.

Example: Let $F(x, y) = x^2 + y^2$


Take the *total derivative* of F :

$$dF = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy = F_x dx + F_y dy.$$

So $dF = 2x dx + 2y dy$.

The differential equation for $F(x, y) = C$ is

$$dF = 0.$$



Solutions to $F(x, y) = x^2 + y^2 = C$.

Example: Let $F(x, y) = x^2 + y^2$

Take the *total derivative* of F :

$$dF = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy = F_x dx + F_y dy.$$


So $dF = 2x dx + 2y dy$.

The differential equation for $F(x, y) = C$ is

$$dF = 0.$$

In this case,

$$2x dx + 2y dy = 0 \quad \text{or} \quad 2x + 2y \frac{dy}{dx} = 0$$



Solutions to $F(x, y) = x^2 + y^2 = C$.

In general, $M dx + N dy = 0$, or $M + N \frac{dy}{dx} = 0$ is exact if it is $dF = 0$.

In general, $M dx + N dy = 0$, or $M + N \frac{dy}{dx} = 0$ is exact if it is $dF = 0$.

In other words, $M dx + N dy = 0$ is exact if there is an F such that $M = F_x$ and $N = F_y$.

In general, $M dx + N dy = 0$, or $M + N \frac{dy}{dx} = 0$ is exact if it is $dF = 0$.

In other words, $M dx + N dy = 0$ is exact if there is an F such that $M = F_x$ and $N = F_y$.

So $2x dx + 2y dy = 0$ is exact.

In general, $M dx + N dy = 0$, or $M + N \frac{dy}{dx} = 0$ is exact if it is $dF = 0$.

In other words, $M dx + N dy = 0$ is exact if there is an F such that $M = F_x$ and $N = F_y$.

So $2x dx + 2y dy = 0$ is exact.

$x^2 + y^2 = C$ are implicit solutions.

In general, $M dx + N dy = 0$, or $M + N \frac{dy}{dx} = 0$ is exact if it is $dF = 0$.

In other words, $M dx + N dy = 0$ is exact if there is an F such that $M = F_x$ and $N = F_y$.

So $2x dx + 2y dy = 0$ is exact.

$x^2 + y^2 = C$ are implicit solutions.

$y = \pm\sqrt{C - x^2}$ are the explicit solutions.

In general, $M dx + N dy = 0$, or $M + N \frac{dy}{dx} = 0$ is exact if it is $dF = 0$.

In other words, $M dx + N dy = 0$ is exact if there is an F such that $M = F_x$ and $N = F_y$.

So $2x dx + 2y dy = 0$ is exact.

$x^2 + y^2 = C$ are implicit solutions.

$y = \pm \sqrt{C - x^2}$ are the explicit solutions.

Normally, we start with $M dx + N dy = 0$ and we wish to find the unknown $F(x, y)$.

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

\vec{v} is a *conservative vector field* if it comes with a potential function F , where $\vec{v} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y} \right)$.

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

\vec{v} is a *conservative vector field* if it comes with a potential function F , where $\vec{v} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y} \right)$.

Let γ be a path in the plane from (x_1, y_1) and ending at (x_2, y_2) and think of \vec{v} as force.

The work required to move along γ is
$$\int_{\gamma} \vec{v}(\vec{r}) \cdot d\vec{r} = \int_{\gamma} M dx + N dy = F(x_2, y_2) - F(x_1, y_1).$$

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

\vec{v} is a *conservative vector field* if it comes with a potential function F , where $\vec{v} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y} \right)$.

Let γ be a path in the plane from (x_1, y_1) and ending at (x_2, y_2) and think of \vec{v} as force.

The work required to move along γ is
$$\int_{\gamma} \vec{v}(\vec{r}) \cdot d\vec{r} = \int_{\gamma} M dx + N dy = F(x_2, y_2) - F(x_1, y_1).$$

In other words, work done depends only on endpoints.

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

\vec{v} is a *conservative vector field* if it comes with a potential function F , where $\vec{v} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y} \right)$.

Let γ be a path in the plane from (x_1, y_1) and ending at (x_2, y_2) and think of \vec{v} as force.

The work required to move along γ is
$$\int_{\gamma} \vec{v}(\vec{r}) \cdot d\vec{r} = \int_{\gamma} M dx + N dy = F(x_2, y_2) - F(x_1, y_1).$$

In other words, work done depends only on endpoints.

Example: Force of gravity is conservative (assuming 2D) — F is the gravitational potential.

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

\vec{v} is a *conservative vector field* if it comes with a potential function F , where $\vec{v} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y} \right)$.

Let γ be a path in the plane from (x_1, y_1) and ending at (x_2, y_2) and think of \vec{v} as force.

The work required to move along γ is
$$\int_{\gamma} \vec{v}(\vec{r}) \cdot d\vec{r} = \int_{\gamma} M dx + N dy = F(x_2, y_2) - F(x_1, y_1).$$

In other words, work done depends only on endpoints.

Example: Force of gravity is conservative (assuming 2D) — F is the gravitational potential. Work done by moving a mass only depends on the change in elevation.

Interpretation: At each point (x, y) there is a vector $\vec{v} = (M, N)$ giving a *vector field*.

\vec{v} is a *conservative vector field* if it comes with a potential function F , where $\vec{v} = \left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y} \right)$.

Let γ be a path in the plane from (x_1, y_1) and ending at (x_2, y_2) and think of \vec{v} as force.

The work required to move along γ is
$$\int_{\gamma} \vec{v}(\vec{r}) \cdot d\vec{r} = \int_{\gamma} M dx + N dy = F(x_2, y_2) - F(x_1, y_1).$$

In other words, work done depends only on endpoints.

Example: Force of gravity is conservative (assuming 2D) — F is the gravitational potential. Work done by moving a mass only depends on the change in elevation.

Exact equations are conservative vector fields, and their implicit solutions are given by the potential functions.

The process is:

1) We are given an equation $M + N \frac{dy}{dx} = 0$

The process is:

- 1) We are given an equation $M + N \frac{dy}{dx} = 0$
- 2) We determine if the equation is exact.

The process is:

- 1) We are given an equation $M + N \frac{dy}{dx} = 0$
- 2) We determine if the equation is exact.
- 3) If it is, we try to find an F such that $F_x = M$ and $F_y = N$.

The process is:

1) We are given an equation $M + N \frac{dy}{dx} = 0$

2) We determine if the equation is exact.

3) If it is, we try to find an F such that $F_x = M$ and $F_y = N$.

$F(x, y(x)) = C$ gives an implicit solution.

The process is:

1) We are given an equation $M + N \frac{dy}{dx} = 0$

2) We determine if the equation is exact.

3) If it is, we try to find an F such that $F_x = M$ and $F_y = N$.

$F(x, y(x)) = C$ gives an implicit solution.

Note: Adding a constant to F does not change anything.

The process is:

1) We are given an equation $M + N \frac{dy}{dx} = 0$

2) We determine if the equation is exact.

3) If it is, we try to find an F such that $F_x = M$ and $F_y = N$.

$F(x, y(x)) = C$ gives an implicit solution.

Note: Adding a constant to F does not change anything.

If $F(x, y)$ works, $F(x, y) + 3$ or $F(x, y) - 8$ also work.

Example: Consider $2x + 2y\frac{dy}{dx} = 0$. Forget we knew what F was.

Example: Consider $2x + 2y \frac{dy}{dx} = 0$. Forget we knew what F was.

$M = 2x$ and $N = 2y$.

Example: Consider $2x + 2y \frac{dy}{dx} = 0$. Forget we knew what F was.

$M = 2x$ and $N = 2y$.

If F exists, $F_x(x, y) = 2x$.

Example: Consider $2x + 2y\frac{dy}{dx} = 0$. Forget we knew what F was.

$M = 2x$ and $N = 2y$.

If F exists, $F_x(x, y) = 2x$.

Integrate in x : $F(x, y) = x^2 + A(y)$, for some function $A(y)$ (constant of integration).

Example: Consider $2x + 2y \frac{dy}{dx} = 0$. Forget we knew what F was.

$M = 2x$ and $N = 2y$.

If F exists, $F_x(x, y) = 2x$.

Integrate in x : $F(x, y) = x^2 + A(y)$, for some function $A(y)$ (constant of integration).

Differentiate in y and set it equal to N :

$$2y = F_y(x, y) = A'(y).$$

Example: Consider $2x + 2y \frac{dy}{dx} = 0$. Forget we knew what F was.

$M = 2x$ and $N = 2y$.

If F exists, $F_x(x, y) = 2x$.

Integrate in x : $F(x, y) = x^2 + A(y)$, for some function $A(y)$ (constant of integration).

Differentiate in y and set it equal to N :

$$2y = F_y(x, y) = A'(y).$$

Integrate $2y = A'(y)$: $A(y) = y^2$ (no need for a constant of integration).

Example: Consider $2x + 2y \frac{dy}{dx} = 0$. Forget we knew what F was.

$M = 2x$ and $N = 2y$.

If F exists, $F_x(x, y) = 2x$.

Integrate in x : $F(x, y) = x^2 + A(y)$, for some function $A(y)$ (constant of integration).

Differentiate in y and set it equal to N :

$$2y = F_y(x, y) = A'(y).$$

Integrate $2y = A'(y)$: $A(y) = y^2$ (no need for a constant of integration).

So

$$F(x, y) = x^2 + A(y) = x^2 + y^2.$$

The procedure (if equation is exact):

The procedure (if equation is exact):

- (i) Integrate $F_x = M$ in x resulting in $F(x, y) = \text{something} + A(y)$.

The procedure (if equation is exact):

- (i) Integrate $F_x = M$ in x resulting in $F(x, y) = \text{something} + A(y)$.
- (ii) Differentiate F in y , and set that equal to N to find $A(y)$ by integration.

The procedure (if equation is exact):

- (i) Integrate $F_x = M$ in x resulting in $F(x, y) = \text{something} + A(y)$.
- (ii) Differentiate F in y , and set that equal to N to find $A(y)$ by integration.

Roles of x and y (and so M and N) can be reversed.

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

$$F_x(x, y) = 2x + y.$$

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

$$F_x(x, y) = 2x + y.$$

$$\text{Integrate in } x: \quad F(x, y) = x^2 + xy + A(y)$$

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

$$F_x(x, y) = 2x + y.$$

$$\text{Integrate in } x: \quad F(x, y) = x^2 + xy + A(y)$$

Differentiate in y and set equal to N :

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

$$F_x(x, y) = 2x + y.$$

$$\text{Integrate in } x: \quad F(x, y) = x^2 + xy + A(y)$$

Differentiate in y and set equal to N :

$$N = xy = F_y(x, y) = x + A'(y).$$

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

$$F_x(x, y) = 2x + y.$$

$$\text{Integrate in } x: \quad F(x, y) = x^2 + xy + A(y)$$

Differentiate in y and set equal to N :

$$N = xy = F_y(x, y) = x + A'(y).$$

But there is no way to write $xy = x + A'(y)$, no matter what $A'(y)$ is.

Example: Consider $2x + y + xy \frac{dy}{dx} = 0$.

$$M = 2x + y, \quad N = xy.$$

We try as before. Suppose F exists.

$$F_x(x, y) = 2x + y.$$

$$\text{Integrate in } x: \quad F(x, y) = x^2 + xy + A(y)$$

Differentiate in y and set equal to N :

$$N = xy = F_y(x, y) = x + A'(y).$$

But there is no way to write $xy = x + A'(y)$, no matter what $A'(y)$ is.

The equation is not exact! F does not exist.

How to check if F exists?

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y}$$

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x}$$

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y}$$

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}.$$

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}.$$

In fact:

Theorem (Poincaré)

If M and N are continuously differentiable functions of (x, y) , and $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then near any point there is a function $F(x, y)$ such that $M = \frac{\partial F}{\partial x}$ and $N = \frac{\partial F}{\partial y}$.

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}.$$

In fact:

Theorem (Poincaré)

If M and N are continuously differentiable functions of (x, y) , and $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then near any point there is a function $F(x, y)$ such that $M = \frac{\partial F}{\partial x}$ and $N = \frac{\partial F}{\partial y}$.

Note: The theorem doesn't say a *global* F exists, only *local*.

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}.$$

In fact:

Theorem (Poincaré)

If M and N are continuously differentiable functions of (x, y) , and $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then near any point there is a function $F(x, y)$ such that $M = \frac{\partial F}{\partial x}$ and $N = \frac{\partial F}{\partial y}$.

Note: The theorem doesn't say a *global* F exists, only *local*.

Back to example: If $M = 2x + y$ and $N = xy$, then $M_y = 1$ and $N_x = y$.

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}.$$

In fact:

Theorem (Poincaré)

If M and N are continuously differentiable functions of (x, y) , and $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then near any point there is a function $F(x, y)$ such that $M = \frac{\partial F}{\partial x}$ and $N = \frac{\partial F}{\partial y}$.

Note: The theorem doesn't say a *global* F exists, only *local*.

Back to example: If $M = 2x + y$ and $N = xy$, then $M_y = 1$ and $N_x = y$. Not equal.

How to check if F exists?

If F exists, then $M = F_x$ and $N = F_y$.

Then

$$\frac{\partial M}{\partial y} = \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}.$$

In fact:

Theorem (Poincaré)

If M and N are continuously differentiable functions of (x, y) , and $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then near any point there is a function $F(x, y)$ such that $M = \frac{\partial F}{\partial x}$ and $N = \frac{\partial F}{\partial y}$.

Note: The theorem doesn't say a *global* F exists, only *local*.

Back to example: If $M = 2x + y$ and $N = xy$, then $M_y = 1$ and $N_x = y$. Not equal. Not exact.

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

$M_y = 1 = N_x$ so the equation is exact.

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works.

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0$, so $M = 2x + y$ and $N = x - 1$.

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works. Take $F(x, y) = x^2 + xy - y$.

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works. Take $F(x, y) = x^2 + xy - y.$

Implicit (general) solution is $x^2 + xy - y = C$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0,$ so $M = 2x + y$ and $N = x - 1.$

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works. Take $F(x, y) = x^2 + xy - y.$

Implicit (general) solution is $x^2 + xy - y = C$

$y(0) = 1 \Rightarrow F(0, 1) = C$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0$, so $M = 2x + y$ and $N = x - 1$.

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works. Take $F(x, y) = x^2 + xy - y$.

Implicit (general) solution is $x^2 + xy - y = C$

$y(0) = 1 \Rightarrow F(0, 1) = C \Rightarrow 0^2 + 0 \times 1 - 1 = C$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0$, so $M = 2x + y$ and $N = x - 1$.

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works. Take $F(x, y) = x^2 + xy - y$.

Implicit (general) solution is $x^2 + xy - y = C$

$y(0) = 1 \Rightarrow F(0, 1) = C \Rightarrow 0^2 + 0 \times 1 - 1 = C \Rightarrow C = -1.$

Example: Solve $\frac{dy}{dx} = \frac{-2x - y}{x - 1}, \quad y(0) = 1.$

Write the equation as $(2x + y) + (x - 1)\frac{dy}{dx} = 0$, so $M = 2x + y$ and $N = x - 1$.

$M_y = 1 = N_x$ so the equation is exact.

Integrate M in x : $F(x, y) = x^2 + xy + A(y)$

Differentiate in y and set to N to find $x - 1 = x + A'(y)$

$A'(y) = -1$ so $A(y) = -y$ works. Take $F(x, y) = x^2 + xy - y$.

Implicit (general) solution is $x^2 + xy - y = C$

$y(0) = 1 \Rightarrow F(0, 1) = C \Rightarrow 0^2 + 0 \times 1 - 1 = C \Rightarrow C = -1.$

Finally, solve $x^2 + xy - y = -1$ for y :

$$y = \frac{-x^2 - 1}{x - 1}.$$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0)$$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy$$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy$$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi$$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

The theorem only guaranteed F defined *locally*.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

The theorem only guaranteed F defined *locally*.

Suppose $x > 0$ (note the IC).

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

The theorem only guaranteed F defined *locally*.

Suppose $x > 0$ (note the IC).

Integrate M in x to find $F(x, y) = \arctan(y/x)$.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

The theorem only guaranteed F defined *locally*.

Suppose $x > 0$ (note the IC).

Integrate M in x to find $F(x, y) = \arctan(y/x)$.

Implicit solution: $\arctan(y/x) = C$.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0, \quad y(1) = 2.$ (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.

If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

The theorem only guaranteed F defined *locally*.

Suppose $x > 0$ (note the IC).

Integrate M in x to find $F(x, y) = \arctan(y/x)$.

Implicit solution: $\arctan(y/x) = C$.

Explicit solution: $y = \tan(C)x$.

Example: Solve $\frac{-y}{x^2 + y^2}dx + \frac{x}{x^2 + y^2}dy = 0$, $y(1) = 2$. (Exercise: $M_y = N_x$)

The vector field (M, N) is **not** conservative on the entire plane minus the origin.

Problem: Let γ be a circle around the origin counterclockwise starting and ending at $(1, 0)$.
If F existed we would have (computation left to interested student)

$$0 = F(1, 0) - F(1, 0) = \int_{\gamma} F_x dx + F_y dy = \int_{\gamma} \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy = 2\pi \quad \text{Nonsense!}$$

There is no F defined for all $(x, y) \neq (0, 0)$.

The theorem only guaranteed F defined *locally*.

Suppose $x > 0$ (note the IC).

Integrate M in x to find $F(x, y) = \arctan(y/x)$.

Implicit solution: $\arctan(y/x) = C$.

Explicit solution: $y = \tan(C)x$.

As $y(1) = 2$, we get $\tan(C) = 2$,

so $y = 2x$ is the solution (only for $x > 0$).

