

## 9. Autonomous equations (Notes on Diffy Qs, 1.6)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.


$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.

**Example:** The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

$x$ =temperature of coffee,  $t$ =time,

$A$ =ambient temperature,  $k$ =constant



$$x' = 0.3(5 - x) \quad (\text{i.e., } k = 0.3, A = 5)$$

$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.


**Example:** The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

$x$ =temperature of coffee,  $t$ =time,

$A$ =ambient temperature,  $k$ =constant

The solution  $x = A$  is called  
an *equilibrium solution*.



$$x' = 0.3(5 - x) \quad (\text{i.e., } k = 0.3, A = 5)$$

$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.

**Example:** The cooling coffee problem


$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

$x$ =temperature of coffee,  $t$ =time,

$A$ =ambient temperature,  $k$ =constant

The solution  $x = A$  is called  
an *equilibrium solution*.

Equilibrium solutions happen when  $f(x) = 0$ .



$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.

**Example:** The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$


$x$ =temperature of coffee,  $t$ =time,

$A$ =ambient temperature,  $k$ =constant

The solution  $x = A$  is called  
an *equilibrium solution*.

Equilibrium solutions happen when  $f(x) = 0$ .

Call an  $x$  such that  $f(x) = 0$  a *critical point*  
( $f(5) = 0$  in the plot)



$$x' = 0.3(5 - x) \quad (\text{i.e., } k = 0.3, A = 5)$$

$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.

**Example:** The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

$x$ =temperature of coffee,  $t$ =time,


$A$ =ambient temperature,  $k$ =constant

The solution  $x = A$  is called  
an *equilibrium solution*.

Equilibrium solutions happen when  $f(x) = 0$ .

Call an  $x$  such that  $f(x) = 0$  a *critical point*  
( $f(5) = 0$  in the plot)

The critical point  $x = A = 5$  is *stable*,  
small changes in initial condition still lead to the solution approaching  $x = 5$  as  $t \rightarrow \infty$ .



$\frac{dx}{dt} = f(x)$  (equation independent of time) is called an *autonomous equation*.

**Example:** The cooling coffee problem

$$\frac{dx}{dt} = k(A - x) \quad (\text{Newton's law of cooling})$$

$x$ =temperature of coffee,  $t$ =time,

$A$ =ambient temperature,  $k$ =constant


The solution  $x = A$  is called  
an *equilibrium solution*.

Equilibrium solutions happen when  $f(x) = 0$ .

Call an  $x$  such that  $f(x) = 0$  a *critical point*  
( $f(5) = 0$  in the plot)

The critical point  $x = A = 5$  is *stable*,  
small changes in initial condition still lead to the solution approaching  $x = 5$  as  $t \rightarrow \infty$ .

A critical point that is not stable is called *unstable*.



Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .


Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

**Example:**  $x' = 0.1x(5 - x)$




Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .




Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .  
 $x = 0$  is unstable,




Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .  
 $x = 0$  is unstable,  $x = 5$  is stable.



Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .


Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .  
 $x = 0$  is unstable,  $x = 5$  is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$



Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )


**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .  
 $x = 0$  is unstable,  $x = 5$  is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$

**Note:** Picture not enough to decide for  $x(0) < 0$ .



Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .

Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )


**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .  
 $x = 0$  is unstable,  $x = 5$  is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$

**Note:** Picture not enough to decide for  $x(0) < 0$ .  $x(t)$  may not exist for all time  $t$ .



Consider the *logistic equation*:  $x' = kx(M - x)$ , for some  $k > 0$  and  $M > 0$ .


Common model for population where there is a maximum sustainable population  $M$ .  
(less catastrophic than  $x' = kx$ )

**Example:**  $x' = 0.1x(5 - x)$

Two critical points  $x = 0$  and  $x = 5$ .  
 $x = 0$  is unstable,  $x = 5$  is stable.

By inspection,

$$\lim_{t \rightarrow \infty} x(t) = \begin{cases} 5 & \text{if } x(0) > 0, \\ 0 & \text{if } x(0) = 0, \\ \text{DNE or } -\infty & \text{if } x(0) < 0. \end{cases}$$



**Note:** Picture not enough to decide for  $x(0) < 0$ .  $x(t)$  may not exist for all time  $t$ .

So to find long term behavior,  $x(t)$  for very large  $t$ , we don't need to solve.

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

1) Draw the  $x$  axis (vertical).

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):


- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$



To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):


- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$



$x = 5$


$x = 0$



To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$




Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$




Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .

Graph  $f(x) = 0.1x(5 - x)$ ,

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$




Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .  
Graph  $f(x) = 0.1x(5 - x)$ , or plug in a few points,

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$



Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .


Graph  $f(x) = 0.1x(5 - x)$ , or plug in a few points,

e.g.,  $f(6) = -0.6 < 0$ ,

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$



Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .


Graph  $f(x) = 0.1x(5 - x)$ , or plug in a few points,

e.g.,  $f(6) = -0.6 < 0$ ,  $f(1) = 0.4 > 0$ ,

To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$




Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .

Graph  $f(x) = 0.1x(5 - x)$ , or plug in a few points,  
e.g.,  $f(6) = -0.6 < 0$ ,  $f(1) = 0.4 > 0$ ,  $f(-1) = -0.6 < 0$


To find  $\lim_{t \rightarrow \infty} x(t)$ , enough to look at a *phase diagram* (or *phase portrait*):

- 1) Draw the  $x$  axis (vertical).
- 2) Mark the critical points.
- 3) Draw an up arrow between where  $f(x) > 0$  and down arrow where  $f(x) < 0$ .

E.g.:  $x' = f(x) = 0.1x(5 - x)$




Enough to note that  $x = 0$  and  $x = 5$  solve  $0.1x(5 - x) = 0$ .  
Graph  $f(x) = 0.1x(5 - x)$ , or plug in a few points,  
e.g.,  $f(6) = -0.6 < 0$ ,  $f(1) = 0.4 > 0$ ,  $f(-1) = -0.6 < 0$




Graph of  $y = f(x) = 0.1x(5 - x)$

Armed with a phase diagram, easy to sketch solutions.

Classifying critical points is also easy:




Classifying critical points is also easy:



**Remark:** If one arrow points in and one out, sometimes called *semistable*.


Classifying critical points is also easy:



**Remark:** If one arrow points in and one out, sometimes called *semistable*.

Unstable points tend to be bad news:

Classifying critical points is also easy:



**Remark:** If one arrow points in and one out, sometimes called *semistable*.

Unstable points tend to be bad news:

Small changes in initial conditions lead to different outcomes.

**Example:** Consider the logistic equation with harvesting.

**Example:** Consider the logistic equation with harvesting.  
An alien race really likes to eat humans,

**Example:** Consider the logistic equation with harvesting.  
An alien race really likes to eat humans,  
they harvest  $h$  million humans per year.

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,  
they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,  
they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

$M$ =limiting population when no harvesting.

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,  
they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

$M$ =limiting population when no harvesting.

$k > 0$  is a number depending on how quickly humans multiply.

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,  
they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

$M$ =limiting population when no harvesting.

$k > 0$  is a number depending on how quickly humans multiply.

Equation becomes:  $x' = kx(M - x) - h$ .

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,  
they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

$M$ =limiting population when no harvesting.

$k > 0$  is a number depending on how quickly humans multiply.

Equation becomes:  $x' = kx(M - x) - h$ .

Find critical points (solve  $kx(M - x) - h = -kx^2 + kMx - h = 0$ )

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

$M$ =limiting population when no harvesting.

$k > 0$  is a number depending on how quickly humans multiply.

Equation becomes:  $x' = kx(M - x) - h$ .

Find critical points (solve  $kx(M - x) - h = -kx^2 + kMx - h = 0$ )

There are two (quadratic formula). Give them names:

$$A = \frac{kM + \sqrt{(kM)^2 - 4hk}}{2k}, \quad B = \frac{kM - \sqrt{(kM)^2 - 4hk}}{2k}.$$

**Example:** Consider the logistic equation with harvesting.

An alien race really likes to eat humans,

they harvest  $h$  million humans per year.

$x$ =millions of humans on the planet,  $t$ =time in years,

$M$ =limiting population when no harvesting.

$k > 0$  is a number depending on how quickly humans multiply.

Equation becomes:  $x' = kx(M - x) - h$ .

Find critical points (solve  $kx(M - x) - h = -kx^2 + kMx - h = 0$ )

There are two (quadratic formula). Give them names:


$$A = \frac{kM + \sqrt{(kM)^2 - 4hk}}{2k}, \quad B = \frac{kM - \sqrt{(kM)^2 - 4hk}}{2k}.$$

3 possibilities:  $A > B$ , or  $A = B$ , or  $A$  and  $B$  both complex (i.e. no real solutions).

For example, let  $M = 8$  and  $k = 0.1$ .

For example, let  $M = 8$  and  $k = 0.1$ .


Harvest  $h = 1$  million:



$$x' = 0.1x(8-x) - 1$$

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:




$$x' = 0.1x(8-x) - 1$$

$$A \approx 6.45,$$

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$ , stable

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$ , stable       $B \approx 1.55$ ,

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:




$$x' = 0.1x(8 - x) - 1$$

$A \approx 6.45$ , stable

$B \approx 1.55$ , unstable

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:




$$x' = 0.1x(8 - x) - 1$$

$A \approx 6.45$ , stable       $B \approx 1.55$ , unstable

As long as there are at least 1.55 million humans, the harvesting is sustainable.

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:



$$x' = 0.1x(8 - x) - 1$$


$A \approx 6.45$ , stable       $B \approx 1.55$ , unstable

As long as there are at least 1.55 million humans, the harvesting is sustainable.

If there is an earthquake and population drops below  $B$ ,

For example, let  $M = 8$  and  $k = 0.1$ .

Harvest  $h = 1$  million:




$$x' = 0.1x(8-x) - 1$$

$A \approx 6.45$ , stable       $B \approx 1.55$ , unstable

As long as there are at least 1.55 million humans, the harvesting is sustainable.


If there is an earthquake and population drops below  $B$ ,  
the alien race becomes vegetarian. (not good for us either)

Now harvest  $h = 1.6$ :



$$x' = 0.1x(8-x) - 1.6.$$


Now harvest  $h = 1.6$ :



$$x' = 0.1x(8 - x) - 1.6.$$

$A = B = 4$ , only one (unstable) critical point.

Now harvest  $h = 1.6$ :




$$x' = 0.1x(8 - x) - 1.6.$$

$A = B = 4$ , only one (unstable) critical point.

If population starts above 4 million, it will tend towards 4 million.

Now harvest  $h = 1.6$ :



$$x' = 0.1x(8 - x) - 1.6.$$


$A = B = 4$ , only one (unstable) critical point.

If population starts above 4 million, it will tend towards 4 million.

If the population drops below 4 million, humans die out.


Now harvest  $h = 2$  million per year.

Now harvest  $h = 2$  million per year.



$$x' = 0.1x(8-x) - 2.$$


Now harvest  $h = 2$  million per year.



$$x' = 0.1x(8-x) - 2.$$

No critical points.

Now harvest  $h = 2$  million per year.



$$x' = 0.1x(8-x) - 2.$$

No critical points. Population always goes to 0.